

TRANSACTIONS
OF THE
LINNAEAN SOCIETY
OF
NEW YORK
Volume VII

The Species of
Middle American Birds

A list of all species recorded
from Mexico to Panama, with
suggested English names, outlines of range,
and a distributional bibliography.

By Eugene Eisenmann

With the collaboration in the selection
of English names of Emmet R. Blake
and Edward L. Chalif

New York

APRIL, 1955

17

QL

687

E 36

THE LINNAEAN SOCIETY
OF NEW YORK

Regular meetings of the Society are held on the second and fourth Tuesdays of each month from October to May. Informal meetings are held at least once a month during June to September, inclusive. All meetings (except the Annual dinner meeting on the second Tuesday of March) are held at the American Museum of Natural History, and are open to the public.

231849

Persons interested in natural history are eligible to membership in the Society. Annual dues for Active Members are \$4.00, and for Associate Members (residing more than fifty miles from New York City) are \$1.50.

The Society conducts field trips and maintains a small circulating library for its members. It distributes free to all members a monthly *News-Letter*, and every two or three years an issue of *Proceedings* containing articles and notes of ornithological interest. At irregular intervals longer papers and monographs called *Transactions* are published, which members receive free or at a substantial discount.

Communications, and inquiries regarding publications, should be addressed to:

Secretary, Linnaean Society of New York,
c/o American Museum of Natural History,
79th Street at Central Park West,
New York 24, N. Y.

PRINTED IN THE U. S. A.

Corrigenda to
“The Species of Middle American Birds”
(Trans. Linnaean Soc. N. Y. vol. 7, 1955)

The following *corrigenda* are limited to such typographical errors as might be misleading, and such material published before the work went to press which, but for oversight, would have affected the range or nomenclature given. Various persons have advised me of several unpublished records, and there have been several papers recently published that affect the ranges, nomenclature, or taxonomy of a few of the species listed. Though I am very anxious to receive any information that serves to correct, modify or extend the data contained in the list, changes based on unpublished data or on material published after Jan. 1955 would be in the nature of *addenda* rather than *corrigenda*, and are therefore not included here. I am grateful to the following for calling pertinent matters to my attention: J. L. Bull, H. H. Collins, T. R. Howell, E. Mayr, K. C. Parkes, A. R. Phillips, P. Slud, A. Wetmore.

TYPOGRAPHICAL

- p. iii: lines 30 and 31. For “41” read “42.”
- p. 45: line 17. For “Powell” read “Howell.”
- p. 57: line 9. For “thyroides” read “thyroideus.”
- p. 60: line 5. For “Syndactila” read “Syndactyla.”
- p. 75: line 23. For “argenticula” read “argentigula.”
- p. 111: line 24. Omit “A.” after “Grinnell, J.”.
- p. 118: line 25. For “Auk, 23” read “Auk, 18.”
- p. 118: line 31. For “(1951)” read “(1941).”
- p. 123: Index, third column. For “Galinule” read “Gallinule.”
- p. 123: Index, third column. For “Gmpsonyx” read “Gampsonyx.”
- p. 125: Index, first column. After “Muscivora,” for “64” read “66.”
- p. 127: Index, third column. For “Syndactila” read “Syndactyla.”
- p. 128: Index, second column. After “Woodcreeper” and “Woodhewer,” for “57” read “58.”

OTHER CORRECTIONS or CLARIFICATIONS

- p. 11: line 9. *Puffinus puffinus*. Replace period by semicolon and add “ranging n. to California; recorded off Pac. coast Costa Rica (*opisthomelas*, Cabo Blanco, *fide* R. C. Murphy).”
- p. 12: Footnote 5. *Oceanodroma leucorhoa*. After “*socorroensis*” insert “the form breeding on Guadalupe I.”
- p. 17: lines 3-4. *Phoenicopterus ruber*. After “Greater Antilles” insert “Bonaire.”; at end replace period by semicolon and add “recorded in Florida, Colombia, Guianas, n. Brazil.”
- p. 17: Footnote 8. *Branta hutchinsii*. Add “Includes *minima* as well as true *hutchinsii*, both of which occur in Mexico and each of which Conover considers a separate species; Aldrich and Blake are here followed.”

- p. 24: line 19. *Falco sparverius*. After "Guatemala" insert "Honduras."
- p. 27: line 17. *Porzana flaviventer*. Before "and" insert "Panama and n."
- p. 56: line 32. *Centurus rubricomus*. Replace "rubricomus" by "pygmaeus." In Footnote 10, replace period by semicolon and add "but *pygmaeus*, the name of the Cozumel I. race, has priority and becomes the specific name."
- p. 60: line 15. *Automolus rubiginosus*. After "Brit. Honduras" add semicolon and "not recorded from Costa Rica."
- p. 67: line 20. *Myiarchus cinerascens*. Strike out "casually to n.w. Costa Rica." (The old record mentioned in Bent probably relates to *nuttingi*).
- p. 73: Footnote 5. *Progne subis*. Substitute for entire footnote: "Treatment above based on Hellmayr. Includes *sinaloae* of w. Mexico (chiefly mts. Chihuahua, Sonora, Sinaloa, Jalisco, Nayarit), one record Guatemala (Peten); considered by some conspecific with West Indian *dominicensis*, if so Snowy-bellied Martin may be used for the complex. Includes also *cryptoleuca* of Cuba, recorded in winter from Brit. Honduras and Guatemala; considered by some a separate species, by others a race of *dominicensis*, and by still others conspecific (including *dominicensis* complex) with *subis*." (For a recent review, see Zimmer, Amer. Mus. Novit., no. 1723: 1-7, 1955).
- p. 74: line 24. *Calocitta formosa*. For "S.w. Mexico (Colima southward)" substitute "W. Mexico (s. Sonora southward)."
- p. 95: line 35. *Icterus chrysater*. After "Trop. Mid. Am." insert "(Costa Rican record doubtful.)"
- p. 95: Footnote 10. Replace period by semicolon and add "corrected to *Pendulinus* (Wilson Bull., 64: 115, 1952)."
- p. 98: line 14. *Tangara chrysophrys*. Footnote 2: Add new sentence: "Some authors consider *guttata* to have priority as the specific name."

In addition to instances mentioned in the footnotes, many genera listed in this work, which are accepted in the A.O.U. Check-list, are merged with other genera by some ornithologists, particularly in Europe. Those most often merged are: *Puffinus* (in *Procellaria*), *Morus* (in *Sula*), *Chen* (in *Anser*), *Lophodytes* (in *Mergus*), *Squatarola* (in *Pluvialis*, and sometimes both in *Charadrius*), *Totanus* and sometimes also *Actitis* (in *Tringa*), *Ereunetes* and *Erolia* (in *Calidris*), *Steganopus* and *Lobipes* (in *Phalaropus*), *Catharacta* (in *Stercorarius*), *Thalasseus* (in *Sterna*), *Spinus* (in *Carduelis*).

BIBLIOGRAPHY

- p. 114: MEXICO: Under "South central" add:
- * Davis, W. B. and R. J. Russell. 1953. Aves y mamiferos del estado de Morelos. Rev. Soc. Mex. Hist. Nat., 14 (1-4): 77-147. A list of the birds and mammals of Morelos, with localities and bibliography.
- p. 116. EL SALVADOR. Add:
- ** Rand, A. L. and M. A. Traylor. 1954. Manual de las Aves de El Salvador. Univ. de El Salvador. A guide to the birds, with keys and descriptions.
- p. 117. NICARAGUA. Add:
- Nutting, C. C. 1883-1884. On a collection of birds from Nicaragua. (Edited by R. Ridgway). Proc. U. S. Nat. Mus., 6: 372-410. Birds taken or observed at three representative localities.

—EUGENE EISENMAN

CONTENTS

	PAGE
Introduction	1
Method of Selecting English Names	2
Technical Names	4
Ranges	4
Comments on Middle American Bird Distribution	5
Tinamiformes (5)*	
Tinamidae—Tinamous (5)	9
Gaviiformes (3)	
Gaviidae—Loons (3)	9
Colymbiformes (5)	
Colymbidae—Grebes (5)	9
Procellariiformes (31)	
Diomedeidae—Albatrosses (6)	10
Procellariidae—Shearwaters, Petrels (14)	10
Hydrobatidae—Storm-Petrels (11)	11
Pelecaniformes (18)	
Phaethontidae—Tropicbirds (3)	12
Pelecanidae—Pelicans (2)	13
Sulidae—Gannets, Boobies (5)	13
Phalacrocoracidae—Cormorants (5)	14
Anhingidae—Anhingas (1)	14
Fregatidae—Frigatebirds (2)	14
Ciconiiformes (31)	
Ardeidae—Herons (20)	14
Cochleariidae—Boat-billed Herons (1)	16
Ciconiidae—Storks (2)	16
Threskiornithidae—Ibises, Spoonbills (7)	16
Phoenicopteridae—Flamingos (1)	17
Anseriformes (41)	
Anatidae—Ducks, Geese, Swans (41)	17
Falconiformes (65)	
Cathartidae—American Vultures (5)	20
Accipitridae—Hawks (44)	20
Pandionidae—Ospreys (1)	23
Falconidae—Falcons (15)	23

*The numbers in parentheses indicate the number of species here listed from Middle America in each order and family.

	PAGE
Galliformes (31)	
Cracidae—Guans, Curassows (8)	24
Phasianidae—Pheasants, Partridges (21)	25
Meleagrididae—Turkeys (2)	26
Gruiformes (22)	
Gruidae—Cranes (2)	26
Aramidae—Limpkins (1)	26
Rallidae—Rails (17)	26
Heliornithidae—Sungrebes (1)	28
Eurypygidae—Sunbitterns (1)	28
Charadriiformes (87)	
Jakanidae—Jacanas (1)	28
Haematopodidae—Oystercatchers (2)	28
Charadriidae—Plovers (11)	28
Scolopacidae—Snipe, Sandpipers (29)	29
Recurvirostridae—Avocets, Stilts (2)	31
Phalaropodidae—Phalaropes (3)	31
Burhinidae—Thick-knees (1)	32
Stercorariidae—Skuas, Jaegers (4)	32
Laridae—Gulls, Terns (29)	32
Rynchopidae—Skimmers (1)	35
Alcidae—Auks, Murres (4)	35
Columbiformes (32)	
Columbidae—Pigeons (32)	36
Psittaciformes (32)	
Psittacidae—Parrots (32)	38
Cuculiformes (15)	
Cuculidae—Cuckoos (15)	40
Strigiformes (30)	
Tytonidae—Barn Owls (1)	41
Strigidae—Owls (29)	41
Caprimulgiformes (18)	
Nyctibiidae—Potoos (2)	43
Caprimulgidae—Nightjars (16)	43
Apodiformes (117)	
Apodidae—Swifts (17)	44
Trochilidae—Hummingbirds (100)	45
Trogoniformes (14)	
Troganidae—Trogons (14)	52
Coraciiformes (14)	
Alcedinidae—Kingfishers (6)	53
Momotidae—Motmots (8)	53

	PAGE
Piciformes (60)	
<i>Galbulidae—Jacamars (3)</i>	54
<i>Bucconidae—Puffbirds (8)</i>	54
<i>Capitonidae—Barbets (3)</i>	54
<i>Ramphastidae—Toucans (8)</i>	55
<i>Picidae—Woodpeckers (38)</i>	55
Passeriformes (752)	
<i>Dendrocolaptidae—Woodcreepers (20)</i>	58
<i>Furnariidae—Ovenbirds, Spinetails (23)</i>	59
<i>Formicariidae—Antbirds (39)</i>	61
<i>Rhinocryptidae—Tapaculos (2)</i>	63
<i>Pipridae—Manakins (15)</i>	63
<i>Cotingidae—Cotingas (23)</i>	64
<i>Tyrannidae—Tyrant Flycatchers (106)</i>	65
<i>Oxyruncidae—Sharpbills (1)</i>	72
<i>Alaudidae—Larks (1)</i>	73
<i>Hirundinidae—Swallows (14)</i>	73
<i>Corvidae—Crows, Jays (25)</i>	74
<i>Paridae—Titmice (8)</i>	75
<i>Sittidae—Nuthatches (3)</i>	76
<i>Certhiidae—Creepers (1)</i>	76
<i>Chamaeidae—Wrentits (1)</i>	76
<i>Cinclidae—Dippers (1)</i>	76
<i>Troglodytidae—Wrens (45)</i>	77
<i>Mimidae—Thrashers, Mockingbirds (18)</i>	80
<i>Turdidae—Thrushes (31)</i>	81
<i>Zeledoniidae—Wren-Thrushes (1)</i>	83
<i>Sylviidae—Old World Warblers, Kinglets, Gnatcatchers (10)</i>	83
<i>Motacillidae—Wagtails, Pipits (5)</i>	84
<i>Bombycillidae—Waxwings (1)</i>	84
<i>Ptylogonatidae—Silky-flycatchers (4)</i>	84
<i>Laniidae—Shrikes (1)</i>	85
<i>Sturnidae—Starlings (1)</i>	85
<i>Cyclarhidae—Peppershrikes (1)</i>	85
<i>Vireolaniidae—Shrike-Vireos (3)</i>	85
<i>Vireonidae—Vireos (26)</i>	85
<i>Coerebidae—Honeycreepers (9)</i>	87
<i>Parulidae—Wood-Warblers (80)</i>	88
<i>Ploceidae—House Sparrows, Weavers (1)</i>	94
<i>Icteridae—American Orioles, Blackbirds (46)</i>	94
<i>Tersinidae—Swallow-Tanagers (1)</i>	97
<i>Thraupidae—Tanagers (64)</i>	97
<i>Fringillidae—Finches, Grosbeaks, Sparrows (121)</i>	101
Bibliography	
<i>General</i>	109
<i>Mexico</i>	
<i>Entire country</i>	110

	PAGE
Mexico—<i>continued</i>	
Northwest Pacific	111
West central	112
South Pacific	112
Northeast	113
South central	114
Southern Gulf and Caribbean	114
British Honduras	115
Guatemala	116
El Salvador	116
Honduras	116
Nicaragua	117
Costa Rica	117
Panama	118
Neighboring countries	119
Index to genera and group names	121

INTRODUCTION

Improved travel facilities have greatly stimulated interest in the bird life of Middle America—the region from Mexico to Panama. Before long the Inter-American Highway will make possible automobile traffic from the United States to beyond the Canal Zone; even today it is only a few hours flight to the cities of Mexico and Panama. A large proportion of the several thousand forms currently recognized from Middle America are but slightly varying races of species found through much of the region, and in many cases well into South America. Amateurs and ornithological writers have long felt the need for a list of English names by which each species could be designated throughout its range, without regard to locality or the subspecies involved. Actually no list of Middle American species, even by their technical names, has been available since the outdated tabular lists that appeared fifty years ago in *Biologia Centrali-Americana: Aves*.

The great multi-volumed reference works, *Birds of North and Middle America* and *Catalogue of Birds of the Americas*, contain much of the basic data from which a distributional list may be prepared, but they fail to provide English names for species as such. Instead, contrary to modern practice, each subspecies is given a separate common name, without any consistent policy of indicating conspecific relationships. Frequently the names adopted in these works disagree; often they are perfunctory translations of the technical designations without regard to appropriateness; and occasionally the same name is applied to two distinct species. Those wishing to use an English name for the species unit, rather than for each subspecies, have been forced to make their own selections. Selection of a suitable name for a polytypic species requires examination of specimens, or at least literature, covering the range of the species. As the individual writer usually lacks the time or the facilities to make an adequate check for this purpose, he is often likely to choose or invent a name for the species, which, though seemingly appropriate for the subspecies he knows best, may be grossly misleading for the species as a whole. Such individual selections, even when otherwise satisfactory, create a multiplicity of names which handicaps popular ornithology. The use of differing names tends to limit the value of published observations, for others may fail to recognize the species discussed. Now that popular bird guides for the area

are beginning to appear, it seems desirable to lay a foundation, before it is too late, for some measure of uniformity.

This paper provides a convenient list of the fourteen hundred species presently known to have been recorded in Middle America, giving the technical name, a suggested English name, and a brief outline of range. The English names were chosen, in almost all instances, from those already in the literature. Great weight was given to the maintenance of well-settled usage, but as relatively few truly neotropical birds have established English names, very substantial consideration was accorded to the factors of appropriateness to the entire species and utility to the amateur.

The English names here recommended (save for a few variations resulting chiefly from taxonomic or orthographic considerations) have been adopted for their respective areas by the only current handbook covering a Middle American country, Blake's *Birds of Mexico* (1953), by Eisenmann's *Annotated List of Birds of Barro Colorado Island, Panama Canal Zone* (1952), and by Chalif's projected *Field Guide to the Birds of Mexico and Northern Central America*, which will cover the birds of Mexico, British Honduras, Guatemala and El Salvador. The birds treated in these works comprise nearly all species having a wide range in Middle America.

METHOD OF SELECTING ENGLISH NAMES

1. For species reaching Middle America chiefly as migrants from the area north of Mexico the current preferences of the A.O.U. Committee on Classification and Nomenclature have been followed. As no official list of such English species names has yet been issued and there may be changes of view before the publication of the next *A.O.U. Check-List*, in a few instances there is also listed an alternate name regarded by many students as superior. Occasionally, to distinguish the northern species from a closely allied species also found in Middle America, an A.O.U. name has been slightly modified, by adding some such simple prefix as "American", "Northern", "Common", or the like.

2. For neotropical or essentially Middle American species (even though ranging slightly into southwestern United States) the basic idea was to select from the literature an appropriate name, already used for the species as a whole or for one of its subspecies, having if possible some associative value. Names suggesting a characteristic of the species in regard to appearance, behavior, or habitat were favored over patronymics, because such names are often aids in identifying or remembering a species. Geographic names were generally avoided

(except for representative or very local species), as they tend to be misleading or to suggest subspecific relationship.

When more than one appropriate name for a species appeared in the literature, preference was given (other factors being about equal) to the name used in the *A.O.U. Check-List*, in Ridgway's *Birds of North and Middle America*, or in Hellmayr's *Catalogue of Birds of the Americas*, in that order. Though the overwhelming majority of names were drawn from these three works, some were taken also from the writings of Beebe, Bond, Brabourne and Chubb, Gould, Griscom, Penard, Skutch, and van Rossem, and particularly the *Distributional Check-List of the Birds of Mexico* (Part I) and Sutton's *Mexican Birds*. Very few names are wholly new. The availability of the vast neotropical collection of the American Museum of Natural History, to which its custodians kindly offered free access, made possible checking of proposed names against most races of each species. Helpful ideas and comments were received from D. Amadon, R. S. Arbib, Jr., R. A. Paynter, Jr., F. A. Pitelka, A. F. Skutch, R. W. Storer, G. M. Sutton, J. Van Tyne, and particularly Ludlow Griscom and Alexander Wetmore. John L. Bull read the entire manuscript and supplied many useful suggestions, both with respect to names and other matters.

After preparation of a preliminary list, containing alternate English names for many species, it was sent to Emmet R. Blake and Edward L. Chalif (each then working independently on a Middle American bird book) for indication of their respective preferences and suggestions. It was gratifying to find that in almost all cases our preferences were the same. In the relatively few instances where initially we did not concur, agreement was ultimately reached (often after much discussion) on names felt by all three to be acceptable.

It is inevitable that not all our selections will please everyone. In fact they do not all please our little committee. We felt it desirable, for the sake of uniformity, to defer to the views of the A.O.U. Check-List Committee in many instances where we would have favored another name. Certain names of doubtful usefulness we accepted because they had appeared in the literature and we could not find any really good name not conflicting with that of another species. Generally speaking, we preferred a reasonably associative name previously employed in the literature to the invention of a wholly new, though possibly better, name.

Regarding orthography, the recommendations of Cheesman and Oehser (Auk, 54: 333–340, 1937) have been followed in the main. Compound *group names* are generally written as one word, being hyphenated only when a single word seemed clumsy or likely to

obscure relationship, meaning, or pronunciation. If hyphenated, the second word of a group name is capitalized only when designating a subdivision of birds to which the group actually belongs or is allied (*cf.* Common Tody-Flycatcher and Gray Silky-flycatcher, the former is, the latter is not, a member of Tyrannidae, the Tyrant Flycatchers). At the suggestion of R. S. Arbib, in hyphenated *specific names*, doubling of the past participle is avoided for simplicity and euphony (*e.g.*, Bar-tailed and Stripe-crowned are used, rather than Barred-tailed and Striped-crowned).

TECHNICAL NAMES

The technical nomenclature is essentially that of the Hellmayr volumes of the *Catalogue of Birds of the Americas*, supplemented and modified chiefly by Peters' *Check-List of Birds of the World*, the Friedmann volumes of *Birds of North and Middle America*, Zimmer's *Studies of Peruvian Birds*, and the recent Supplements (1944-1954) to the *A.O.U. Check-List of North American Birds*. Since one major purpose of this list is to provide English names for birds that, on the basis of present knowledge, *may* be regarded as species, some forms or groups of forms are here listed as species that many ornithologists (including members of our own committee) consider to be subspecies. Where there is such current disagreement as to the specific rank or relationship of a Middle American bird, this is indicated in a footnote, and if the appropriateness of the suggested English name would be affected by a different taxonomic opinion an alternate name is supplied.

RANGES

Ranges are limited to the barest outline, for the main purpose is to indicate the countries of Middle America where each species is known to occur. The ranges are derived chiefly from the *Catalogue of Birds of the Americas*, with numerous modifications based on other published papers, and in many instances from unpublished data—usually labelled specimens in the American Museum of Natural History or information from qualified ornithologists. More detailed distributional data can, of course, be found in the works already mentioned and in various local papers listed in the annexed bibliography. The following ornithologists have generously given me information as to ranges of certain species: H. Friedmann, F. C. Lincoln (North American migrants); M. Alvarez del Toro, E. R. Blake, L. I. Davis, G. M. Sutton (Mexico); R. A. Paynter, Jr. (Yucatan Peninsula); K. C. Parkes (British Honduras); M. J. Carr, J. C. Dickinson, Jr. (Honduras); L. Griscom (Mexico, Guatemala, Nicaragua, Panama); A. R. Phillips (Mexico,

Guatemala); T. R. Howell (Nicaragua); A. F. Skutch, M. A. Carriker, Jr. (Costa Rica); R. C. Murphy (sea-birds off Panama); C. H. Rogers (Costa Rica, Panama); A. Wetmore (Panama); J. Bond (West Indies, Panama). They should not be held responsible, however, for any errors in the present paper.

Abbreviations: Geographical abbreviations have been freely employed. The constantly iterated "Mid. Am." (Middle America) indicates—unless expressly qualified—that the species occurs in *all* the Middle American countries: Mexico, British Honduras, Guatemala, El Salvador, Honduras, Nicaragua, Costa Rica, and Panama. The same abbreviation preceded by "Trop." (Tropical) imports breeding in some part of the tropical lowlands of each of these countries, though in many cases the species may also range into the mountains, while in others distribution may be very local. "Highlands" indicates that in Middle America the species usually breeds above 2,500–3,000 feet. "Mts." means that the usual breeding range is above 5,000 feet. The breeding range runs to the first semicolon; or, if there is no such mark, to the end.

Many of the gaps in the ranges of individual species doubtless indicate merely gaps in present knowledge or at least in published data. This is certainly true of sea-birds, of northern migrants generally, and of a number of resident species in British Honduras and Nicaragua (for which no published check-list exists), and in Honduras (for which the published check-list is very incomplete). Calling attention to the apparent gaps may help elicit the true situation. While it is unlikely that many *breeding* species wholly new to the Middle American avifauna will be added (except perhaps in the Darien region of eastern Panama adjacent to South America), in *detailed* distributional knowledge the ornithology of Middle America is still at the pioneer stage.

COMMENTS ON MIDDLE AMERICAN BIRD DISTRIBUTION

The very brevity of the range outlines emphasizes certain facts of faunal interest. Thus the exclusion of El Salvador from the range of a lowland species generally indicates that it is a bird of humid forest, absent from the semi-arid Pacific slope over which El Salvador extends. Similarly the absence from British Honduras, El Salvador and Nicaragua of certain widely distributed montane species may be explained by the small area and low altitude of the mountains in those countries.

Generally speaking, plateaus or mountain ranges separate the tropical lowlands of the Atlantic (Gulf and Caribbean) from those of the Pacific slope. The true tropical climate in Middle America has an almost uniform year-round temperature (governed chiefly by altitude),

with two seasons—a rainy season and a dry. The dry season, controlled by the northeast trade winds, comes during the northern winter.

The fact is sometimes overlooked that much of Mexico is within the Temperate Zone geographically, and that the high interior plateaus and mountains carry the climate, vegetation, and fauna of that zone far south of the Tropic of Cancer. Many northern species, including such well-known birds as Red-shafted Flicker (*Colaptes cafer*), Brown Creeper (*Certhia familiaris*), Common Bluebird (*Sialia sialis*), Red Crossbill (*Loxia curvirostra*) and Chipping Sparrow (*Spizella passerina*), range as breeders as far as Nicaragua, which is also the southern limit of pine trees. A few species of northern affinities, like Red-tailed Hawk (*Buteo jamaicensis*), Hairy Woodpecker (*Dendrocopos villosus*), American Dipper (*Cinclus mexicanus*) and Volcano Junco (*Junco vulcani*), breed south to the mountains of western Panama.

The montane avifauna of tropical Middle America is characterized not only by certain northern species, but by some endemics, and by a number of so-called subtropical forms with close relatives in the mountains of western South America. As the Middle American highlands are interrupted by lowland gaps (notably the Isthmus of Tehuantepec, southern Nicaragua and central Panama), this montane fauna is not continuous; yet, as Chapman long ago pointed out, the same species frequently occur from Mexico to Ecuador.

The Caribbean lowlands have a relatively short dry season, relieved by showers, so that the area naturally supports, from Veracruz in Mexico to western Colombia, a perpetually green forest.* The rather recent derivation of the rain forest avifauna from South America is indicated not only by the identity of the species, but also by the increase in variety as one approaches that continent. Major exceptions on the Caribbean slope to the humid climate are the northern part of the Yucatan Peninsula and certain interior valleys in Guatemala, Honduras and Nicaragua, the bird-life of which resembles that of the drier Pacific slope.

As the intervening mountains drain the northeast trade winds of their moisture, there prevails over most of the Pacific lowlands a longer, more severe, dry season. This results, depending on local conditions, in a variety of semi-arid environments, ranging from deciduous forest to grassland and desert scrub. The avifauna includes (in addition to the more widely distributed birds) a number of endemic species, some of which range into southwestern United States. This endemic avifauna

*Man is increasingly destroying this forest; but in Middle America south of Mexico most of the population is on the Pacific slope, and it is on that slope that the forests are most reduced.

ends in northwestern Costa Rica with the appearance on the Pacific coast south of the Gulf of Nicoya of heavy forest, which (locally interrupted by savannas) continues into western Panama. Much of the Pacific coast of Panama is relatively open, with forest on the hillsides and near permanent water. Typical rain forest appears in Darien province, which has but a brief dry season and adjoins the wettest section of South America (northwestern Colombia), with a rather similar and enormously varied bird life. As is to be expected from the geographic situation, in Panama a number of Central American forms meet their South American representatives, raising interesting problems from the viewpoint of speciation.

But present conditions cannot wholly account for present bird distribution. For example, there are absent from the Pacific coast of Panama, where much of the environment seems suitable, a number of species of semi-arid country that breed not only from Mexico to northern Costa Rica but also in South America; e.g., Rufescent Tinamou (*Crypturellus cinnamomeus*), Double-striped Thick-knee (*Burhinus bistriatus*), Common Ground Dove (*Columbigallina passerina*), Grayish Saltator (*Saltator coerulescens*). The gap in some cases extends from Honduras to South America; viz., Vermilion Flycatcher (*Pyrocephalus rubinus*), Green Jay (*Cyanocorax yncas*). To achieve such ranges, it seems likely that at one time these species occupied the intervening area, which thereafter became unsuitable because of climatic or topographic changes. Similarly, the presence in northern Yucatan Peninsula of many species otherwise restricted in Mexico to the Pacific coast suggests a former semi-arid connection through country that today supports humid forest. The surprising number of endemics (lowland as well as mountain species) in southern Costa Rica and the adjacent Chiriquí province of western Panama must indicate that at one time the Costa Rica-Chiriquí area was much more isolated than today.

The occurrence as breeders in Nicaragua of a few birds, otherwise unknown in Middle America, having their nearest relatives in northern South America; e.g., Pearl Kite (*Gampsonyx swainsoni*), Nicaraguan Seed-Finch (*Oryzoborus nuttingi*), is even more puzzling. The explanation may possibly lie in the fact that Nicaragua, occupying the southeastern part of a hump-like projection into the Caribbean, doubtless intercepts storm-driven South American strays, which, in rare instances, may have become established and, through isolation, occasionally have developed a local form.

Regarding North American migrants, it is striking that very few passerine species breeding only in the West winter farther south than

Nicaragua, and most do not seem to go regularly beyond Guatemala. On the other hand, almost all eastern species migrating as far as Guatemala also reach Panama, and many winter to South America. The reason for this pattern in winter ranges is not apparent. But our actual information of winter habitats, migration dates and routes in Middle America is extremely sparse.

In fact so little is recorded of the ecological preferences, local distribution, and behavior of most birds in Middle America that any conscientious observer can add greatly to knowledge.

It is hoped that the following list of some 1,424 species may stimulate study of the living birds, and help to accelerate the day when the *A.O.U. Check-List of North American Birds* will include the avifauna of Middle America—ornithologically the richest part of our North American continent.*

*The number of species here listed in each order and family is indicated in the table of contents, *supra*.

Order TINAMIFORMES

Family TINAMIDAE TINAMOUS

Tinamus major	GREAT TINAMOU
Trop. Mid. Am. (except El Salvador) and n. S. Am.	
Nothocercus bonapartei	HIGHLAND TINAMOU
Highlands Costa Rica, w. Panama and n.w. S. Am.	
Crypturellus soui	LITTLE TINAMOU
Trop. Mid. Am. (except El Salvador) and n.w. S. Am.	
Crypturellus boucardi	SLATY-BREASTED TINAMOU
Carib. slope trop. Mid. Am. to n. Costa Rica (both slopes).	
Crypturellus cinnamomeus	RUFESCENT TINAMOU
Trop. Mid. Am. to n.w. Costa Rica; also n.e. Colombia and Venez.	

Order GAVIIFORMES

Family GAVIIDAE LOONS

Gavia stellata	RED-THROATED LOON
Circumpolar regions; winters to Pac. coast Mexico.	
Gavia arctica	ARCTIC OR BLACK-THROATED LOON
Circumpolar regions; winters to Pac. coast Mexico.	
Gavia immer	COMMON LOON
N. N. Am., Greenland, Iceland and islands n. of Europe; in Am. winters to Pac. coast Mexico.	

Order COLYMBIFORMES

Family COLYMBIDAE GREBES

Colymbus¹ dominicus	LEAST GREBE
S.e. Texas, Mid. Am., Greater Antilles, Bahamas, and trop. S. Am.	
Colymbus caspicus	EARED OR BLACK-NECKED GREBE
Old World, w. N. Am. to n.w. Mexico; winters in Am. to Guatemala; also Colombia.	

¹ It has been proposed that *Podiceps* be adopted for this genus, which would result in changing the family name to Podicipididae and the ordinal name to Podicipidiformes.

<i>Aechmophorus occidentalis</i>	WESTERN GREBE
W. N. Am.; winters to Mexico.	
<i>Podilymbus podiceps</i>	PIED-BILLED GREBE
N. Am., Mid. Am., West Indies and S. Am.	
<i>Podilymbus gigas</i>	ATITLAN GREBE
Guatemala (Lake Atitlan).	

Order PROCELLARIIFORMES

Family DIOMEDEIDAE ALBATROSSES

<i>Diomedea exulans</i>	WANDERING ALBATROSS
Antarctic is.; ranges in s. oceans; once Panama Bay.	
<i>Diomedea irrorata</i>	GALAPAGOS ALBATROSS
Galapagos Is.; ranges n. to Panama and s. to Peru.	
<i>Diomedea albatrus</i>	SHORT-TAILED ALBATROSS
Is. cent. and w. N. Pacific; formerly ranged to Pac. coast N. Am. and Mexico (off Baja California); now nearly extinct.	
<i>Diomedea nigripes</i>	BLACK-FOOTED ALBATROSS
Is. cent. and w. N. Pacific; ranges to Pac. coast N. Am. and Mexico (Baja Calif.).	
<i>Diomedea immutabilis</i>	LAYSAN ALBATROSS
Is. cent. N. Pacific; ranges to Mexico (Baja Calif.).	
<i>Diomedea chrysostoma</i>	GRAY-HEADED ALBATROSS
Is. w. S. Pacific; ranges over s. oceans; once Pac. coast Panama.	

Family PROCELLARIIDAE SHEARWATERS AND PETRELS

<i>Fulmarus glacialis</i>	NORTHERN FULMAR
Circumpolar; winters casually s. to Mexico (Pac. coast).	
<i>Fulmarus² antarcticus</i>	ANTARCTIC FULMAR
Antarctic; ranges to Peru and once to Mexico (Mazatlan).	
<i>Puffinus creatopus</i>	PINK-FOOTED SHEARWATER
Mas Atierra Is., Chile; ranges to Alaska, Mexico (Baja Calif.).	
<i>Puffinus pacificus</i>	WEDGE-TAILED SHEARWATER
Is. of cent. and w. Pacific and of Indian Oceans, and off Mexico (Revilla Gigedo Is.); recorded off Pac. coast Mexico and Panama.	
<i>Puffinus griseus</i>	SOOTY SHEARWATER
Is. S. Pacific; ranges off both coasts N. Am., Pac. coast Mexico, Costa Rica (sight, A. P. Smith) and Panama (sight, R. C. Murphy).	

² Sometimes placed in a separate genus *Priocella*; also called Silver-gray Fulmar.

Puffinus tenuirostris	SHORT-TAILED³ SHEARWATER
Is. off Australia; ranges to Pac. coast N. Am., casually to Mexico and Panama.	
Puffinus nativitatis	CHRISTMAS ISLAND SHEARWATER
Is. trop. (chiefly cent.) Pac.; taken once between Clipperton I. and Mexican coast.	
Puffinus puffinus⁴	COMMON SHEARWATER
Is. cooler waters of the Old World, in America breeds in Mexico (Is. in Gulf of Calif. and off Baja Calif.).	
Puffinus auricularis⁴	REVILLA GIGEDO SHEARWATER
Mexico (Revilla Gigedo Is.).	
Puffinus lherminieri	DUSKY-BACKED SHEARWATER
Is. in pan-tropical waters, in America West Indies, Tobago and Galapagos; ranges to s.e. U. S., in Caribbean Sea, and Pac. coast of n. S. Am.; sight reports off Panama (Pac. coast, R. C. Murphy).	
Pterodroma neglecta	VARIABLE PETREL
Is. warmer waters of Indian and Pac., incl. Juan Fernandez Is. off Chile; taken once off Revilla Gigedo Is., Mexico (<i>fide</i> R. C. Murphy).	
Pterodroma phaeopygia	DARK-RUMPED PETREL
Hawaiian and Galapagos Is.; ranges in America off Pac. coast n. S. Am. to vicinity of Clipperton I. off Pac. coast of Mexico, sight reports off Pac. coast Panama (R. C. Murphy).	
Pterodroma externa	WHITE-NECKED PETREL
Kermadec and Juan Fernandez Is.; ranges in America off Pac. coast S. Am. n. to vicinity of Clipperton I. off Mexico.	
Pterodroma cookii	BLUE-FOOTED PETREL
Is. S. Pacific; ranges in America off Pac. coast S. Am. n. to Mexico (off Baja Calif.); recorded Aleutians.	

Family HYDROBATIDAE STORM-PETRELS

Oceanites oceanicus	WILSON'S PETREL
Antarctic Is.; ranges in America commonly into N. Atl. and casually N. Pac., taken Veracruz, Mexico.	
Oceanites gracilis	WHITE-VENTED PETREL
Probably breeds Peru and Galapagos Is.; ranges off Pac. coast S. Am. n. to Panama (sight, R. C. Murphy).	

³ This, the standard name in Australia where the species breeds, is also used by Murphy in "Oceanic Birds of South America," and is therefore favored over "Slender-billed" Shearwater.

⁴ The A.O.U. is followed in treating *P. opisthomelas*, the Black-vented Shearwater, as a race of *P. puffinus*. R. C. Murphy would also treat *auricularis* as a subspecies.

Pelagodroma marina	WHITE-FACED PETREL
Coasts of Australia, New Zealand and is. in S. Atl., trop. e. N. Atl. and w. S. Pac.; ranges in America to Cocos I., Costa Rica and Galapagos.	
Oceanodroma tethys	WEDGE-RUMPED PETREL
Galapagos and Peruvian Is.; ranges off Pac. coast Mexico, Costa Rica (Cocos I.) and Panama.	
Oceanodroma castro	BAND-RUMPED PETREL
Is. in pan-tropical waters of e. and cent. Atl. (Madeira, Azores to Ascension and St. Helena) and cent. and e. Pac. (Hawaiian and Galapagos Is.); in Mid. Am. recorded off Pac. coast Costa Rica (Cocos I., possibly breeding).	
Oceanodroma leucorhoa	LEACH'S PETREL
Is. N. Atl. and e. N. Pac., Mexico (taxonomy of forms disputed) (Is. in Gulf of Calif., off Baja Calif. and Guadalupe I.); migrates to S. Atl., Caribbean Sea and e. S. Pac. to Galapagos; recorded once off Pac. coast Guatemala. ⁵	
Oceanodroma macroura	GUADALUPE PETREL
Mexico (Guadalupe I.); possibly extinct.	
Oceanodroma markhami	SOOTY PETREL
Ranges off Peru and Chile; taken near Clipperton I. off Pac. coast Mexico and Costa Rica (Cocos I.).	
Oceanodroma homochroa	ASHY PETREL
Is. off Calif. and Mexico (Baja Calif.).	
Oceanodroma Melania	BLACK PETREL
Mexico (Is. off Baja Calif. and in Gulf of Calif.); ranges to Pac. coast Panama and S. Am.	
Halocyptena microsoma	LEAST PETREL
Mexico (Is. off Baja Calif. and in Gulf of Calif.); ranges to Pac. coast Panama and S. Am.	

Order PELECANIFORMES

Family PHAETHONTIDAE TROPICBIRDS

Phaethon aethereus	RED-BILLED TROPICBIRD
Is. in trop. Atl. (including West Indies), Pac. and Indian Oceans: Mexico (Pac. coast), Panama (Swan Key, Caribbean coast); recorded also Pac. coast El Salvador, Honduras, Costa Rica (Gulf of Nicoya).	

⁵ The form *socorroensis*, treated by Peters as a race of the Asiatic *O. monorhis*.

Phaethon lepturus	WHITE-TAILED TROPICBIRD
Is. in trop. Atl. (including West Indies), cent. Pac. and Indian Oceans; in Mid. Am. recorded off Caribbean Guatemala (Puerto Barrios).	
Phaethon rubricauda	RED-TAILED TROPICBIRD
Is. in trop. cent. Pac. and Indian Oceans; in Mid. Am. casual off Pac. coast Mexico (Baja Calif. and Revilla Gigedo Is.).	

Family PELECANIDAE PELICANS

Pelecanus erythrorhynchus	AMERICAN WHITE PELICAN
N. cent. N. Am.; winters to Mexico and Guatemala. ⁶	
Pelecanus occidentalis	BROWN PELICAN
S.w. Canada, w. and s. U. S., West Indies, Mid. Am. and S. Am.	

Family SULIDAE GANNETS AND BOOBIES

Morus⁷ bassanus	NORTHERN GANNET
Local, is. off N. Atl. coasts; winters to Gulf of Mexico (Veracruz).	
Sula nebouxii	BLUE-FOOTED BOOBY
Local, trop. is. Pac.: Mexico, Costa Rica, Panama and n.w. S. Am.	
Sula dactylatra	BLUE-FACED OR MASKED BOOBY
Local, trop. is. Atl. (incl. Caribbean), Pac. and Indian Oceans: Mexico (Yucatan Pen., Baja Calif.); recorded from Guatemala (Pac. sight, A. J. van Rossem), Nicaragua and Panama (both coasts).	
Sula sula	RED-FOOTLED BOOBY
Local, trop. is. Atl. (incl. Caribbean), Pac. and Indian Oceans: Mexico (Tres Marias and Revilla Gigedo Is.), Brit. Honduras, Honduras, Costa Rica (Cocos I.); recorded off coast of Mexico (Pac.), Nicaragua (Caribbean), Panama (Caribbean; sight off Pearl Is., B. B. Sturgis).	
Sula leucogaster	BROWN BOOBY
Local, trop. is. Atl. (incl. Caribbean), Pac. and Indian Oceans: Mexico (Yucatan Pen., Gulf of Calif., Pac. coast), Brit. Honduras, and both coasts of Honduras, Nicaragua, Costa Rica and Panama; ranging off Guatemala (Caribbean, sight, L. Griscom), and El Salvador (sight, O. Salvin).	

⁶ There seems to be no evidence that this species occurs farther south.

⁷ Some authors merge this genus in *Sula*.

Family PHALACROCORACIDAE CORMORANTS

Phalacrocorax auritus	DOUBLE-CRESTED CORMORANT
N. Am., Bahamas, Cuba, Mexico (Yucatan Pen. and Pac. coast), Brit. Honduras.	
Phalacrocorax olivaceus	OLIVACEOUS OR NEOTROPIC CORMORANT
Louisiana, Texas, Bahamas, Cuba, Mid. Am. and S. Am.	
Phalacrocorax penicillatus	BRANDT'S CORMORANT
Pac. coast N. Am. to Mexico (Gulf of Calif. and Baja Calif.).	
Phalacrocorax pelagicus	PELAGIC CORMORANT
Pac. coast N. Am. to n. Mexico (n.w. Baja Calif.).	
Phalacrocorax bougainvillii	GUANAY CORMORANT
Is. off Pac. coast, Peru and Chile; ranges n. to Panama (sight off Darien, once, R. C. Murphy).	

Family ANHINGIDAE ANHINGAS

Anhinga anhinga	ANHINGA
S. U. S., Cuba, Mid. Am. and S. Am.	

Family FREGATIDAE FRIGATEBIRDS

Fregata magnificens	MAGNIFICENT FRIGATEBIRD
Coastal is. of trop. Atl. and e. Pac., Mid. Am. and S. Am.; ranging along both coasts of Mid. Am.	
Fregata minor	GREAT FRIGATEBIRD ⁸
Is. of Pac., Indian and S. Atl. oceans, Mexico (Revilla Gigedo Is.); casually ranging off Pac. coast Mexico.	

Order CICONIIFORMES

Family ARDEIDAE HERONS

Ardea herodias	GREAT BLUE HERON
N. Am., West Indies, Mexico, Galapagos Is.; regular throughout the year in Mid. Am., though breeding uncertain; winters to n. S. Am.	
Ardea occidentalis⁹	GREAT WHITE HERON
S. Florida and Greater Antilles; Mexico (Yucatan Pen.).	

⁸ This is the established name in the literature of Australia, where the species is common, and where "Lesser" Frigatebird is used for *F. ariel*.

⁹ Perhaps a color phase of *A. herodias*.

Ardea cocoi	WHITE-NECKED HERON
S. Am.; e. Panama (sight, once, A. Wetmore).	
Pilherodius pileatus	CAPPED HERON
Panama and S. Am.	
Butorides virescens	GREEN HERON
N. Am., West Indies, Mid. Am.; winters to n. S. Am.	
Butorides striatus¹⁰	STRIATED HERON
Qld World tropics, e. Panama and S. Am.	
Florida caerulea	LITTLE BLUE HERON
S. U. S., West Indies, Mid. Am. and S. Am.	
Dichromana rufescens	REDDISH EGRET
S. U. S., Bahamas, Greater Antilles and Mexico (both coasts); winters to Pac. coast Guatemala and El Salvador.	
Casmerodus¹ albus	COMMON OR LARGE EGRET
Old World, U. S., West Indies, Mid. Am. and S. Am.	
Leucophoyx¹ thula	SNOWY EGRET
U. S., West Indies, Mid. Am. and S. Am.	
Bubulcus² ibis	CATTLE EGRET
Warmer parts of Old World, recently breeding in n. S. Am. and Florida; present (possibly breeding) in e. U. S. and Panama (sight, E. Eisenmann, F. O. Chapelle, repeatedly and in numbers).	
Hydranassa tricolor	TRICOLORED HERON
S. U. S., West Indies, Mid. Am. and S. Am.	
Agamia agami	CHESTNUT-BELLIED HERON
Local: trop. Mid. Am. (not recorded El Salvador, Honduras and Nicaragua) and S. Am.	
Nycticorax nycticorax	BLACK-CROWNED NIGHT-HERON
Old World, N. Am., West Indies, Mid. Am. and S. Am.	
Nyctanassa violacea	YELLOW-CROWNED NIGHT-HERON
U. S., West Indies, Mid. Am. and S. Am.	
Tigrisoma lineatum	BANDED TIGER-HERON
Honduras, Nicaragua, Costa Rica, Panama and S. Am.; Mexico (Chiapas, once).	
Heterocnus mexicanus	BARE-THROATED TIGER-HERON
Trop. Mid. Am.	
Ixobrychus exilis	LEAST BITTERN
N. Am., Mid. Am. and S. Am.; the northern race <i>exilis</i> winters through Mid. Am.	

¹⁰ If considered conspecific with *B. virescens*, Green Heron may be used for the entire complex.

¹ Placed by some authors in the genus *Egretta*.

² Placed by some authors in the genus *Ardeola*.

Botaurus lentiginosus	AMERICAN BITTERN
N. Am.; winters through Mid. Am.	
Botaurus pinnatus	PINNATED BITTERN
S. Am.; recorded from e. Nicaragua and s.e. Mexico (Quintana Roo, <i>fide</i> R. A. Paynter).	

Family COCHLEARIIDAE³ BOAT-BILLED HERONS

Cochlearius cochlearius	BOAT-BILLED HERON
Trop. Mid. Am. and S. Am.	

Family CICONIIDAE STORKS

Mycteria americana	WOOD IBIS OR WOOD STORK
S. U. S., Greater Antilles, Mid. Am. and S. Am.	
Jabiru mycteria	JABIRU
Rare: trop. Mid. Am. (not reported from Brit. Honduras and Honduras), and S. Am.	

Family THRESKIORNITHIDAE IBISES AND SPOONBILLS

Theristicus caudatus	BUFF-NECKED IBIS
S. Am.; recorded from e. Panama (once).	
Mesembrinibis cayennensis	GREEN IBIS
Panama and S. Am.	
Eudocimus albus	WHITE IBIS
S. U. S., Greater Antilles, Mid. Am. and n. S. Am.	
Eudocimus ruber	SCARLET IBIS
Trop. S. Am.; accidental in Honduras and West Indies.	
Plegadis falcinellus	GLOSSY IBIS
Old World; s.e. U. S., Greater Antilles; e. Mexico (?); recorded from e. Panama once, n. Colombia.	
Plegadis mexicana⁴	WHITE-FACED IBIS
W. U. S. to Mexico, also s. S. Am.; winter records in Guatemala, El Salvador and Costa Rica.	
Ajaia ajaja	ROSEATE SPOONBILL
S. U. S., Greater Antilles, Mid. Am. and S. Am.	

³ Some authors believe this group is best merged in Ardeidae.

⁴ If a race of *P. falcinellus*, the entire complex would be called Glossy Ibis. Hellmayr and Conover say the proper name of this form is *P. chihi*.

Family PHOENICOPTERIDAE FLAMINGOS

Phoenicopterus ruber⁵ AMERICAN FLAMINGO
Bahamas, Greater Antilles, Mexico (Yucatan Pen.), Venezuela,
Guianas, Galapagos Is.

Order ANSERIFORMES

Family ANATIDAE DUCKS, GEESE AND SWANS

Cygnus columbianus	WHISTLING SWAN
Arctic N. Am.; winters casually to Mexico (Baja Calif.).	
Cygnus buccinator	TRUMPETER SWAN
Local: n.w. N. Am.; one record from Mexico (Tamaulipas).	
Chen caerulescens⁶	BLUE GOOSE
Arctic N. Am.; winters to Mexico.	
Chen hyperborea⁶	SNOW GOOSE
Arctic N. Am.; winters to Mexico.	
Chen rossii	Ross's GOOSE
Arctic n.w. Canada; winters casually to Mexico (Chihuahua).	
Anser albifrons	WHITE-FRONTED GOOSE
Arctic regions; in America winters to Mexico.	
Branta bernicla⁷	BRANT
Arctic regions; winters to Mexico (Baja California).	
Branta canadensis	CANADA GOOSE
N. N. Am.; winters to Mexico.	
Branta hutchinsii⁸	CACKLING GOOSE
Arctic America and n.e. Siberia; in America winters to Mexico.	
Dendrocygna viduata	WHITE-FACED TREE-DUCK
Trop. Africa, Madagascar and S. Am.; occurs in Costa Rica and Panama.	
Dendrocygna bicolor	FULVOUS TREE-DUCK
Trop. Asia, Madagascar and E. Africa, s.w. U. S., Mexico, Guatemala and Honduras (<i>fide</i> M. J. Carr), also S. Am.; casual in e. Panama.	

* If the Old World *P. antiquorum* is deemed a race of this species, the name Greater Flamingo may be used.

* These birds may be color phases of the same species. If so, the scientific name would be *C. caerulescens*, the English name Snow Goose.

⁷ Birds recorded from Mexico are the western form, sometimes regarded as a species and called Black Brant, *B. nigricans*, which Zimmer and Delacour say should be called *orientalis*.

⁸ Generally regarded as conspecific with the Canada Goose, *B. canadensis*.

Dendrocygna autumnalis	BLACK-BELLIED TREE-DUCK
S.w. U. S., Mid. Am., and S. Am.	
Sarkidiornis sylvicola⁹	SOUTH AMERICAN COMB-DUCK
Trop. S. Am.; recorded from e. Panama once.	
Cairina moschata	MUSCOVY
Trop. Mid. Am. and S. Am.	
Anas carolinensis¹⁰	GREEN-WINGED TEAL
N. N. Am.; winters to West Indies, Mexico, Guatemala, Brit. Honduras and Honduras; casual in Colombia.	
Anas acuta	COMMON PINTAIL
Holarctic regions; winters through Mid. Am. (not reported El Salvador) to Colombia.	
Anas platyrhynchos	COMMON MALLARD
Holarctic regions; in America winters regularly to Mexico; recorded from Guatemala, Nicaragua, Costa Rica and Panama.	
Anas wyvilliana	HAWAIIAN DUCK
Hawaiian Is.; once at Mazatlan, Sinaloa, Mexico.	
Anas diazi	MEXICAN DUCK
S.w. U. S. and Mexico.	
Anas fulvigula	MOTTLED DUCK
S. U. S. and e. Mexico.	
Anas strepera	GADWALL
Holarctic regions; winters to Mexico.	
Anas cyanoptera	CINNAMON TEAL
W. N. Am. to n. Mexico, also S. Am.; northern birds winter regularly at least to Mexico, with scattered records from Guatemala, Nicaragua, Costa Rica, Panama and n. S. Am.	
Anas discors	BLUE-WINGED TEAL
N. Am. and possibly Mexico; winters through Mid. Am. and S. Am.	
Spatula¹ clypeata	NORTHERN SHOVELER
Holarctic regions; winters through Mid. Am. (not recorded Brit. Honduras) to n. S. Am.	
Mareca¹ penelope	EUROPEAN WIDGEON
Europe and n. Asia; in America winters to U. S.; once in Mexico (Baja Calif.).	

⁹ May be a race of the Old World Comb-Duck, *S. melanotos*; if so the name Comb Duck will suffice.

¹⁰ If deemed a race of the Old World *A. crecca*, the species can be called Green-winged Teal.

¹ These genera should perhaps be merged in *Anas* (Delacour and Mayr, Wilson Bull., 57:17, 1945).

Mareca americana	BALDPATE OR AMERICAN WIDGEON
N. N. Am.; winters sparingly through Mid. Am. (not recorded Brit. Honduras, El Salvador) to Colombia.	
Aix sponsa	WOOD DUCK
N. Am., Cuba; winters to Mexico.	
Aythya valisineria	CANVASBACK
N. N. Am.; winters to Mexico and Guatemala.	
Aythya americana	REDHEAD
N.w. N. Am.; winters to Mexico and Guatemala.	
Aythya collaris	RING-NECKED DUCK
N. N. Am.; winters to Mexico, Guatemala, Honduras; Panama (sight, R. T. and K. T. Scholes).	
Aythya marila	GREATER SCAUP
Holarctic regions; in America winters to West Indies and n.w. Mexico.	
Aythya affinis	LESSER SCAUP
N.w. N. Am.; winters through Mid. Am. to Colombia.	
Melanitta deglandi²	WHITE-WINGED SCOTER
N. w. N. Am.; winters in America to Mexico (Baja Calif.).	
Melanitta perspicillata	SURF SCOTER
N. N. Am.; winters to n.w. Mexico.	
Melanitta nigra	BLACK SCOTER
N. Holarctic; winters in America to Mexico (Baja Calif., seen C. L. Hubbs).	
Bucephala clangula	COMMON GOLDENEYE
Holarctic regions; in America winters to w. Mexico.	
Bucephala albeola	BUFFLEHEAD
N.w. N. Am.; winters to Mexico.	
Lophodytes cucullatus	HOODED MEGANSER
N. N. Am.; winters to Mexico.	
Mergus merganser	COMMON MEGANSER OR GOOSANDER
Holarctic regions; winters in America to Mexico.	
Mergus serrator	RED-BREASTED MEGANSER
Holarctic regions; winters in America to Mexico.	
Oxyura dominica	MASKED DUCK
Local: Texas, Greater Antilles, Mexico, Guatemala, Costa Rica, Panama, and S. Am.	
Oxyura jamaicensis	RUDDY DUCK
N. Am., West Indies, Mexico, Brit. Honduras, Guatemala; re- corded Costa Rica.	

² By some considered a race of the Old World *M. fusca*.

Order FALCONIFORMES

Family CATHARTIDAE AMERICAN VULTURES

Sarcoramphus papa	KING VULTURE
Trop. Mid. Am. and S. Am.	
Coragyps atratus	BLACK VULTURE
S. U. S., Mid. Am., S. Am.	
Cathartes burrovianus³	YELLOW-HEADED VULTURE
S. Am.; recorded from Mexico (Tamaulipas, Veracruz), Panama (Pac. slope, common, may breed).	
Cathartes aura	TURKEY VULTURE
N. Am., West Indies, Mid. Am., S. Am.	
Gymnogyps californianus	CALIFORNIA CONDOR
California and formerly Mexico (Baja Calif.).	

Family ACCIPITRIDAE HAWKS

Elanus leucurus	WHITE-TAILED KITE
Calif., s. U. S., Mexico (Baja Calif. and Gulf Coast), also S. Am.; winters to Guatemala and Brit. Honduras.	
Gampsonyx⁴ swainsonii	PEARL KITE
W. Nicaragua and n. S. Am.	
Elanoides forficatus	SWALLOW-TAILED KITE
S. U. S., Mid. Am. (not reported El Salvador), and S. Am.	
Leptodon cayanensis⁵	GRAY-HEADED KITE
Trop. Mid. Am. and S. Am.	
Chondrohierax uncinatus	HOOK-BILLED KITE
Trop. Mid. Am., S. Am., Grenada	
Harpagus bidentatus	DOUBLE-TOOTHED KITE
Trop. Mid. Am. (not reported Brit. Honduras and Honduras), and S. Am.	
Ictinia mississippiensis	MISSISSIPPI KITE
S. U. S.; winters at least to Mexico and Guatemala; recorded Paraguay.	
Ictinia plumbea	PLUMBEOUS KITE
Trop. Mid. Am. and S. Am.	
Rostrhamus sociabilis	EVERGLADE OR SNAIL KITE
Local in marshes: Fla., Cuba, Mid. Am. (except El Salvador) and S. Am.	

³ *C. urubitinga* of many authorities.

⁴ Peters and Hellmayr and Conover place this genus in *Falconidae*; Friedmann is here followed.

⁵ *Odontorchis palliatus* of many authorities.

Accipiter gentilis	NORTHERN GOSHAWK
Holarctic regions, in America to mts. of n.w. Mexico.	
Accipiter bicolor	BICOLORED HAWK
Trop. Mid. Am. (except El Salvador) and S. Am.	
Accipiter cooperii	COOPER'S HAWK
N. Am. and n. Mexico; winters to Guatemala; recorded Costa Rica.	
Accipiter superciliosus	TINY HAWK
Nicaragua, Costa Rica, Panama and S. Am.	
Accipiter striatus	SHARP-SHINNED HAWK
N. Am., Greater Antilles, and highlands of Mexico; winters in Mid. Am. to w. Panama (not reported Brit. Honduras and Honduras).	
Accipiter chionogaster⁶	WHITE-BREASTED HAWK
Pine woods s. Mexico (Chiapas), Guatemala, Honduras, El Salvador and Nicaragua, also mts. of n. S. Am.	
Heterospizias meridionalis	SAVANNA HAWK
Panama and S. Am.	
Buteo albicaudatus	WHITE-TAILED HAWK
Local, s.w. U. S., Mid. Am. (not recorded Brit. Honduras) and S. Am.	
Buteo regalis	FERRUGINOUS HAWK
W. N. Am.; winters to Mexico.	
Buteo jamaicensis	RED-TAILED HAWK
N. Am., West Indies, and highlands of Mid. Am. to w. Panama; northern birds migrate to Nicaragua, perhaps occasionally to Panama.	
Buteo albonotatus	ZONE-TAILED HAWK
S.w. U. S., local in Mid. Am. (breeding status uncertain in Costa Rica and Panama; not reported Brit. Honduras) to S. Am.	
Buteo lineatus	RED-SHOULDERED HAWK
N. Am. to Mexico.	
Buteo swainsoni	SWAINSON'S HAWK
W. N. Am. and n.w. Mexico; migrates through Mid. Am. (not reported Brit. Honduras, Honduras, Nicaragua) to S. Am.	
Buteo platypterus	BROAD-WINGED HAWK
N. Am., West Indies; migrates and winters through Mid. Am. and S. Am.	
Buteo magnirostris	ROADSIDE HAWK
Trop. Mid. Am. and S. Am.	

⁶This bird has been treated as a race of *A. erythronemius*, the Rufous-thighed Hawk, of s. S. Am.; and may be a race of *A. striatus*, according to Storer.

Buteo brachyurus	SHORT-TAILED HAWK
Local: Florida, Mid. Am. (not reported El Salvador, Honduras), and S. Am.	
Buteo nitidus	GRAY HAWK
S.w. U. S., Mid. Am. and S. Am.	
Parabuteo unicinctus	BAY-WINGED HAWK
S.w. U.S., Mid. Am. (not reported Brit. Honduras, Honduras) and S. Am.	
Leucopternis albicollis	WHITE HAWK
Trop. Mid. Am. (except El Salvador) and S. Am.	
Leucopternis semiplumbea	SEMIPLUMBEOUS HAWK
Honduras, Nicaragua, Costa Rica, Panama and n.w. S. Am.	
Leucopternis plumbea⁷	PLUMBEOUS HAWK
E. Panama to n.w. Peru.	
Leucopternis princeps	BARRED HAWK
Costa Rica, Panama to Ecuador.	
Busarellus nigricollis	BLACK-COLLARED HAWK
Local: trop. Mid. Am. and S. Am.	
Buteogallus anthracinus	COMMON BLACK HAWK
S.w. U. S., Mid. Am., S. Am., Lesser Antilles (St. Vincent).	
Hypomorphus⁸ urubitinga	GREAT BLACK HAWK
Trop. Mid. Am. and S. Am.	
Harpyhaliaetus⁹ solitarius	SOLITARY EAGLE
Rare, (chiefly highlands?) Mexico, Guatemala, Nicaragua, Costa Rica, Panama and n. S. Am.	
Morphnus guianensis¹⁰	CRESTED EAGLE
Honduras, Nicaragua, Costa Rica, Panama and S. Am.	
Harpia harpyja	HARPY EAGLE
Trop. Mid. Am. (not recorded Brit. Honduras, El Salvador, Honduras) and S. Am.	
Spizastur melanoleucus	BLACK-AND-WHITE HAWK-EAGLE¹
Trop. Mid. Am. (not recorded Brit. Honduras, El Salvador) and S. Am.	

⁷ Regarded by Hellmayr and Conover as a race of *L. schistacea*, the Slate-colored Hawk, of eastern South America.

⁸ Hellmayr and Conover use *Urubitinga*; the genus may yet be merged in *Buteogallus*, according to Amadon.

⁹ A separate genus *Urubitornis* is by some authors erected for this species; while others make it a race of *H. coronatus* of s. S. Am.

¹⁰ If *M. taeniatus*, the Banded Crested Eagle (known from Panama to Guiana, Brazil and Bolivia), be a good species, rather than a color phase of *M. guianensis*, then the latter may be distinguished by the name Common Crested Eagle.

¹ "Eagle-Hawk" is also used for this and the next genus; but "Hawk-Eagle" seems preferable to conform with the older usage and that in the Old World for the same group.

Spizaetus ornatus	ORNATE HAWK-EAGLE
Trop. Mid. Am. and S. Am.	
Spizaetus tyrannus	BLACK HAWK-EAGLE
Trop. Mid. Am. (except El Salvador) and S. Am.	
Aquila chrysaetos	GOLDEN EAGLE
Mts. (chiefly) of Eurasia and N. Am. to Mexico.	
Haliaeetus leucocephalus	BALD EAGLE
N. Am. to n.w. Mexico (Baja Calif.; recorded Veracruz).	
Circus cyaneus	MARSH HAWK OR WHITE-RUMPED HARRIER
Eurasia and N. Am. to Mexico (Baja Calif.); winters through Mid. Am. to Colombia.	
Geranospiza nigra?	BLACKISH CRANE-HAWK
Trop. Mid. Am. and n.w. S. Am.	

Family PANDIONIDAE OSPREYS

Pandion haliaetus	OSPREY
Old World, N. Am., Cuba, Bahamas, Mexico (Pac. Coast and Yucatan Pen.), and Brit. Honduras; winters, and frequently summers, through Mid. Am. and S. Am.	

Family FALCONIDAE FALCONS

Herpetotheres cachinnans	LAUGHING FALCON
Trop. Mid. Am. and S. Am.	
Micrastur semitorquatus	COLLARED FOREST-FALCON
Trop. Mid. Am. and S. Am.	
Micrastur mirandollei	SLATY-BACKED FOREST-FALCON
Costa Rica, Panama and n. S. Am.	
Micrastur ruficollis	BARRED FOREST-FALCON
Trop. Mid. Am. and S. Am.	
Daptrius americanus	RED-THROATED CARACARA
Trop. Mid. Am. (except Brit. Honduras and El Salvador) and S. Am.	
Milvago chimachima	YELLOW-HEADED CARACARA
Panama (Pac. slope grasslands) and S. Am.	
Polyborus cheriway *	CRESTED CARACARA
S. U. S., Cuba, Mid. Am. and n. S. Am.	
Polyborus lutosus	GUADALUPE CARACARA
Mexico (Guadalupe I.); now extinct.	

* If but one species in this genus is recognized, the specific name *caerulescens* must be used and the "Blackish" in the English name omitted.

* Hellmayr and Conover use *Caracara* for the genus (but cf. Amadon, Auk 71: 203-4, 1954), and suggest that *cheriway* may be conspecific with *plancus* of S. Am.

Falco mexicanus	PRAIRIE FALCON
W. N. Am. and n.w. Mexico; winters to s. Mexico.	
Falco peregrinus	PEREGRINE FALCON
Old World, N. Am., Mexico (Baja Calif.), s. Chile and Argentina; the N. Am. race, <i>anatum</i> , has been recorded in winter or on migration through Mid. Am. (except Nicaragua) and in S. Am.	
Falco deiroleucus	ORANGE-BREASTED FALCON
Local: Guatemala, Nicaragua, Costa Rica, Panama and S. Am.; Mexico (Veracruz, once).	
Falco albicularis^{3a}	BAT FALCON
Trop. Mid. Am. and S. Am.	
Falco femoralis	APLOMADO FALCON
Local: s.w. U. S., Mexico, Brit. Honduras, Guatemala, Nicaragua, w. Panama and S. Am.	
Falco columbarius	PIGEON HAWK OR MERLIN
Eurasia and N. Am.; in America winters through Mid. Am. (not reported Brit. Honduras and El Salvador) and S. Am.	
Falco sparverius	AMERICAN SPARROW HAWK OR AMERICAN KESTREL
N. Am., West Indies, Mexico, Guatemala, also S. Am.; winters through Mid. Am. and S. Am.	

Order GALLIFORMES

Family CRACIDAE GUANS AND CURASSOWS

Crax rubra	GREAT CURASSOW
Trop. Mid. Am. to Ecuador.	
Penelope purpurascens	CRESTED GUAN
Trop. Mid. Am. and n. S. Am.	
Ortalis wagleri	RUFOUS-BELLIED CHACHALACA
W. Mexico.	
Ortalis vetula	PLAIN CHACHALACA
S.e. Texas, Mexico, Guatemala, Brit. Honduras, Honduras, El Salvador, Nicaragua.	
Ortalis garrula	CHESTNUT-WINGED CHACHALACA
Nicaragua, Costa Rica, Panama and Colombia.	
Penelopina nigra	BLACK CHACHALACA
Mts. s. Mexico (e. Oaxaca, Chiapas), Guatemala, El Salvador, Honduras and Nicaragua.	
Chamaepetes unicolor	BLACK GUAN
Mts. Costa Rica and w. Panama.	

^{3a} Hellmayr and Conover consider *F. rufigularis* the correct name. Friedmann is here followed.

Oreophasis derbianus HORNED GUAN
Mts. s. Mexico (Chiapas) and Guatemala.

Family PHASIANIDAE PARTRIDGES AND PHEASANTS

Dendrocyx barbatus	BEARDED WOOD-PARTRIDGE
Mexico (mts. Veracruz).	
Dendrocyx macroura	LONG-TAILED WOOD-PARTRIDGE
Mexico (mts. Jalisco and Veracruz to Oaxaca).	
Dendrocyx leucophrys	BUFFY-CROWNED WOOD-PARTRIDGE
Highlands s. Mexico (Chiapas), Guatemala, Honduras, El Salvador, Nicaragua and Costa Rica.	
Oreortyx picta	MOUNTAIN QUAIL
Mts. w. U. S. and Mexico (Baja Calif.).	
Callipepla squamata	SCALED QUAIL
Arid. s.w. U. S. and Mexico (s. to Jalisco and Morelos).	
Lophortyx californica	CALIFORNIA QUAIL
Pac. U. S. and n.w. Mexico	
Lophortyx gambelii	GAMBEL'S OR DESERT QUAIL
Deserts s.w. U. S. to n.w. Mexico.	
Lophortyx douglasii	ELEGANT QUAIL
W. Mexico (Sonora and Chihuahua to Jalisco).	
Philortyx fasciatus	BANDED QUAIL
Highlands south central Mexico.	
Colinus virginianus	COMMON BOBWHITE
N. Am., Cuba, Mexico and Guatemala.	
Colinus nigrogularis	BLACK-THROATED BOBWHITE
Mexico (Yucatan Pen.), Brit. Honduras, and Caribbean slope of Guatemala, Honduras (Segovia River) and (?) Nicaragua.	
Colinus leucopogon ⁴	SPOT-BELLIED BOBWHITE
Highlands and Pacific slope Guatemala, Honduras, El Salvador, Nicaragua and Costa Rica.	
Colinus cristatus	CRESTED BOBWHITE
Panama and n. S. Am.	
Odontophorus gujanensis	MARBLED WOOD-QUAIL
S.w. Costa Rica, Panama and S. Am.	
Odontophorus erythrops	RUFOUS-FRONTED WOOD-QUAIL
Caribbean Honduras, Nicaragua, Costa Rica, Panama and n.w. S. Am.	
Odontophorus leucolaemus	WHITE-THROATED WOOD-QUAIL
Highlands Costa Rica and w. Panama.	

⁴ Considered forms of the Crested Bobwhite (*C. cristatus*) by some authorities.

Odontophorus guttatus	SPOTTED WOOD-QUAIL
Trop. s. Mexico and highlands of Mid. Am. (except El Salvador) to w. Panama.	
Dactylortyx thoracicus	SINGING QUAIL
Highlands (chiefly) s. Mexico, Guatemala, El Salvador and Honduras.	
Cyrtonyx montezumae	HARLEQUIN QUAIL
Highlands s.w. U. S. and Mexico (s. to Oaxaca).	
Cyrtonyx ocellatus	OCELLATED QUAIL
Mts. s.w. Mexico, Guatemala, Honduras, El Salvador and Nicaragua.	
Rhynchortyx cinctus	TAWNY-FACED QUAIL
Caribbean Honduras, Nicaragua, Costa Rica, Panama (both slopes) and n.w. S. Am.	

Family MELEAGRIDIDAE TURKEYS

Meleagris gallopavo	COMMON TURKEY
E. and s.w. U. S. and Mexico (chiefly mts.)	
Agriocharis ocellata	OCELLATED TURKEY
Mexico (Yucatan Pen.), Brit. Honduras and Guatemala (Peten).	

Order GRUIFORMES

Family GRUIDAE CRANES

Grus americana	WHOOPING CRANE
Local: w. Canada; wintering to Gulf coast Mexico.	
Grus canadensis	SANDHILL CRANE
N.e. Siberia, w. and s.e. N. Am., Cuba; winters to Mexico.	

Family ARAMIDAE LIMPKINS

Aramus guarauna	LIMPKIN
S. U. S., West Indies, Mid. Am. and S. Am.	

Family RALLIDAE RAILS

Rallus elegans	KING RAIL
E. N. Am. and Cuba; winters to Mexico.	
Rallus longirostris ⁵	CLAPPER RAIL
U. S., West Indies, Mexico (both coasts and interior marshes), Brit. Honduras, and n. S. Am.	

⁵ The rails of western U.S. and the Pacific coast and interior marshes of Mexico are by some considered races of *R. elegans*. Perhaps they should be treated as a third species *R. obsoletus*, Western Rail.

Rallus limicola	VIRGINIA RAIL
N. Am. to cent. Mexico, also S. Am.; winters to Guatemala.	
Pardirallus maculatus	SPOTTED RAIL
Cuba, Mexico (Chiapas), Brit. Honduras and S. Am.	
Amaurolimnas concolor	UNIFORM CRAKE
Local: trop. Mid. Am. (except El Salvador) and S. Am.	
Aramides cajanea	GRAY-NECKED WOOD-RAIL
Trop. Mid. Am. and S. Am.	
Aramides axillaris	RUFOUS-NECKED WOOD-RAIL
Local: trop. Mid. Am. (unrecorded Guatemala and El Salvador), and n. S. Am.	
Porzana carolina	SORA
N. Am. to Mexico (Baja Calif.); recorded in winter from Mexico, Brit. Hond., Guatemala, Nicaragua, Costa Rica, Panama and S. Am.	
Porzana flaviventer	YELLOW-BREASTED CRAKE
Local: Greater Antilles, El Salvador, Nicaragua, and e. S. Am.	
Laterallus jamaicensis	BLACK RAIL OR CRAKE
Local: U. S., Mexico (Baja Calif.), Jamaica, Peru, Chile; winters to Guatemala.	
Laterallus exilis	GRAY-BREASTED CRAKE
Local: Caribbean Honduras, Nicaragua and n. S. Am.	
Laterallus albicularis*	WHITE-THROATED CRAKE
Nicaragua, Costa Rica, Panama and n.w. S. Am.	
Laterallus ruber	RUDDY CRAKE
Trop. Mexico, Brit. Honduras, Guatemala, El Salvador, Honduras and Nicaragua.	
Coturnicops noveboracensis	YELLOW RAIL
Local: N. Am., Mexico (Lerma, Mexico).	
Gallinula chloropus	COMMON GALLINULE
Old World, N. Am., West Indies, local Mid. Am. (not reported Brit. Honduras) and S. Am.; winters through Mid. Am. to S. Am.	
Porphyrrula martinica	PURPLE GALLINULE
S. U. S., West Indies, Mid. Am. (not reported Brit. Honduras) and S. Am.	
Fulica americana	AMERICAN COOT
N. Am., West Indies, Mexico, Guatemala, Honduras, Nicaragua, Costa Rica, w. Panama and S. Am.; winters through Mid. Am. (not reported Brit. Honduras).	

* By some deemed conspecific with the wide-ranging *L. melanophaius* of S. Am.

Family **HELIORNITHIDAE** SUNGREBES

Heliornis fulica SUNGREBE⁷
Local: trop. Mid. Am. (except El Salvador), and S. Am.

Family **EURYPYGIDAE** SUNBITTERNS

Eurypyga helias SUNBITTERN
Trop. Mid. Am. (except El Salvador), and S. Am.

Order CHARADRIIFORMES⁸

Family **JACANIDAE** JACANAS

Jacana spinosa⁹ AMERICAN JACANA
Texas, Greater Antilles, Mid. Am. and S. Am.

Family **HAEMATOPODIDAE** OYSTERCATCHERS

Haematopus palliatus¹⁰ AMERICAN OYSTERCATCHER
Local: N. Am., West Indies, Mexico, Guatemala, Costa Rica,
Panama and S. Am.

Haematopus bachmani¹⁰ BLACK OYSTERCATCHER
Pac. coast N. Am. to Mexico (n. Baja Calif.).

Family **CHARADRIIDAE** PLOVERS

Belonopterus chilensis¹¹ SOUTHERN LAPWING
S. Am.; a few Panama records.

Hoploxypterus cayanus PIED PLOVER
Trop. S. Am.; doubtfully reported from Honduras (Aloor River).

Squatarola squatarola BLACK-BELLIED PLOVER
Arctic; migrates and winters through Mid. Am. and S. Am.

Pluvialis dominica AMERICAN GOLDEN PLOVER
Arctic N. Am.; in Am. winters in S. Am., recorded on migration
from Mexico, Guatemala, Nicaragua, Costa Rica and Panama
(sight, T. A. Imhof).

⁷ Sometimes called American Finfoot.

⁸ In this order, particularly in the families Charadriidae, Scolopacidae and Laridae,
many non-breeding individuals, especially immatures, summer within their winter
range.

⁹ Birds from central Panama through S. Am. perhaps constitute a distinct species,
J. jacana, the Wattled Jacana; if so, *J. spinosa* may be called Middle American
Jacana. Both *spinosa* and a race of the *jacana* group (*hypomelaena*, formerly called
nigra) have been taken in w. Panama.

¹⁰ Many authorities deem these races of the Common Oystercatcher, *H. ostralegus*,
of the Old World.

¹¹ Some authorities deem *cayennensis* the proper specific name.

Charadrius semipalmatus ²	SEMIPALMATED PLOVER
Arctic N. Am.; winters through Mid. Am. and S. Am.	
Charadrius melanotos	PIPING PLOVER
E. N. Am.; winters to n. Mexico and West Indies.	
Charadrius alexandrines	SNOWY PLOVER
Old World, Pac. and Gulf coasts of U. S., West Indies, Mexico (Baja Calif. and Tamaulipas?), Peru, Chile; winters in Mexico.	
Charadrius collaris	COLLARED PLOVER
Local: trop. Mid. Am., S. Am., s. Lesser Antilles.	
Charadrius vociferus	KILLDEER
N. Am., Mexico, West Indies, Peru and n. Chile; winters through Mid. Am. and n.w. S. Am.	
Charadrius wilsonia	THICK-BILLED PLOVER
S. U. S., West Indies, Mexico, Brit. Honduras, n. S. Am.; migrates in Mid. Am., West Indies, to n. S. Am.; winters (and may possibly breed) in El Salvador, Honduras, Costa Rica and Panama.	
Eupoda montana	MOUNTAIN PLOVER
Arid plains of w. U. S.; winters to Mexico.	

Family SCOLOPACIDAE SNIPE, SANDPIPER AND ALLIES

Bartramia longicauda	UPLAND PLOVER OR SANDPIPER
N. Am.; migrates through Mid. Am., winters in s. S. Am.	
Numenius borealis	ESKIMO CURLEW
Arctic Am.; winters in S. Am.; recorded Mexico and Guatemala; possibly extinct.	
Numenius phaeopus ³	WHIMBREL
Arctic regions; in America winters through Mid. Am. (not re- ported Brit. Honduras) and S. Am.	
Numenius americanus	LONG-BILLED CURLEW
W. N. Am.; winters to Mexico and Guatemala; reported from Honduras (sight, <i>fide</i> W. Stone).	
Limosa haemastica	HUDSONIAN GODWIT
Canada; migrates through U. S. and West Indies to s. S. Am.; once Mexico (Salina Cruz sight, L. I. Davis, R. A. Herbert).	
Limosa fedoa	MARBLED GODWIT
Int. N. Am.; winters s. U. S. to n. Chile, recorded from Mexico, Guatemala, Brit. Honduras, Honduras, Panama (sight repeatedly, E. Eisenmann, F. O. Chapelle).	

² Perhaps a race of the Old World *C. hiaticula*; if so, call Ringed Plover.

³ The Middle American birds are the North American *hudsonicus*, long regarded
as a species and called Hudsonian Curlew.

Totanus flavipes	LESSER YELLOWLEGS
Canada; winters s. U. S. through Mid. Am. to S. Am.	
Totanus melanoleucus⁴	GREATER YELLOWLEGS
Canada and Alaska; winters s. U. S. through Mid. Am. to S. Am.	
Tringa solitaria	SOLITARY SANDPIPER
N. N. Am.; winters through Mid. Am. to S. Am.	
Actitis macularia	SPOTTED SANDPIPER
N. Am.; winters through Mid. Am. to S. Am.	
Catoptrophorus semipalmatus	WILLET
Local: N. Am., n.e. Mexico; winters through Mid. Am. to S. Am.	
Heteroscelus incanus	WANDERING TATTLER
Alaska; winters chiefly on islands of w. Pacific, also w. Mexico, Costa Rica (Cocos I.), Colombia (Malpelo I.), Galapagos Is.	
Aphriza virgata	SURFBIRD
Alaska; winters to Pacific coast S. Am., recorded Mexico, Guate- mala, Panama (sight repeatedly, E. Eisenmann, T. A. Imhof, F. O. Chapelle).	
Arenaria interpres	RUDDY TURNSTONE
Arctic regions; winters through Mid. Am. to S. Am.	
Arenaria melanocephala	BLACK TURNSTONE
Alaska; winters to Mexico (Baja Calif. and Sonora).	
Limnodromus griseus	COMMON DOWITCHER
Canada and Alaska; winters through Mid. Am. to S. Am.	
Limnodromus scolopaceus⁵	LONG-BILLED DOWITCHER
N. and w. Alaska; migrates chiefly through w. N. Am. to Mexico and Guatemala.	
Capella⁶ gallinago	COMMON SNIPE
Holarctic regions to n. Mexico; the N. Am. form, <i>delicata</i> , winters through Mid. Am. to n. S. Am.	
Calidris canutus	RED KNOT
Arctic regions; in America winters chiefly on both coasts of s. S. Am.; recorded Mexico, Guatemala (sight, A. W. Anthony), Honduras, Costa Rica (Pac. coast).	
Crocethia alba	SANDERLING
Arctic regions; winters through Mid. Am. and S. Am.	
Ereunetes pusillus	SEMIPALMATED SANDPIPER
Arctic N. Am.; winters through Mid. Am. and S. Am.	

⁴ *Tringa melanoleuca* of some authorities, who also place *T. flavipes* in *Tringa*.

⁵ Until recently regarded as a race of *L. griseus* (Auk, 71: 311; Pitelka, Univ. Cal. Publ. Zool., 50: 1-11).

⁶ *Gallinago* of some authorities; the American form *delicata* has been known as Wilson's Snipe.

Ereunetes mauri	WESTERN SANDPIPER
Alaska; winters through Mid. Am. (not reported Brit. Honduras, Honduras) and n. S. Am.	
Erolia minutilla	LEAST SANDPIPER
Arctic N. Am.; winters through Mid. Am. to S. Am.	
Erolia fuscicollis	WHITE-RUMPED SANDPIPER
Arctic N. Am.; migrates through Mid. Am. (not reported Guatemala) to s. S. Am.	
Erolia bairdii	BAIRD'S SANDPIPER
Arctic N. Am.; recorded on migration Mexico, El Salvador, Costa Rica and Panama; winters in s. S. Am.	
Erolia melanotos	PECTORAL SANDPIPER
Arctic N. Am.; winters in s. S. Am., migrates through Mid. Am.	
Erolia alpina	DUNLIN
Arctic regions; the N. Am. race <i>pacifica</i> ⁷ winters to Mexico; one record from Nicaragua.	
Micropalama himantopus	STILT SANDPIPER
Arctic N. Am.; winters s. S. Am., recorded in Mexico, Guatemala, Nicaragua, Panama (sight, R. S. Arbib, F. Loetscher.).	
Tryngites subruficollis	BUFF-BREASTED SANDPIPER
Arctic N. Am.; winters s. S. Am., recorded Mexico, Honduras, El Salvador, Costa Rica, Panama.	

Family RECURVIROSTRIDAE STILTS AND AVOCETS

Himantopus mexicanus⁸	BLACK-NECKED STILT
U. S., West Indies, locally Mexico and S. Am.; ranges (perhaps breeds) through Mid. Am. (not reported Brit. Honduras).	
Recurvirostra americana	AMERICAN AVOCET
W. N. Am. and n. Mexico; winters to Mexico and Guatemala.	

Family PHALAROPODIDAE PHALAROPES

Phalaropus fulicarius	RED PHALAROPE
Arctic regions; in America winters at sea s. to Chile, migrates off Pacific coast of Mexico.	
Steganopus tricolor	WILSON'S PHALAROPE
W. N. Am.; winters in s. S. Am., recorded from Mexico, Guatemala, El Salvador, Panama (sight, R. S. Arbib, F. Loetscher).	

⁷ Often called Red-backed Sandpiper.

⁸ By some all Stilts are considered races of the Old World *H. himantopus*; if so call simply Stilt.

Lobipes lobatus NORTHERN PHALAROPE
Arctic regions; in America winters at sea s. to Peru; recorded from Mexico, Guatemala, Costa Rica.

Family BURHINIDAE THICK-KNEES

Burhinus bistriatus DOUBLE-STRIPED THICK-KNEE
Arid trop. Mid. Am. to n.w. Costa Rica, also in S. Am. and Hispaniola; (?) Panama (a captive bird).

Family STERCORARIIDAE SKUAS AND JAEGERS

Catharacta skua^{8a} GREAT SKUA
Arctic, Antarctic, and Sub-Antarctic; *chilensis* ranges to N. Pac.: Panama (sight, A. Wetmore).

Stercorarius pomarinus POMARINE JAEGER
Arctic; in America winters at sea to s. S. Am.; recorded off Mexico and Panama (Caribbean, *fide* C. H. Rogers; sight, Pac. coast R. C. Murphy).

Stercorarius parasiticus PARASITIC JAEGER
Arctic; winters at sea to s. S. Am.; recorded off Mexico and Panama (both coasts sight, L. Griscom, R. C. Murphy).

Stercorarius longicaudus LONG-TAILED JAEGER
Arctic; winters at sea to s. S. Am.; recorded off Mexico (?), and Panama (Caribbean coast, sight, L. Griscom).

Family LARIDAE GULLS AND TERNS

Larus modestus GRAY GULL
Deserts n. Chile and Peru; ranges along Pac. coast S. Am. n. (at least occasionally) to Panama (seen Panama Bay near Canal entrance, R. C. Murphy).

Larus heermanni HEERMANN'S GULL
Pac. coast Mexico; ranges to Br. Col. and winters to Pac. Guatemala.

Larus delawarensis RING-BILLED GULL
Interior n. N. Am.; winters to Mexico (Pac. coast); reported El Salvador (sight, A. Van Rossem).

^{8a} *C. chilensis*, the Chilean Skua, breeding at the southern tip of S. Am., may be a species distinct from the northern *C. skua*, and there may be several southern species (R. C. Murphy, Oceanic Birds of South America, 2: 1006-1010, 1936).

Larus argentatus ⁹	HERRING GULL
Europe, n. Asia, n. N. Am.; winters to Mid. Am., on Pac. coast to El Salvador, on Caribbean coast to Panama (not reported Nicaragua and Costa Rica).	
Larus californicus	CALIFORNIA GULL
Interior N. Am.; winters to Mexico (chiefly Pac. coast).	
Larus occidentalis ¹⁰	WESTERN GULL
Pac. coast U. S. and Mexico.	
Larus glaucescens	GLAUCOUS-WINGED GULL
Pac. coast n. N. Am. and n.e. Asia; in America winters to Pac. coast Mexico.	
Larus atricilla	LAUGHING GULL
E. and s. U. S., West Indies, Mexico (Yucatan Pen., Sinaloa), Brit. Honduras; winters both coasts Mid. Am. and S. Am.	
Larus pipixcan	FRANKLIN'S GULL
Interior N. Am.; winters chiefly n.w. S. Am.; recorded Mexico, Guatemala, El Salvador, Costa Rica, Panama (both coasts).	
Larus ridibundus	BLACK-HEADED OR BROWN-HEADED GULL
Old World; casual off Atl. coast U. S.; accidental in Mexico (Veracruz).	
Larus philadelphicus ¹	BONAPARTE'S GULL
Interior N. Am.; winters to West Indies and Mexico (both coasts).	
Rissa tridactyla	BLACK-LEGGED KITTIWAKE
Arctic; in America winters off Atl. coast of U. S. and Pac. coast to Mexico.	
Creagrus furcatus	SWALLOW-TAILED GULL
Galapagos Is. and Malpelo I. (off Pac. coast Colombia); casually to Panama (?) (vaguely reported by H. S. Swarth to occur occasionally off Pac. coast).	
Xema sabini	SABINE'S GULL
Arctic; in America winters off Pac. coast n.w. S. Am.; recorded Mexico (Baja Calif.), Panama (sight, R. C. Murphy).	
Chlidonias niger	BLACK TERN
Europe, Asia and N. Am.; migrates through Mid. Am.; in America winters in Panama and S. Am.	

⁹ Nelson's Gull (*L. nelsoni*), supposedly a hybrid of this species with the Glaucous Gull (*L. hyperboreus*), has been taken once off Baja Calif.

¹⁰ Some authors would lump this group in *L. marinus*, the Great Black-backed Gull of the North Atlantic.

¹ Published reports that Col. Grayson collected the Eurasian Little Gull (*L. minutus*) near Mazatlan on the Pac. coast of Mexico on March 27, 1868 are erroneous; the specimen was *L. philadelphicus* (*fide* F. Pitelka and L. C. Stone).

Gelochelidon nilotica	GULL-BILLED TERN
Local: Old World, s. U. S., West Indies, Mexico, also eastern S. Am.; in America winters Mexico, Guatemala, Costa Rica, Panama, West Indies, and S. Am.	
Hydroprogne caspia²	CASPIAN TERN
Local: Old World and N. Am.; winters to Mexico.	
Sterna hirundo	COMMON TERN
Old World, N. Am.; in America winters to S. Am. recorded through Mid. Am. on migration or in winter.	
Sterna forsteri	FORSTER'S TERN
Local: Interior N. Am. and Atl. coast U. S.; winters to Mexico and Guatemala.	
Sterna dougallii	ROSEATE TERN
Local: Old World, Atl. and Gulf coasts N. Am., West Indies, Brit. Honduras; in America winters from La. and Bahamas to e. S. Am.; accidental Mexico (Ventosa Bay).	
Sterna anaethetus	BRIDLED TERN
Local: Pan-tropical islands, West Indies, Brit. Honduras; recorded Pac. coast Mexico, Nicaragua (sight, L. Griscom), Costa Rica, Panama.	
Sterna fuscata	SOOTY TERN
Local: Pan-tropical islands, Florida Keys, West Indies, Mexico (Yucatan Pan., Revilla Gigedo Is.), Brit. Honduras, Honduras, Clipperton I., Galapagos Is.; recorded Pac. coast Mexico (Baja Calif.), Costa Rica (off Cocos I.), Panama.	
Sterna albifrons	LEAST TERN
Old World, U. S., West Indies, Mexico (both coasts), Brit. Honduras; in America winters to Guatemala, casually to Nicaragua (sight, L. Griscom), and e. S. Am.	
Thalasseus maximus	ROYAL TERN
S. U. S., West Indies, Mexico, West Africa; in America winters through Mid. Am. to S. Am.	
Thalasseus elegans³	ELEGANT TERN
W. Mexico; ranges n. to Calif. and winters off Pac. coast of S. Am. s. to Chile; recorded Pac. coast Guatemala, El Salvador, Honduras, Nicaragua (sight, L. Griscom).	
Thalasseus sandvicensis	SANDWICH OR YELLOW-NIBBED TERN
Local: Old World, s. U. S., Bahamas, Mexico (Yucatan Pen.), Brit. Honduras; in America winters to Mexico (Gulf coast), Guatemala, Honduras, Panama (both coasts), West Indies, S. Am.	

² Some authors use the name *tschegrava*.

³ Perhaps properly called *T. comatus*.

Anous stolidus	BROWN NODDY
Local: Pan-tropical islands, Florida Keys, West Indies, Mexico (is. off Yucatan Pen. and Pac. coast), Brit. Honduras, Costa Rica (Cocos I.), n.w. Colombia (Octavia Rocks off Pac. coast), Galapagos Is.; recorded Guatemala, Honduras, Panama (Pac. coast, sight, B. B. Sturgis).	
Anous minutus	WHITE-CAPPED NODDY
Local: small trop. islands in Pac. and S. Atl. and Caribbean, Clipperton Is. off Pac. coast of Mexico, Brit. Honduras, Costa Rica (Cocos I.); ranges in Pac. s. to Colombia.	
Gygis alba	WHITE TERN ⁴
Local: small oceanic islands in trop. Pac., Indian and S. Atl., Clipperton I., Mexico (Revilla Gigedo Is.), Costa Rica (Cocos I.), Galapagos Is.	

Family RYNCHOPIDAE SKIMMERS

Rynchops nigra	BLACK SKIMMER
U. S., Mexico, Guatemala, S. Am.; recorded also El Salvador, Nicaragua, Panama (sight, both coasts, L. Griscom, E. Eisenmann, J. L. Bull), West Indies.	

Family ALCIDAE AUKS, MURRES AND ALLIES

Endomychura hypoleuca⁵	XANTUS'S MURRELET
Calif. and Mexico (islands off Baja Calif.); ranges off Pac. coast of Mexico.	
Synthliboramphus antiquus	ANCIENT MURRELET
Coasts and islands of Alaska and n.e. Asia; wintering casually to Mexico (Baja Calif.).	
Ptychoramphus aleuticus	CASSIN'S AUKLET
Pac. coast N. Am. to Mexico (islands off Baja Calif.).	
Cerorhinca monocerata	RHINOCEROS AUKLET
Coasts of N. Pac.; in America winters to Mexico (n. Baja Calif.).	

⁴ "Fairy" Tern, sometimes used, is confusing with *Sterna nereis*, to which that name is applied in Australia where both occur.

⁵ *Brachyramphus hypoleucus* of some authorities. The form *craveri*, breeding on islands in the Gulf of Calif., is sometimes regarded as a separate species, called Craveri's Murrelet.

Order COLUMBIIFORMES

Family COLUMBIDAE PIGEONS

Columba livia	ROCK DOVE OR DOMESTIC PIGEON
Old World; domesticated in the New World, semiferal about many Mexican towns.	
Columba leucocephala	WHITE-CROWNED PIGEON
Florida Keys, West Indies, islands off Caribbean coast of Mexico (Yucatan Pen.), Brit. Honduras, Honduras, Nicaragua and w. Panama; casual mainland of Mexico.	
Columba flavirostris	RED-BILLED PIGEON
Texas and Mid. Am. (except Brit. Honduras) to n. Costa Rica.	
Columba cayennensis	PALE-VENTED PIGEON
Trop. Mid. Am. (except El Salvador), and S. Am.	
Columba fasciata ⁶	BAND-TAILED PIGEON
Mts. w. N. Am., Mid. Am. (except Brit. Honduras) and S. Am.	
Columba speciosa	SCALED PIGEON
Trop. Mid. Am. (except El Salvador), and S. Am.	
Columba nigrirostris	SHORT-BILLED PIGEON
Trop. Mid. Am. (except El Salvador).	
Columba subvinacea	RUDDY PIGEON
Mts. Costa Rica, Panama and S. Am.	
Columba chiriquensis	CHIRIQUI PIGEON
Known only from one spec. Panama (Chiriqui).	
Ectopistes migratoria ⁷	PASSENGER PIGEON
Extinct: N. Am.; recorded in Mexico and (?) Guatemala.	
Zenaidura macroura	MOURNING DOVE
N. Am., West Indies, in Mid. Am. breeding locally Mexico, Brit. Honduras, Honduras, w. Panama; winters through Mid. Am.	
Zenaidura graysoni	SOCORRO DOVE
Mexico (Socorro I. in the Revilla Gigedos).	
Zenaida aurita	ZENAIDA DOVE
Florida Keys, West Indies, and Mexico (Yucatan Pen. and adjacent islands).	
Zenaida asiatica	WHITE-WINGED DOVE
Greater Antilles, Bahamas, s.w. U. S., and locally Mid. Am. (not reported Brit. Honduras) to w. S. Am.; northern birds winter at least to El Salvador.	

⁶ Birds from Costa Rica, Panama and S. Am. are sometimes regarded as a separate species *C. albilinea*, White-naped Pigeon.

⁷ *E. canadensis* of some authors.

Scardafella inca ⁸	INCA DOVE
S.w. U. S., arid areas of Mid. Am. (except Brit. Honduras) to n.w. Costa Rica.	
Columbigallina passerina	COMMON GROUND-DOVE
S. U. S., West Indies, Mid. Am. to Costa Rica, also S. Am.	
Columbigallina minuta	PLAIN-BREASTED GROUND-DOVE
S.e. Mexico, Brit. Honduras, Guatemala, El Salvador, s.w. Costa Rica, Panama and S. Am.	
Columbigallina talpacoti	RUDDY GROUND-DOVE
Trop. Mid. Am. and S. Am.	
Claravis pretiosa	BLUE GROUND-DOVE
Trop. Mid. Am. and S. Am.	
Claravis mondetoura	MAROON-CHESTED GROUND-DOVE
Mts. trop. Mexico, Guatemala, Honduras, Costa Rica, Panama, and n. S. Am.	
Lepotila verreauxi	WHITE-TIPPED DOVE ⁹
S.w. U. S., Mid. Am. and S. Am.	
Lepotila jamaicensis	CARIBBEAN DOVE
Jamaica, Grand Cayman I., St. Andrew's I., Mexico (Yucatan Pen.).	
Leptotila plumbeiceps	GRAY-HEADED DOVE
Trop. Mid. Am. (except El Salvador) to w. Panama and w. Colombia.	
Leptotila cassinii	GRAY-CHESTED DOVE
Trop. Mid. Am. (except El Salvador), and Colombia.	
Geotrygon veraguensis	OLIVE-BACKED QUAIL-DOVE
Costa Rica, Panama and n.w. S. Am.	
Geotrygon lawrencii	PURPLISH-BACKED QUAIL-DOVE
Local: highlands trop. Mexico (Veracruz), Costa Rica and w. Panama.	
Geotrygon goldmani	RUSSET-CROWNED QUAIL-DOVE
Mts. e. Panama and adjacent Colombia.	
Geotrygon costaricensis	BUFF-FRONTED QUAIL-DOVE
Highlands Costa Rica and w. Panama (Pacific slope).	
Geotrygon violacea	VIOLACEOUS QUAIL-DOVE
Local: Nicaragua, Costa Rica, Panama and S. Am.	
Geotrygon montana	RUDDY QUAIL-DOVE
Trop. Mid. Am. (except El Salvador), and S. Am. and West Indies.	

⁸ Considered a subspecies of the S. Am. *S. squammata* by some.

⁹"White-fronted" Dove, sometimes used, is misleading for most races.

Geotrygon albifacies ¹⁰	WHITE-FACED QUAIL-DOVE
Mts. trop. Mexico, Guatemala, Honduras, El Salvador and Nicaragua.	
Geotrygon chiriquensis ¹⁰	RUFOUS-BREASTED QUAIL-DOVE
Mts. Costa Rica and W. Panama.	

Order PSITTACIFORMES

Family PSITTACIDAE PARROTS

Ara ararauna	BLUE-AND-YELLOW MACAW
E. Panama and S. Am.	
Ara militaris	MILITARY MACAW
Mexico (chiefly mts. and semi-arid areas) and w. S. Am.	
Ara ambiguus	GREAT GREEN MACAW
Nicaragua, Costa Rica, Panama and n.w. S. Am.	
Ara macao	SCARLET MACAW
Trop. Mid. Am. (except Brit. Honduras) and S. Am.	
Ara chloroptera	RED-BLUE-AND-GREEN MACAW
E. Panama and S. Am.	
Ara severa	CHESTNUT-FRONTED MACAW
E. Panama and S. Am.	
Aratinga holochlora ¹	GREEN PARAKEET
Trop. Mexico, Guatemala, Honduras, El Salvador and Nicaragua.	
Aratinga finschi	CRIMSON-FRONTED PARAKEET
Nicaragua, Costa Rica and w. Panama.	
Aratinga astec	OLIVE-THROATED PARAKEET
Gulf and Caribbean slope of trop. Mid. Am. to w. Panama.	
Aratinga canicularis	ORANGE-FRONTED PARAKEET
Pacific slope trop. Mid. Am. to w. Costa Rica.	
Aratinga pertinax ²	BROWN-THROATED PARAKEET
W. Panama (Pac. slope), n. S. Am. and West Indies.	
Rhynchopsitta pachyrhyncha	THICK-BILLED PARROT
Mts. n.w. Mexico; wandering n. to Arizona and to e. Mexico.	
Rhynchopsitta terrisi	MAROON-FRONTED PARROT
Mts. n.e. Mexico (Nuevo Leon and Tamaulipas).	

¹⁰ Races of *G. linearis* of e. S. Am., according to some authors; if so, White-faced Quail-Dove may still be used.

¹ Birds of the Pacific slope from Oaxaca southward are sometimes considered a separate species *A. strenua*, Pacific Parakeet, as are those from e. Guatemala southward *A. rubritorquis*, Red-throated Parakeet.

² The Panama bird *A. ocularis*, Veragua Parakeet, has been considered a separate species.

Pyrrhura hoffmanni	SULPHUR-WINGED PARAKEET
Mts. Costa Rica and w. Panama.	
Bolborhynchus lineola	BARRED PARAKEET
Mts. s. Mexico, Guatemala, Honduras, Costa Rica, w. Panama and S. Am.	
Forpus cyanopygius	BLUE-RUMPED PARROTLET
N.w. Mexico.	
Forpus passerinus	BLUE-WINGED PARROTLET
N. and e. S. Am.; one doubtful Panama record.	
Forpus conspicillatus	SPECTACLED PARROTLET
E. Panama and w. Colombia.	
Brotogeris jugularis	ORANGE-CHINNED PARAKEET
Trop. Mid. Am. (except Brit. Honduras) and n. S. Am.	
Touit costaricensis³	RED-FRONTED PARROTLET
Highlands Costa Rica and w. Panama.	
Touit dilectissima³	BLUE-FRONTED PARROTLET
Mts. e. Panama and n. S. Am.	
Pionopsitta haematocephala	BROWN-HOODED PARROT
Trop. Mid. Am. (except El Salvador), and n.w. S. Am.	
Pionopsitta pyrilia	SAFFRON-HEADED PARROT
E. Panama, Colombia and Venezuela.	
Pionus menstruus	BLUE-HEADED PARROT
Costa Rica, Panama and S. Am.	
Pionus senilis	WHITE-CROWNED PARROT
Trop. Mid. Am. (except El Salvador) to w. Panama.	
Amazona xantholora	YELLOW-LORED PARROT
Mexico (Yucatan Pen.) and Brit. Honduras.	
Amazona albifrons	WHITE-FRONTED PARROT
Chiefly arid trop. areas Mid. Am. to w. Costa Rica.	
Amazona viridigenalis	RED-CROWNED PARROT
N.e. Mexico.	
Amazona finschi	LILAC-CROWNED PARROT
W. Mexico.	
Amazona autumnalis	RED-LORED PARROT
Trop. Mid. Am. (except El Salvador), and n.w. S. Am.	
Amazona ochrocephala⁴	YELLOW-HEADED PARROT
Trop. Mid. Am. and S. Am.	

³ If *T. costaricensis* be treated as a race of *T. dilectissima*, as is done by some, then the name Red-winged Parrotlet may be used for the species.

⁴ The wholly yellow-headed birds from most of Mexico, *A. oratrix*, have often been regarded as specifically separable from *A. auropalliata*, "Yellow-naped" Parrot, of Oaxaca and Chiapas to n.w. Costa Rica. Birds from Panama southward belong to the true *ochrocephala* group, which, if the species be subdivided, could be called "Yellow-crowned" Parrot.

Amazona farinosa MEALY PARROT
Trop. Mid. Am. (except El Salvador), and S. Am.

Order CUCULIFORMES
Family CUCULIDAE CUCKOOS

Coccyzus erythrophthalmus	BLACK-BILLED CUCKOO
N. Am.; on migration recorded through Mid. Am. (except Brit. Honduras and El Salvador); winters chiefly in n.w. S. Am.	
Coccyzus americanus	YELLOW-BILLED CUCKOO
N. Am. and n. Mexico; migrates through Mid. Am. (not reported Brit. Honduras); winters in S. Am.	
Coccyzus minor	MANGROVE CUCKOO
S. Florida, West Indies, trop. Mid. Am. and n. S. Am.	
Coccyzus lansbergi	GRAY-CAPPED CUCKOO
N.w. S. Am.; one doubtful Panama record.	
Piaya cayana	SQUIRREL CUCKOO
Trop. Mid. Am. and S. Am.	
Piaya minuta	LITTLE CUCKOO
Central Panama to S. Am.	
Crotophaga major	GREATER ANI
Central Panama to S. Am.	
Crotophaga ani	SMOOTH-BILLED ANI
Florida, West Indies, Mexico (is. off Yucatan Pen.), Honduras (Caribbean is.), Nicaragua (Caribbean is.), s.w. Costa Rica (Pac. slope, rare), Panama, and S. Am.	
Crotophaga sulcirostris	GROOVE-BILLED ANI
Texas, Mid. Am. and n. S. Am.	
Tapera naevia	STRIPED CUCKOO
Trop. Mid. Am. (except Brit. Honduras) and S. Am.	
Morococcyx erythropygus	LESSER ^b GROUND-CUCKOO
Trop. Mid. Am. (except Brit. Honduras) to w. Costa Rica.	
Dromococcyx phasianellus	PHEASANT CUCKOO
Trop. Mid. Am. (except Brit. Honduras) and S. Am.	
Geococcyx californianus	GREATER ROADRUNNER
S.w. U. S. to cent. Mexico.	
Geococcyx velox^a	LESSER ROADRUNNER
W. and s. Mexico, Guatemala, El Salvador, Honduras, n. Nicaragua.	

^a Van Rossem criticized as misleading Ridgway's name "Rufous-rumped" Cuckoo.

^b According to Stresemann *vaticus* may have priority (Condor, 1954: 90).

Neomorphus geoffroyi RUFOUS-VENTED GROUND-CUCKOO
Nicaragua, Costa Rica, Panama and S. Am.

Order STRIGIFORMES

Family TYTONIDAE BARN OWLS

Tyto alba BARN OWL
Practically cosmopolitan, N. Am., West Indies, Mid. Am. (not reported Brit. Honduras), S. Am.

Family STRIGIDAE OWLS

Otus flammeolus⁶ FLAMMULATED OWL
Mts. w. N. Am., Mexico and Guatemala.

Otus asio COMMON SCREECH-OWL
N. Am. to Central Plateau of Mexico.

Otus vinaceus⁷ VINACEOUS SCREECH-OWL
Pacific coast of Mexico (s.w. Sonora to Guerrero).

Otus trichopsis SPOTTED SCREECH-OWL
Mts. Arizona, Mexico, Guatemala, Honduras and El Salvador.

Otus barbarus BEARDED SCREECH-OWL
Mts. n. Guatemala.

Otus guatemalae⁸ VERMICULATED SCREECH-OWL
Trop. Mid. Am. and S. Am.

Otus cooperi⁹ PACIFIC SCREECH-OWL
Arid Pacific slope s. Mexico (Chiapas) to n.w. Costa Rica.

Otus choliba TROPICAL SCREECH-OWL
Costa Rica, Panama and S. Am.

Otus clarkii BARE-SHANKED¹⁰ SCREECH-OWL
Mts. Costa Rica and w. Panama.

Lophostrix cristata CRESTED OWL
Trop. Mid. Am. (not reported Brit. Honduras and Honduras), and S. Am.

Bubo virginianus GREAT HORNED OWL
N. Am., Mid. Am. (chiefly in mts.), and S. Am.

⁶ By some considered a race of the Old World *O. scops*, Common Scops-Owl.

⁷ By some treated as conspecific with *O. asio*.

⁸ Some consider birds from Costa Rica, Panama and S. Am. to be a separate species, *O. vermiculatus*. On that view the birds from Mexico to Nicaragua may be called Middle American Screech-Owl.

⁹ Van Rossem suggests this is conspecific with *O. choliba*.

¹⁰ There is another *Otus* called "Bare-legged". *Clarkii* was formerly called *nudipes*.

Pulsatrix perspicillata	SPECTACLED OWL
Trop. Mid. Am. and S. Am.	
Glaucidium gnoma	NORTHERN PYGMY-OWL
W. N. Am. and mts. Mexico and Guatemala.	
Glaucidium minutissimum	LEAST PYGMY-OWL
Trop. Mid. Am. (not reported Nicaragua and El Salvador), and S. Am.	
Glaucidium jardinii	ANDEAN PYGMY-OWL
Mts. Costa Rica, Panama and n.w. S. Am.	
Glaucidium brasiliandum	FERRUGINOUS PYGMY-OWL
S.w. U. S., Mid. Am. and S. Am.	
Micrathene whitneyi	ELF OWL
Deserts s.w. U. S. and Mexico.	
Speotyto cunicularia	BURROWING OWL
Local: U. S., Bahamas, Hispaniola, Mid. Am. (not reported Brit. Honduras and Nicaragua), and S. Am.	
Ciccaba virgata	MOTTLED OWL
Trop. Mid. Am. and n. S. Am.	
Ciccaba nigrolineata	BLACK-AND-WHITE OWL
Trop. Mid. Am. and n.w. S. Am.	
Strix occidentalis	SPOTTED OWL
W. N. Am. and mts. w. and n. Mexico from Sonora and Nuevo Leon s. to Michoacan.	
Strix varia	BARRED OWL
E. and s. N. Am. and mts. cent. Mexico from Durango to Oaxaca and Veracruz.	
Strix fulvescens¹	FULVOUS OWL
Mts. s. Mexico (Oaxaca and Chiapas), Guatemala, El Salvador and Honduras.	
Rhinoptynx clamator	STRIPED OWL
Local: trop. Mid. Am. (not reported Brit. Honduras, Honduras) and S. Am.	
Asio otus	LONG-EARED OWL
Eurasia, N. Am., n.w. Mexico (n. Baja Calif.); winters to central Mexico.	
Asio stygius	STYGIAN OWL
Local: Greater Antilles, mts. Mexico, Guatemala, Nicaragua, S. Am.	
Asio flammeus	SHORT-EARED OWL
N. Eurasia, N. Am., Greater Antilles, S. Am.; northern birds winter to Mexico and Guatemala.	

¹ By some considered a race of *S. varia*.

Aegolius acadicus	NORTHERN SAW-WHET OWL
N. Am. and mts. of Mexico to Veracruz and Oaxaca.	
Aegolius ridgwayi	UNSPOTTED SAW-WHET OWL
Highlands s. Mexico (Chiapas), Guatemala, El Salvador, Costa Rica.	

Order CAPRIMULGIFORMES

Family NYCTIBIIDAE POTOOS

Nyctibius grandis	GREAT POTOO
Cent. Panama s. to S. Am.	
Nyctibius griseus	COMMON POTOO
Greater Antilles, trop. Mid. Am. and S. Am.	

Family CAPRIMULGIDAE NIGHTJARS

Lurocalis semitorquatus	SEMI-COLLARED NIGHTHAWK
Local: Nicaragua, Panama and S. Am.	
Chordeiles acutipennis	LESSER NIGHTHAWK
S.w. U. S., Mexico, Brit. Honduras, Guatemala, Honduras (Bay Is.?), Nicaragua, S. Am.; migrates through Mid. Am.; winters s. to S. Am. Apparently present year-round in Panama, possibly breeding.	
Chordeiles minor	COMMON NIGHTHAWK
N. Am., West Indies, Mexico (s. to Veracruz, Chiapas); migrates through Mid. Am.; winters in S. Am.	
Nyctidromus albicollis	PAURAQUE
S.w. U. S., Mid. Am. and S. Am.	
Phalaenoptilus nuttallii	COMMON POORWILL
W. N. Am. and w. Mexico.	
Otophanes mcleodii	EARED POORWILL
Mexico (Chihuahua, Jalisco and Guerrero).	
Otophanes yucatanicus	YUCATAN POORWILL
Mexico (Yucatan Pen.) and Guatemala (Peten).	
Nyctiphrynus ocellatus	OCELLATED POORWILL
Nicaragua (one spec.), and S. Am.	
Caprimulgus carolinensis	CHUCK-WILL'S-WIDOW
S.e. U. S.; winters to Greater Antilles, through Mid. Am. (not reported Brit. Honduras), and Colombia.	
Caprimulgus rufus	RUFOUS NIGHTJAR
Costa Rica, Panama, St. Lucia, and S. Am.	

<i>Caprimulgus salvini</i> ²	TAWNY-COLLARED NIGHTJAR
E. Trop. Mexico, Brit. Honduras and n. Nicaragua.	
<i>Caprimulgus ridgwayi</i>	BUFF-COLLARED NIGHTJAR
W. Mexico, Guatemala and Honduras.	
<i>Caprimulgus vociferus</i>	WHIP-POOR-WILL
E. N. Am. and in mts. of Mexico, Guatemala, El Salvador, Honduras; northern birds winter through Mid. Am. to Costa Rica.	
<i>Caprimulgus saturatus</i>	DUSKY NIGHTJAR
Mts. Costa Rica and w. Panama.	
<i>Caprimulgus cayennensis</i>	WHITE-TAILED NIGHTJAR
Costa Rica, Panama and S. Am.	
<i>Caprimulgus maculicaudus</i>	SPOT-TAILED NIGHTJAR
Recorded Mexico (Oaxaca, Veracruz and Chiapas) and S. Am.	

Order APODIFORMES

Family APODIDAE SWIFTS

<i>Streptoprocne zonaris</i>	WHITE-COLLARED SWIFT
Highlands Greater Antilles, Mid. Am., and S. Am.	
<i>Streptoprocne semicollaris</i>	WHITE-NAPED SWIFT
Mts. w. and cent. Mexico; seen Chiapas (M. Alvarez del Toro).	
<i>Chaetura chapmani</i>	DARK-BREASTED SWIFT
N. S. Am.; recorded in Panama.	
<i>Chaetura pelasgica</i>	CHIMNEY SWIFT
E. N. Am.; migrates through Mid. Am. (not reported Honduras and El Salvador); winters in S. Am.	
<i>Chaetura vauxi</i>	VAUX'S SWIFT
N. w. N. Am. s. to cent. Calif.; winters to Louisiana, Mexico and Guatemala.	
<i>Chaetura richmondi</i> ³	DUSKY-BACKED SWIFT
Trop. Mid. Am. (except Brit. Honduras) and n. S. Am.	
<i>Chaetura gaumeri</i> ³	YUCATAN SWIFT
Mexico (Yucatan Pen. and Cozumel I.).	
<i>Chaetura cinereiventris</i>	GRAY-RUMPED SWIFT
Caribbean slope Nicaragua, Costa Rica, w. Panama, S. Am., Lesser Antilles (Grenada).	
<i>Chaetura spinicauda</i> ⁴	BAND-RUMPED SWIFT
Costa Rica, Panama and S. Am.	

² Considered by Peters conspecific with *C. serico-caudatus*, of uncertain origin.

³ Treated by many as races of *C. vauxi*, Vaux's Swift (Sutton, Wilson Bull. 53: 231-233, 1941), but the distance from the breeding range of that form raises doubt.

⁴ Includes *C. fumosa* of Costa Rica and Panama.

Chaetura andrei	ASHY-TAILED SWIFT
S. Am.; recorded once from Panama:	
Cypseloides⁵ rutilus	CHESTNUT-COLLARED SWIFT
Mts. w. and s. Mexico, Guatemala, Costa Rica, Panama and S. Am.	
Cypseloides cherriei	SPOT-FRONTED SWIFT
Recorded from Costa Rica, Colombia and Venezuela.	
Cypseloides cryptus	WHITE-CHINNED SWIFT
Recorded from Nicaragua, Costa Rica, Panama and n. S. Am.	
Cypseloides niger	BLACK SWIFT
W. N. Am., mts. Mexico, Guatemala, Honduras, Nicaragua, Costa Rica, West Indies.	
Aeronautes saxatalis	WHITE-THROATED SWIFT
Mts. w. N. Am., Mexico, Guatemala, Honduras and El Salvador	
Panyptila sancti-hieronymi⁶	GREAT SWALLOW-TAILED SWIFT
Mts. s. Mexico (Michoacan, Chiapas), w. Guatemala, Honduras, w. Nicaragua (near Jalapa, <i>fide</i> T. R. Powell, G. Montrello).	
Panyptila cayennensis	LESSER SWALLOW-TAILED SWIFT
Mexico (Veracruz), Caribbean slope, Honduras, Nicaragua, Costa Rica, Panama (both slopes) and S. Am.	

Family TROCHILIDAE HUMMINGBIRDS

Doryfera ludovicae	GREEN-FRONTED LANCEBILL
Highlands Costa Rica, w. Panama and S. Am.	
Androdon aequatorialis	TOOTH-BILLED HUMMINGBIRD
E. Panama and Pac. slope Colombia and Ecuador.	
Glaucis aenea⁷	BRONZY HERMIT
Nicaragua, Costa Rica (both slopes) and Pac. slope w. Panama (Chiriqui), w. Colombia, n.w. Ecuador.	
Glaucis hirsuta	RUFOUS-BREASTED HERMIT
Cent. and e. Panama through most of trop. S. Am., s. Lesser Antilles (Grenada).	
Threnetes ruckeri	BAND-TAILED BARBTHROAT
Nicaragua, Costa Rica, Panama and n.w. S. Am.	
Phaethornis guy	GREEN HERMIT
Costa Rica, Panama and n. S. Am.	

⁵ Peters puts this in *Chaetura*, and some authors in a separate genus *Chaeturella*; Zimmer is here followed.

⁶ Stresemann suggests that this may be a race of *P. cayennensis*.

⁷ Perhaps a race of *G. hirsuta*, as Peters treats it.

Phaethornis superciliosus⁸	LONG-TAILED HERMIT
Trop. Mid. Am. and S. Am.	
Phaethornis anthophilus	PALE-BELLIED HERMIT
Panama (Pearl Is.), Colombia and w. Venezuela.	
Phaethornis longuemareus	LITTLE HERMIT
Trop. Mid. Am. (except El Salvador), and S. Am.	
Eutoxeres aquila	WHITE-TIPPED SICKLEBILL
Costa Rica, Panama and n.w. S. Am.	
Phaeochroa cuvierii	SCALY-BREASTED HUMMINGBIRD
Guatemala, Brit. Honduras, Honduras, Nicaragua, Costa Rica, Panama, n. Colombia.	
Campylopterus curvipennis⁹	WEDGE-TAILED SABREWING
Trop. e. Mexico and Guatemala.	
Campylopterus rufus	RUFOUS SABREWING
Highlands s. Mexico (Chiapas), Guatemala and El Salvador.	
Campylopterus hemileucurus	VIOLET SABREWING
Highlands s. Mexico through Mid. Am. to w. Panama.	
Florisuga mellivora	WHITE-NECKED JACOBIN
Trop. Mid. Am. (except El Salvador), and S. Am.	
Colibri delphinae	BROWN VIOLET-EAR
Guatemala, Brit. Honduras, Honduras, Nicaragua, Costa Rica, Panama and S. Am.	
Colibri thalassinus¹⁰	GREEN VIOLET-EAR
Highlands Mid. Am. (not reported Brit. Honduras, Nicaragua) and S. Am.	
Anthracothorax prevostii	GREEN-BREASTED MANGO
Trop. Mid. Am. (except Panama) to n. Costa Rica, n.w. Venezuela, Old Providence and St. Andrew's Is. in Caribbean.	
Anthracothorax veraguensis¹¹	VERAGUAN MANGO
W. Panama (Pac. slope Chiriqui and Veraguas).	
Anthracothorax nigricollis	BLACK-THROATED MANGO
Cent. Panama to S. Am.	
Klais guimeti	VIOLET-HEADED HUMMINGBIRD
Highlands Honduras, Nicaragua, Costa Rica, Panama and w. S. Am.	

⁸ Zimmer considers the Middle American and most South American forms to be races of *P. malaris* (of Cayenne), and not of *P. superciliosus*. The English name given is intended to apply to the Middle American birds, regardless of their technical name.

⁹ The form from s. Veracruz, *C. excellens*, Long-tailed Sabrewing, may be a distinct species.

¹⁰ Includes the *C. cyanotus* group of Costa Rica to S. Am., regarded by some as a separate species, Mountain Violet-ear.

¹¹ Possibly a race of *A. prevostii*.

Abeillia abeillei	EMERALD-CHINNED HUMMINGBIRD
Mts. s.e. Mexico, Guatemala, El Salvador, Honduras, and n. Nicaragua.	
Lophornis delattrei	RUFOUS-CRESTED COQUETTE
Mexico (s.w. Guerrero), s.w. Costa Rica, Panama and w. S. Am.	
Paphosia² helena	BLACK-CRESTED COQUETTE
Caribbean slope trop. Mid. Am. to Costa Rica.	
Paphosia adorabilis	WHITE-CRESTED COQUETTE
Cent. and s.w. Costa Rica to s.w. Panama.	
Popelairia conversii	GREEN THORNTAIL
Costa Rica, Panama, Colombia and Ecuador.	
Chlorostilbon canivetii³	FORK-TAILED EMERALD
Trop. Mid. Am. and Caribbean coast of S. Am.	
Cynanthus sordidus	DUSKY HUMMINGBIRD
S.w. Mexico.	
Cynanthus latirostris	BROAD-BILLED HUMMINGBIRD
S.w. U. S. and Mexico.	
Thalurania colombica⁴	BLUE-CROWNED WOODNYMPH
Trop. Mid. Am. (except El Salvador), in Mexico reported only from Jalisco; also Colombia.	
Thalurania fannyi⁴	GREEN-CROWNED WOODNYMPH
E. Panama and w. Colombia and Ecuador.	
Panterpe insignis	FIERY-THROATED HUMMINGBIRD
Mts. Costa Rica and w. Panama.	
Damophila⁵ julie	VIOLET-BELLIED HUMMINGBIRD
Costa Rica, Panama, Colombia and Ecuador.	
Lepidopyga coeruleogularis	SAPPHIRE-THROATED HUMMINGBIRD
Costa Rica, Panama and n. Colombia.	
Hylocharis xanthusii	BLACK-FRONTED HUMMINGBIRD
Mexico (s. Baja Calif.).	

² This genus (and sometimes also *Popelairia*) is often merged in the genus *Lophornis*.

³ Birds from s.w. Costa Rica and Panama are often regarded as a separate species, *C. assimilis*, Garden Emerald. If so treated, the South American population must also be separated from *C. canivetii*. Zimmer lumps all these forms with *C. mellisugus* of Cayenne; if this view is followed Common Emerald is the best name, as typical *mellisugus* has the tail truncate, not forked.

⁴ Considered by Peters, Zimmer and Blake to be races of *T. furcata*, Common Woodnymph, of e. S. Am. Some authors lump only the glittering-crowned forms (including *fannyi*), of Mid. Am. and n.w. Colombia and Ecuador, in *T. colombica*, which may be called Crowned Woodnymph.

⁵ *Juliamyia juliae* of some authors.

Hylocharis leucotis	WHITE-EARED HUMMINGBIRD
Highlands and mts. s. Arizona, Mexico, Guatemala, El Salvador, Honduras and Nicaragua.	
Hylocharis eliciae	BLUE-THROATED GOLDENTAIL
Trop. Mid. Am.	
Hylocharis grayi	BLUE-HEADED SAPPHIRE
E. Panama (Darien) and n.w. S. Am.	
Goldmania violiceps	VIOLET-CAPPED HUMMINGBIRD
E. Panama.	
Goethalsia bella	RUFous-CHEEKED HUMMINGBIRD
E. Panama (Darien).	
Amazilia candida	WHITE-BELLIED EMERALD
Trop. Mid. Am. (except El Salvador) to Costa Rica.	
Amazilia luciae	HONDURAS EMERALD
Honduras.	
Amazilia amabilis⁶	BLUE-CHESTED HUMMINGBIRD
Nicaragua, Costa Rica, Panama, Colombia and Ecuador.	
Amazilia boucardi	MANGROVE HUMMINGBIRD
Costa Rica (Pac. coast).	
Amazilia cyanocephala	RED-BILLED AZURECROWN
Mts. s.e. Mexico, Brit. Honduras, Guatemala, El Salvador, Hon- duras, Nicaragua.	
Amazilia microrhyncha	SMALL-BILLED AZURECROWN
Honduras ?. Known only from the type.	
Amazilia cyanifrons	BLUE-FRONTED HUMMINGBIRD
Costa Rica (one spec.), Colombia.	
Amazilia beryllina	BERYLLINE HUMMINGBIRD
Mexico, Guatemala, El Salvador and Honduras.	
Amazilia cyanura	BLUE-TAILED HUMMINGBIRD
Pac. slope s. Mexico (Chiapas), Guatemala, El Salvador, Hon- duras, Nicaragua; once in Costa Rica.	
Amazilia saucerrottei	BLUE-VENTED HUMMINGBIRD
W. and s. Nicaragua and Costa Rica, also Colombia and Vene- zuela.	
Amazilia edward⁷	SNOWY-BREASTED HUMMINGBIRD
Costa Rica and Panama.	

⁶ Includes *A. decora*, Charming Hummingbird, Pacific slope s.w. Costa Rica and w. Panama, regarded by some as a distinct species.

⁷ Includes *niveoventer* of Costa Rica and w. Panama, by some held to be a species distinct from *A. edward*. The *edward* group has been called White-bellied Hummingbird.

Amazilia rutila	CINNAMON HUMMINGBIRD
Mid. Am. from s. Mexico (Chiapas) to cent. Costa Rica.	
Amazilia yucatanensis	FAWN-BREASTED HUMMINGBIRD
S. Texas, Mexico, Guatemala and Brit. Honduras.	
Amazilia tzacatl	RUFOUS-TAILED HUMMINGBIRD
Trop. Mid. Am. (except El Salvador), Colombia and Ecuador.	
Amazilia violiceps⁸	VIOLET-CROWNED HUMMINGBIRD
W. and s. Mexico (Sonora and Chihuahua to Chiapas); accidental in Arizona.	
Amazilia viridifrons⁸	GREEN-FRONTED HUMMINGBIRD
S. Mexico (Guerrero to Chiapas).	
Eupherusa eximia⁹	STRIPE-TAILED HUMMINGBIRD
Highlands Mid. Am. (except El Salvador) from s. Mexico to w. Panama.	
Eupherusa nigriventris	BLACK-BELLIED HUMMINGBIRD
Highlands Costa Rica and w. Panama.	
Elvira chionura	WHITE-TAILED EMERALD
Highlands s.w. Costa Rica and w. Panama.	
Elvira cupreiceps	COPPERY-HEADED EMERALD
Caribbean slope Costa Rica.	
Microchera albo-coronata	SNOWCAP
Caribbean slope Nicaragua, Costa Rica and w. Panama.	
Chalybura buffonii	WHITE-VENTED PLUMELETEER
Cent. Panama to n.w. S. Am.	
Chalybura melanorrhao¹⁰	DUSKY PLUMELETEER
Caribbean slope Nicaragua, Costa Rica and extreme n.w. Panama.	
Chalybura urochrysia	BRONZE-TAILED PLUMELETEER
Panama and n.w. S. Am.	
Lampornis clemenciae	BLUE-THROATED HUMMINGBIRD
Mts. s.w. U. S. and Mexico.	
Lampornis amethystinus	AMETHYST-THROATED HUMMINGBIRD
Mts. trop. Mexico, Guatemala, El Salvador and Honduras.	
Lampornis viridi-pallens	GREEN-THROATED MOUNTAIN-GEM
Mts. s. Mexico (Chiapas), Guatemala, El Salvador, Honduras, Nicaragua.	
Lampornis hemileucus	WHITE-BELLIED MOUNTAIN-GEM
Highlands Caribbean slope Costa Rica and w. Panama.	

⁸ Peters regards *viridifrons* as the immature plumage of *violiceps*. An earlier name of *violiceps* may be *verticalis* (Stresemann, Condor, 1954; 91).

⁹ Includes *E. poliocerca*, White-tailed Hummingbird, of s.w. Mexico, which may be entitled to specific status.

¹⁰ This may be a race of *C. urochrysia*.

Lampornis calolaema ¹	PURPLE-THROATED MOUNTAIN-GEM
Highlands w. Nicaragua, Costa Rica and w. Panama (e. to Veraguas).	
Lampornis castaneoventris ¹	WHITE-THROATED MOUNTAIN-GEM
Highlands w. Panama (w. Chiriqui).	
Lampornis cinereicauda ¹	GRAY-TAILED MOUNTAIN-GEM
Mts. Costa Rica and (?) w. Panama (Volcan de Chiriqui, <i>fide</i> Ridgway, but Peters <i>dubitante</i>).	
Lamprolaima rhami	GARNET-THROATED HUMMINGBIRD
Mts. s. Mexico, Guatemala, El Salvador, Honduras.	
Heliodoxa jacula	GREEN-CROWNED BRILLIANT
Highlands Costa Rica, Panama, Colombia and Ecuador.	
Eugenes fulgens ²	MAGNIFICENT HUMMINGBIRD
Mts. s.w. U. S. and Mid. Am. to w. Panama.	
Haplophaedia aureliae	GREENISH PUFFLEG
Highlands e. Panama (Darien) and w. S. Am.	
Heliothrix barroti ³	PURPLE-CROWNED FAIRY
Trop. Mid. Am. (except Mexico and El Salvador) to Colombia and Ecuador.	
Heliomaster constanti	PLAIN-CAPPED STARTHROAT
Pac. coast trop. Mid. Am. to Costa Rica.	
Heliomaster longirostris	LONG-BILLED STARTHROAT
Trop. Mid. Am. (except Brit. Honduras), and S. Am.	
Philodice bryantae	MAGENTA-THROATED WOODSTAR
Highlands Costa Rica and w. Panama.	
Doricha enicura	SLENDER SHEARTAIL
Highlands s. Mexico (Chiapas), Guatemala, El Salvador and Honduras.	
Doricha eliza	MEXICAN SHEARTAIL
S. e. Mexico (Veracruz and Yucatan Pen.).	

¹ The relationship among these forms (sometimes segregated in a genus *Oreopyra*) is uncertain. Berlioz suggests that all three are partly localized color phases of one species. Most authors treat *calolaema* as a subspecies of *castaneoventris*; yet both occur together in w. Chiriqui. Hartert and Blake suggest that the subspecies of *castaneoventris* is *cinereicauda*. On that view, occasional Costa Rican birds of *castaneoventris* type might be explained as hybrids between *calolaema* and *cinereicauda*, or as mutants of the latter. If two or more of these forms is included in *castaneoventris*, it may be best to call the complex Chestnut-bellied Mountain-gem.

² Includes *E. spectabilis* of Costa Rica and Panama, called the Admirable Hummingbird. The northern form has been called Rivoli's Hummingbird.

³ Zimmer considers this a race of the green-crowned South American *H. aurita*, Black-eared Fairy, which may be used for the entire complex.

Tilmatura dupontii	SPARKLING-TAILED HUMMINGBIRD
Highlands (chiefly) s. and cent. Mexico, Guatemala, El Salvador, Honduras and Nicaragua.	
Calothrax lucifer	LUCIFER HUMMINGBIRD
S.w. U. S. and highlands of s. cent. and s. Mexico.	
Calothrax pulcher	BEAUTIFUL HUMMINGBIRD
S. Mexico (Guerrero to Chiapas).	
Archilochus colubris	RUBY-THROATED HUMMINGBIRD
E. N. Am.; winters through Mid. Am.	
Archilochus alexandri	BLACK-CHINNED HUMMINGBIRD
W. N. Am. to n. Mexico; winters through Mexico.	
Calypte anna	ANNA'S HUMMINGBIRD
Calif. and n. Mexico (n. Baja Calif.); winters to w. Mexico.	
Calypte costae	COSTA'S HUMMINGBIRD
W. U. S. and n.w. Mexico (Baja Calif. and Sonora); winters to n.w. Mexico (Sinaloa).	
Stellula calliope	CALLIOPE HUMMINGBIRD
Mts. w. N. Am. and w. Mexico (Baja Calif. and Guerrero); winters in Mexico.	
Atthis heloisa	BUMBLEBEE HUMMINGBIRD
Mts. Mexico (Sinaloa, Chihuahua, Nuevo Leon and Tamaulipas to Oaxaca and Veracruz); accidental in Arizona.	
Atthis ellioti⁴	WINE-THROATED HUMMINGBIRD
Mts. s. Mexico (Chiapas), Guatemala and Honduras.	
Acestrura heliodor	GORGED WOODSTAR
Highlands e. Panama (Cana, Darien one spec. <i>fide</i> J. Bond) and n.w. S. Am.	
Selasphorus platycercus	BROAD-TAILED HUMMINGBIRD
Mts. w. U. S., Mexico and Guatemala.	
Selasphorus rufus	RUFOUS HUMMINGBIRD
W. N. Am.; winters to Mexico.	
Selasphorus sasin	ALLEN'S HUMMINGBIRD
Calif.; winters to n.w. and cent. Mexico.	
Selasphorus flammula	ROSE-THROATED HUMMINGBIRD
High mts. Costa Rica.	
Selasphorus torridus⁵	HELIOTROPE-THROATED HUMMINGBIRD
High mts. Costa Rica and w. Panama (Volcan de Chiriqui).	
Selasphorus simoni⁶	CERISE-THROATED HUMMINGBIRD
Highlands of Costa Rica.	

⁴ Frequently considered conspecific with *A. heloisa*.

⁵ By some believed to be a color phase of *S. flammula*. If this be so, the species may be known as the Volcano Hummingbird.

⁶ Possibly a race of *S. ardens*.

Selasphorus ardens	GLOW-THROATED HUMMINGBIRD
Highlands w. Panama.	
Selasphorus scintilla	SCINTILLANT HUMMINGBIRD
Highlands Costa Rica and w. Panama.	

Order TROGONIFORMES

Family TROGONIDAE TROGONS

Pharomachrus mocinno	QUETZAL
Mts. Mid. Am. (except Brit. Honduras), from Mexico (Oaxaca, Chiapas) to w. Panama.	
Pharomachrus auriceps⁷	GOLDEN-HEADED TROGON
Mts. e. Panama (Darien), w. and n. S. Am.	
Euptilotis neoxenus	EARED TROGON
Mts. central Mexico (Chihuahua to Michoacan).	
Trogon massena	SLATY-TAILED TROGON
Trop. Mid. Am., Colombia and Ecuador.	
Trogon clathratus	LATTICE-TAILED TROGON
Caribbean coast Costa Rica and w. Panama.	
Trogon melanurus	BLACK-TAILED TROGON
Cent. Panama to S. Am.	
Trogon viridis⁸	WHITE-TAILED TROGON
S.w. Costa Rica, Panama and S. Am.	
Trogon citreolus	CITREOLINE TROGON
Trop. Mid. Am. to n. Costa Rica.	
Trogon mexicanus	MOUNTAIN TROGON
Highlands Mexico, Guatemala, Honduras.	
Trogon elegans⁹	ELEGANT TROGON
S.w. U. S., Mid. Am. (not reported Brit. Honduras) to n.w. Costa Rica.	
Trogon collaris	BAR-TAILED TROGON
Highlands trop. Mid. Am. and S. Am.	
Trogon aurantiventris	ORANGE-BELLIED TROGON
Highlands Costa Rica and w. Panama.	
Trogon rufus	BLACK-THROATED TROGON
S.e. Honduras, Nicaragua, Costa Rica, Panama and S. Am.	

⁷ By some deemed a race of the Amazonian *P. pavoninus*, Pavonine Trogon.

⁸ *T. strigilatus* of some authorities; includes *bairdi*, Baird's Trogon, of s.w. Costa Rica and s.w. Panama.

⁹ Includes *T. ambiguus*, the Coppery-tailed Trogon of Mexico; Cent. American birds lack any coppery tail color.

Trogon violaceus VIOLACEOUS TROGON
Trop. Mid. Am. and S. Am.

Order CORACIIFORMES
Family ALCEDINIDAE KINGFISHERS

Ceryle¹⁰ torquata	RINGED KINGFISHER
Mid. Am., Lesser Antilles, and S. Am.	
Ceryle alcyon	BELTED KINGFISHER
N. Am.; winters through Mid. Am. to n. S. Am.	
Chloroceryle amazona	AMAZON KINGFISHER
Trop. Mid. Am. and S. Am.	
Chloroceryle americana	GREEN KINGFISHER
S.w. U. S., Mid. Am. and S. Am.	
Chloroceryle indica	GREEN-AND-RUFOUS KINGFISHER
Nicaragua, Costa Rica, Panama and S. Am.	
Chloroceryle aenea	PYGMY KINGFISHER
Trop. Mid. Am. and S. Am.	

Family MOMOTIDAE MOTMOTS

Hylomanes momotula	TODY MOTMOT
Trop. Mid. Am. and n.w. Colombia.	
Aspatha gularis	BLUE-THROATED MOTMOT
Highlands s. Mexico (Chiapas), Guatemala, El Salvador and Honduras.	
Electron platyrhynchum	BROAD-BILLED MOTMOT
E. Honduras, Nicaragua, Costa Rica, Panama and S. Am.	
Electron carinatum	KEEL-BILLED MOTMOT
Caribbean slope Mid. Am. from s. Mexico to n. Costa Rica.	
Eumomota superciliosa	TURQUOISE-BROWED MOTMOT
Trop. Mid. Am. (not reported Brit. Honduras) to n.w. Costa Rica.	
Baryphthengus ruficapillus	RUFOUS MOTMOT
Nicaragua, Costa Rica, Panama and S. Am.	
Momotus mexicanus	RUSSET-CROWNED MOTMOT
W. Mexico (Sonora to Chiapas) and Guatemala (arid interior).	
Momotus momota¹	BLUE-CROWNED MOTMOT
Trop. Mid. Am. and S. Am.	

¹⁰ This and the next species are often placed in a separate genus, *Megaceryle*.

¹ Birds from the Pacific slope of cent. and e. Panama to n. Venezuela are by some regarded as a distinct species, *M. subrufescens*, the Tawny-bellied Motmot. Birds from Mexico to w. Panama belong to the *M. lessoni* group.

Order PICIFORMES

Family GALBULIDAE JACAMARS

Brachygalba salmoni	DUSKY-BACKED JACAMAR
E. Panama (Darien) and n.w. Colombia.	
Galbula ruficauda²	RUFOUS-TAILED JACAMAR
Trop. Mid. Am. (except El Salvador) to w. Panama, and S. Am.	
Jacamerops aurea	GREAT JACAMAR
Carribean slope Costa Rica, Panama (both slopes) and S. Am.	

Family BUCCONIDAE PUFFBIRDS

Notharchus macrorhynchos	WHITE-NECKED PUFFBIRD
Trop. Mid. Am. and S. Am.	
Notharchus pectoralis	BLACK-BREASTED PUFFBIRD
Cent. Panama to n.w. S. Am.	
Notharchus tectus	PIED PUFFBIRD
Costa Rica, Panama, to S. Am.	
Nystalus radiatus	BARRED PUFFBIRD
Cent. Panama to n.w. S. Am.	
Malacoptila panamensis	WHITE-WHISKERED PUFFBIRD
Trop. Mid. Am. (except El Salvador) and S. Am.	
Micromonacha lanceolata	LANCEOLATED MONKLET
Rare: Costa Rica, Panama and S. Am.	
Nonnula frontalis³	GRAY-CHEEKED NUNLET
Cent. Panama to Colombia.	
Monasa morphoeus	WHITE-FRONTED NUNBIRD
Nicaragua, Costa Rica, Panama and S. Am.	

Family CAPITONIDAE BARBETS

Capito maculicoronatus	SPOT-CROWNED BARBET
Cent. Panama to Colombia.	
Eubucco bourcierii	RED-HEADED BARBET
Highlands Costa Rica, Panama and n.w. S. Am.	

Semnornis frantzii	PRONG-BILLED BARBET
Highlands Costa Rica and w. Panama.	

² Includes *melanogenia*, the Middle American form ranging to w. Ecuador, formerly regarded as a separate species, Black-chinned Jacamar.

³ Considered by de Schauensee conspecific with *N. ruficapilla*, ranging to Peru.

Family RAMPHASTIDAE TOUCANS

Aulacorhynchus prasinus	EMERALD TOUCANET
Highlands s. Mexico, Guatemala, Brit. Honduras, Honduras, El Salvador and Nicaragua.	
Aulaeorhynchus caeruleogularis⁴	BLUE-THROATED TOUCANET
Highlands of Costa Rica and Panama.	
Pteroglossus torquatus	COLLARED ARAÇARI
Trop. Mid. Am., Colombia and Venezuela.	
Pteroglossus frantzii⁵	FIERY-BILLED ARAÇARI
Pacific slope Costa Rica and w. Panama.	
Selenidera spectabilis	YELLOW-EARED TOUCANET
Honduras, Nicaragua, Costa Rica, Panama and n. w. Colombia.	
Ramphastos sulfuratus	KEEL-BILLED TOUCAN
Trop. Mid. Am. (except El Salvador), and n. S. Am.	
Ramphastos swainsonii	CHESTNUT-MANDIBLED TOUCAN
Honduras, Nicaragua, Costa Rica, Panama and n.w. S. Am.	
Ramphastos ambiguus	BLACK-MANDIBLED TOUCAN
E. Panama and n. S. Am.	

Family PICIDAE WOODPECKERS

Picumnus olivaceus	OLIVACEOUS PICULET
E. Honduras, Nicaragua, Costa Rica, Panama and n. S. Am.	
Colaptes cafer⁶	RED-SHAFTED FLICKER
W. N. Am. and highlands Mexico, Guatemala, El Salvador, Honduras, n. Nicaragua.	
Colaptes chrysoides⁶	GILDED FLICKER
Deserts s.w. U. S. and n.w. Mexico (Baja Calif., Sonora and n. Sinaloa).	
Chrysotilus punctigula	SPOT-BREASTED WOODPECKER
E. Panama (Darien) and S. Am.	
Piculus auricularis	GRAY-CROWNED WOODPECKER
W. Mexico (Sonora to Guerrero).	
Piculus aeruginosus	BRONZE-WINGED WOODPECKER
Trop. n.e. Mexico (s. Nuevo Leon and Tamaulipas to Veracruz and Puebla).	

⁴ Often considered conspecific with *A. prasinus*, as well as certain S. Am. forms; on that view call the complex Emerald Toucanet.

⁵ Peters treats it as a race of *P. torquatus*.

⁶ All North American Flickers may be races of *C. auratus*; if so, call Common Flicker.

Piculus rubiginosus	GOLDEN-OLIVE WOODPECKER
Trop. Mid. Am. (in much of its range chiefly in highlands), and S. Am.	
Piculus simplex	RUFOUS-WINGED WOODPECKER
Honduras, Nicaragua, Costa Rica and w. Panama.	
Piculus callopterus⁷	STRIPE-CHEEKED WOODPECKER
Cent. and e. Panama.	
Piculus chrysochloros	GOLDEN-GREEN WOODPECKER
E. Panama (Darien) to S. Am.	
Celeus castaneus	CHESTNUT-COLORED WOODPECKER
Trop. s.e. Mexico and Caribbean slope of Mid. Am. to n.w. Panama.	
Celeus immaculatus	IMMACULATE WOODPECKER
Panama (?). Unique spec. believed by describer to be of Panama "make".	
Celeus loricatus	CINNAMON WOODPECKER
S.e. Nicaragua, Costa Rica, Panama and n.w. S. Am.	
Dryocopus lineatus	LINEATED WOODPECKER
Trop. Mid. Am. and S. Am.	
Asyndesmus lewisi	LEWIS'S WOODPECKER
W. N. Am.; winters to n.w. Mexico (Baja Calif. and n. Sonora).	
Melanerpes formicivorus	ACORN WOODPECKER
W. U. S. and highlands of Mid. Am. to w. Panama, also Colombia.	
Centurus⁸ uropygialis⁹	GILA OR DESERT WOODPECKER
S.w. U. S. (deserts) and w. Mexico (south to Jalisco).	
Centurus hypoleucus	GRAY-BREASTED WOODPECKER
Arid s.w. and cent. Mexico (Mexico to Guerrero and Oaxaca).	
Centurus aurifrons	GOLDEN-FRONTED WOODPECKER
Texas and Mid. Am. to Costa Rica.	
Centurus chrysogenys	GOLDEN-CHEEKED WOODPECKER
W. Mexico.	
Centurus rubricomus¹⁰	RED-VENTED WOODPECKER
Mexico (Yucatan Pen. and Cozumel I.), Honduras (Bonacca I.).	
Centurus rubricapillus	RED-CROWNED WOODPECKER
S.w. Costa Rica, Panama and n. S. Am.	

⁷ Peters treats this as a race of *leucolaemus*, a west Amazonian species. Yet *callopterus* seems much more closely allied to *P. simplex*, which Peters keeps as a separate species.

⁸ Peters merges *Centurus* (including *Tripsurus*) in *Melanerpes*.

⁹ Peters and Blake treat this bird as conspecific with the Mexican *hypoleucus*.

¹⁰ Regarded by Peters as conspecific with the next species. The old specific name, *ruberiventris*, must be replaced by *rubricomus* if *Centurus* and *Tripsurus* are merged in one genus.

Centurus¹ pucherani	BLACK-CHEEKED WOODPECKER
Trop. Mid. Am. (except El Salvador), and n.w. S. Am.	
Centurus¹ chrysauchen	GOLDEN-NAPED WOODPECKER
S.w. Costa Rica and s.w. Panama. ²	
Sphyrapicus varius	YELLOW-BELLIED SAPSUCKER
N. Am.; winters through Mid. Am. to Panama.	
Sphyrapicus ruber³	RED-BREASTED SAPSUCKER
Pac. coast of N. Am.; winters to n. Mexico (Baja Calif.).	
Sphyrapicus thyroides	WILLIAMSON'S SAPSUCKER
W. N. Am.; winters to w. Mexico.	
Veniliornis fumigatus	SMOKY-BROWN WOODPECKER
Trop. Mid. Am. and S. Am.	
Veniliornis kirkii	RED-RUMPED WOODPECKER
S.w. Costa Rica, Panama and n. S. Am.	
Dendrocopos villosus	HAIRY WOODPECKER
N. Am., Bahamas, mts. Mid. Am. (except Brit. Honduras) to w. Panama.	
Dendrocopos nuttallii	NUTTALL'S WOODPECKER
Calif. and Mexico (n.w. Baja Calif.).	
Dendrocopos scalaris	LADDER-BACKED WOODPECKER
Arid s.w. U. S., Mexico, Brit. Honduras, and Honduras.	
Dendrocopos arizonae⁴	BROWN-BACKED WOODPECKER
Mts. s.w. U. S. and w. Mexico (Sonora and Chihuahua to Michoacan).	
Dendrocopos stricklandi	BROWN-BARRED WOODPECKER
Mts. cent. and s.e. Mexico (Michoacan to Veracruz).	
Phloeoceastes guatemalensis	PALE-BILLED WOODPECKER
Trop. Mid. Am. to w. Panama.	
Phloeoceastes melanoleucus	CRIMSON-CRESTED WOODPECKER
Cent. Panama to S. Am.	
Phloeoceastes haematogaster⁵	CRIMSON-BELLIED WOODPECKER
Panama and w. S. Am.	
Campephilus imperialis	IMPERIAL WOODPECKER
Mexico (Sierra Madre Occidental, from Sonora and Chihuahua to Michoacan).	

¹ These species were formerly placed in a separate genus, *Tripsurus*.

² The Colombian *T. pulcher* is by some deemed a race of *chrysauchen*.

³ Often treated as conspecific with *S. varius*.

⁴ J. Davis suggests that *arizonae* is conspecific with *D. stricklandi*. If so, Brown-backed Woodpecker may be used for both.

⁵ The Panama and lowland S. Am. form *splendens* is by some considered a distinct species, Splendid Woodpecker.

Order PASSERIFORMES*

Family DENDROCOLAPTIDAE WOODCREEPERS⁷

<i>Dendrocincla fuliginosa</i> ⁸	PLAIN-BROWN WOODCREEPER
S.e. Honduras, Nicaragua, Costa Rica, Panama and S. Am.	
<i>Dendrocincla anabatina</i>	TAWNY-WINGED WOODCREEPER
Trop. Mid. Am. (except El Salvador) to w. Panama.	
<i>Dendrocincla homochroa</i>	RUDDY WOODCREEPER
Trop. Mid. Am. (except El Salvador) and Venezuela.	
<i>Deconychura longicauda</i>	LONG-TAILED WOODCREEPER
Nicaragua, s.w. Costa Rica, Panama and S. Am.	
<i>Sittasomus griseicapillus</i>	OLIVACEOUS WOODCREEPER
Trop. Mid. Am. and S. Am.	
<i>Glyphorynchus spirurus</i>	WEDGE-BILLED WOODCREEPER ⁹
Trop. Mid. Am. (except El Salvador), and S. Am.	
<i>Xiphocolaptes promeropirhynchus</i>	STRONG-BILLED WOODCREEPER
Highlands trop. Mid. Am. to w. Panama, also S. Am.	
<i>Dendrocolaptes certhia</i>	BARRED WOODCREEPER
Trop. Mid. Am. and S. Am.	
<i>Dendrocolaptes picumnus</i>	BLACK-BANDED WOODCREEPER
Highlands s. Mexico (Chiapas), Guatemala, Honduras, Costa Rica, w. Panama and S. Am.	
<i>Xiphorhynchus picus</i> ¹⁰	STRAIGHT-BILLED WOODCREEPER
Central Panama (Pacific coast) and S. Am.	
<i>Xiphorhynchus guttatus</i>	BUFF-THROATED WOODCREEPER
Guatemala, Honduras, Nicaragua, Panama and S. Am.	
<i>Xiphorhynchus flavigaster</i>	IVORY-BILLED WOODCREEPER
Trop. Mid. Am. to n.w. Costa Rica.	
<i>Xiphorhynchus striatigularis</i>	STRIPE-THROATED WOODCREEPER
Mexico (Tamaulipas, one spec.).	

* What families should be recognized and how the many genera should be allocated have been much disputed recently (see Mayr and Amadon, Am. Mus. Nov., no. 1496, 1952; Wetmore, Smith. Misc. Coll., 117 no. 4, 1952; Beecher, Auk, 70: 270-333, 1953; Tordoff, Auk, 71: 273-284, 1954).

⁷ The misnomer "Woodhewer," a translation of the technical name, has often been used.

⁸ The name *meruloides* is often used for the Middle American forms, *fuliginosa* of e. S. Am. being regarded as a distinct species (see Todd, Ann. Carn. Mus. 31(2): 17, 1948).

⁹ The unmodified name "Wedge-bill", sometimes used, creates unnecessary confusion with a genus of hummingbirds to which that group-name applies.

¹⁰ The Panama form belongs to the *picrostis* group of arid n. S. Am., sometimes regarded as a separate species. This species is often placed in a separate genus *Dendroplex*.

Xiphorhynchus lachrymosus	BLACK-STRIPED WOODCREEPER
E. Nicaragua, Costa Rica, Panama and S. Am.	
Xiphorhynchus erythropygius¹	SPOTTED WOODCREEPER
Chiefly highlands trop. Mid. Am. (except Brit. Honduras), Colombia and Ecuador.	
Lepidocolaptes leucogaster	WHITE-STRIPED WOODCREEPER
W. and cent. Mexico.	
Lepidocolaptes souleyetii	STREAK-HEADED WOODCREEPER
Trop. Mid. Am. and S. Am.	
Lepidocolaptes affinis	SPOT-CROWNED WOODCREEPER
Highlands trop. Mid. Am. (except Brit. Honduras) to w. Panama. ²	
Campylorhamphus trochilirostris	RED-BILLED SCYTHEBILL ³
Cent. Panama to S. Am.	
Campylorhamphus pusillus	BROWN-BILLED SCYTHEBILL
Costa Rica and w. Panama, also Colombia and Ecuador.	

Family FURNARIIDAE

OVENBIRDS, SPINETAILS AND ALLIES

Synallaxis albescens	PALE-BREASTED SPINETAIL
S.w. Costa Rica, Panama and S. Am.	
Synallaxis brachyura	SLATY SPINETAIL
E. Honduras, e. Nicaragua, Costa Rica, Panama and n.w. S. Am.	
Synallaxis erythrothorax	RUFOUS-BREASTED SPINETAIL
S. Mexico, Guatemala, Brit. Honduras, El Salvador and Honduras.	
Cranioleuca erythrops	RED-FACED SPINETAIL
Highlands Costa Rica, Panama and n.w. S. Am.	
Xenerpestes⁴ minlosi	DOUBLE-BANDED SOFTTAIL
E. Panama (Darien) and Colombia.	
Margarornis bellulus	BEAUTIFUL TREERUNNER
Mts. e. Panama (Darien).	
Margarornis rubiginosus	RUDDY TREERUNNER
Mts. Costa Rica and w. Panama.	
Premnoplex brunneascens	SPOTTED BARBTAIL
Highlands Costa Rica, Panama and w. S. Am.	

¹ Birds from s.e. Nicaragua to w. Ecuador, the *aequatorialis* group, found chiefly in the lowlands, are sometimes deemed a separate species, Spot-throated Woodcreeper. Hellmayr treats all as races of the S. Am. *triangularis*.

² Peters includes in this species the *lacrymiger* group of S. Am.

³ "Sicklebill", sometimes used for this genus, is the name of a genus of hummingbirds *Eutoxeres*, *supra*.

⁴ It is uncertain whether this little-known genus should be included in Furnariidae.

Pseudocolaptes lawrencii⁵	BUFFY TUFTEDCHEEK
Highlands Costa Rica, w. Panama and n.w. S. Am.	
Hyloctistes subulatus	STRIPED WOODHAUNTER
Nicaragua, Costa Rica, Panama and w. S. Am.	
Syndactyla⁶ subalaris	LINEATED FOLIAGE-GLEANER
Highlands Costa Rica, Panama and w. S. Am.	
Anabacerthia variegaticeps⁷	SCALY-THROATED FOLIAGE-GLEANER
Highlands trop. Mexico, Guatemala, Honduras, Costa Rica and w. Panama.	
Philydor erythrocercus⁸	RUFOUS-RUMPED FOLIAGE-GLEANER
Panama and S. Am.	
Philydor rufus	BUFF-FRONTED FOLIAGE-GLEANER
Highlands Costa Rica, Panama and S. Am.	
Automolus rubiginosus⁹	RUDDY FOLIAGE-GLEANER
Highlands Trop. Mid. Am. (except Brit. Honduras) and S. Am.	
Automolus ochrolaemus¹⁰	BUFF-THROATED FOLIAGE-GLEANER
Trop. Mid. Am. (except El Salvador), and S. Am.	
Thripadectes rufobrunneus	STREAK-BREASTED TREEHUNTER
Highlands Costa Rica and w. Panama.	
Xenops rutilans	STREAKED XENOPS
Highlands Costa Rica, Panama and S. Am.	
Xenops minutus	PLAIN XENOPS
Trop. Mid. Am. (except El Salvador), and S. Am.	
Sclerurus albicularis	GRAY-THROATED LEAFSCRAPER
Costa Rica and w. Panama (Chiriquí), also S. Am.	
Sclerurus mexicanus	TAWNY-THROATED LEAFSCRAPER
S. Mexico, Guatemala, Honduras, Costa Rica, Panama, and S. Am.	
Sclerurus guatemalensis	SCALY-THROATED LEAFSCRAPER
Trop. Mid. Am. (except El Salvador), Colombia and Ecuador.	

⁵ Hellmayr treats all *Pseudocolaptes* as conspecific with the white-tufted *P. boissonneautii* of the S. Am. Andes (chiefly Temperate Zone); if so, Tuftedcheek could be used for all. Zimmer and de Schauensee regard the *lawrencii* group as a distinct species.

⁶ *Xenoctistes* of Hellmayr and *Xenicopsis* of Ridgway.

⁷ Sometimes regarded as conspecific with *striaticollis* of w. S. Am. Hellmayr uses the name *Xenicopoides montanus*, including also birds of e. S. Am.

⁸ Birds from Panama and n.w. S. Am. are sometimes treated as a separate species, *P. fuscipennis*, Dusky-winged Foliage-gleaner.

⁹ Birds from e. Panama and n.w. S. Am. are sometimes regarded as a distinct species *A. nigricauda*, Black-tailed Foliage-gleaner.

¹⁰ The birds of Mid. Am. to w. Ecuador belong to the *pallidigularis* group, which Todd regards as specifically distinct from *ochrolaemus* of S. Am. east of the Andes (Ann. Carn. Mus. 31(4): 41, 1948).

Lochmias nematura SHARP-TAILED STREAMCREEPER
Highlands e. Panama (Darien), and S. Am.

Family **FORMICARIIDAE** ANTBIRDS

<i>Cymbilaimus lineatus</i>	FASCIATED ANTSHRIKE
Nicaragua, Costa Rica, Panama and S. Am.	
<i>Taraba major</i>	GREAT ANTSHRIKE
Trop. Mid. Am. (except Brit. Honduras and El Salvador), and S. Am.	
<i>Thamnophilus doliatus</i>	BARRED ANTSHRIKE
Trop. Mid. Am. and S. Am.	
<i>Thamnophilus bridgesi</i>	BLACK-HOODED ANTSHRIKE
S.w. Costa Rica and s.w. Panama.	
<i>Thamnophilus nigriceps</i>	BLACK ANTSHRIKE
E. Panama (Darien), and Colombia.	
<i>Thamnophilus punctatus</i>	SLATY ANTSHRIKE
Brit. Honduras, Guatemala, Honduras, Nicaragua, Costa Rica, Panama and S. Am.	
<i>Xenornis setifrons</i>	GRAY-FACED ANTBIRD
E. Panama and n.w. Colombia.	
<i>Thamnistes anabatinus</i>	RUSSET ANTSHRIKE
Trop. Mid. Am. (except El Salvador), and w. S. Am.	
<i>Dysithamnus mentalis</i>	PLAIN ANTVIREO
S. Mexico (Campeche), Guatemala, Honduras, Costa Rica, Panama and S. Am.	
<i>Dysithamnus striaticeps</i>	STREAK-CROWNED ANTVIREO
Caribbean slope Nicaragua, Costa Rica and n.w. Panama (s of R. Sixaola, <i>fide</i> M. A. Carriker).	
<i>Dysithamnus puncticeps</i>	SPOT-CROWNED ANTVIREO
E. Costa Rica, Panama, Colombia and Ecuador.	
<i>Myrmotherula brachyura</i>	PYGMY ANTWREN
Cent. Panama to S. Am.	
<i>Myrmotherula surinamensis</i>	STREAKED ANTWREN
Panama and S. Am.	
<i>Myrmotherula fulviventris</i>	FULVOUS-BELLIED ANTWREN
S. Honduras, Nicaragua, Costa Rica, Panama, Colombia and Ecuador.	
<i>Myrmotherula axillaris</i>	WHITE-FLANKED ANTWREN
S. Honduras, Nicaragua, Costa Rica, Panama and S. Am.	
<i>Myrmotherula schisticolor</i>	SLATY ANTWREN
Trop. Mid. Am. (from Chiapas south, except Brit. Honduras and El Salvador), and n.w. S. Am.	

Herpsilochmus rufimarginatus	RUFOUS-WINGED ANTWREN
E. Panama (Darien), and S. Am.	
Microrhopias quixensis¹	DOT-WINGED ANTWREN
Trop. Mid. Am. (except El Salvador) and S. Am.	
Formicivora grisea	BLACK-BREASTED ANTWREN
Panama (Pearl Is.), and n. and e. S. Am.	
Terenura callinota	RUFOUS-RUMPED ANTWREN
Highlands w. Panama, n. and n.w. S. Am.	
Cercomacra tyrannina	DUSKY ANTBIRD
Trop. Mid. Am. (except El Salvador), and S. Am.	
Cercomacra nigricans	JET ANTBIRD
Cent. Panama to n. S. Am.	
Gymnichla nudiceps	BARE-CROWNED ANTBIRD
Guatemala, Honduras, Nicaragua, Costa Rica, Panama and n. Colombia.	
Myrmeciza longipes	WHITE-BELLIED ANTBIRD
Cent. Panama to n. S. Am.	
Myrmeciza exsul²	CHESTNUT-BACKED ANTBIRD
Nicaragua, Costa Rica, Panama, Colombia and Ecuador.	
Myrmeciza laemosticta	DULL-MANTLED ANTBIRD
Costa Rica, Panama and n.w. S. Am.	
Myrmeciza immaculata	IMMACULATE ANTBIRD
Costa Rica, Panama and n.w. S. Am.	
Formicarius analis	BLACK-FACED ANTTHRUSH
Trop. Mid. Am. (except El Salvador), and S. Am.	
Formicarius nigricapillus	BLACK-HEADED ANTTHRUSH
Costa Rica, Panama, Colombia and Ecuador.	
Formicarius rufipectus	RUFOUS-BREASTED ANTTHRUSH
Highlands Costa Rica, Panama and n.w. S. Am.	
Gymnopithys leucaspis³	BICOLORED ANTBIRD
Honduras, Nicaragua, Costa Rica, Panama and n.w. S. Am.	
Hylophylax naevioides	SPOTTED ANTBIRD
E. Nicaragua, Costa Rica, Panama, Colombia and Ecuador.	
Phaenostictus mcleannanni	OCELLATED ANTTHRUSH
S.e. Nicaragua, Costa Rica, Panama, Colombia and Ecuador.	

¹ Middle American birds belong to the *boucardi* group, which some authors regard as distinct from S. Am. *quixensis*.

² Birds from e. Panama and S. Am. *M. maculifer*, are by some deemed a separate species. If so, they may be called Wing-spotted Antbird.

³ Middle American birds belong to the *bicolor* group, sometimes regarded as a species distinct from *leucaspis*.

Myrmornis torquata⁴	WING-BANDED ANTHRUSH
S.e. Nicaragua, e. Panama and S. Am.	
Pittasoma michleri	BLACK-CROWNED ANTPITTA
Costa Rica, Panama and Colombia.	
Grallaricula flavirostris	OCHRE-BREASTED ANTPITTA
Mts. Costa Rica, Panama and w. S. Am.	
Grallaria guatimalensis	SCALED ANTPITTA
Highlands trop. Mid. Am. (except Brit. Honduras) and n. S. Am.	
Grallaria fulviventris	FULVOUS-BELLIED ANTPITTA
E. Nicaragua, Costa Rica, Panama, Colombia and Ecuador.	
Grallaria perspicillata	STREAK-CHESTED ANTPITTA
E. Nicaragua, Costa Rica, Panama, Colombia and Ecuador.	

Family RHINOCRYPTIDAE TAPACULOS

Scytalopus argentifrons⁵	SILVERY-FRONTED TAPACULO
Mts. Costa Rica and w. Panama.	
Scytalopus panamensis	PALE-THROATED TAPACULO
Highlands e. Panama (Darien), Colombia and Ecuador.	

Family PIPRIDAE MANAKINS

Piprites griseiceps	GRAY-HEADED MANAKIN
Nicaragua and Costa Rica (chiefly Caribbean slope).	
Pipra coronata	BLUE-CROWNED MANAKIN
Costa Rica, Panama and S. Am.	
Pipra mentalis	RED-CAPPED MANAKIN
Trop. Mid. Am. (except El Salvador), and S. Am.	
Pipra erythrocephala	FLAME-HEADED MANAKIN
E. Panama and S. Am.	
Pipra pipra	WHITE-CROWNED MANAKIN
Highlands Costa Rica, w. Panama and S. Am.	
Chloropipo holochlora	GREEN MANAKIN
E. Panama and n.w. S. Am.	
Chiroxiphia linearis	LONG-TAILED MANAKIN
Trop. Mid. Am. (chiefly Pacific slope, not recorded Brit. Honduras) to Costa Rica.	
Chiroxiphia lanceolata	LANCE-TAILED MANAKIN
Panama (chiefly Pacific slope), n. Colombia and Venezuela.	

⁴ The Middle American form is *stictoptera*, often considered a separate species. Ridgway used the generic name *Rhopoterpe*.

⁵ Includes *chiriquensis* of Panama (e. Chiriqui and Veraguas).

Corapipo leucorrhoea⁶	WHITE-RUFFED MANAKIN
Nicaragua, Costa Rica, Panama, Colombia and Venezuela.	
Manacus aurantiacus⁷	ORANGE-COLLARED MANAKIN
Pacific slope of Costa Rica and w. Panama.	
Manacus cerritus⁷	ALMIRANTE MANAKIN
Caribbean slope of extreme n.w. Panama (Almirante Bay).	
Manacus vitellinus	GOLDEN-COLLARED MANAKIN
Panama and Colombia.	
Manacus candei	WHITE-COLLARED MANAKIN
Trop. Mid. Am. (chiefly Caribbean slope) to n.e. Costa Rica.	
Schiffornis⁸ turdinus	THRUSH-LIKE MANAKIN
Trop. Mid. Am. (except El Salvador), and S. Am.	
Sapayoaaenigma	BROAD-BILLED MANAKIN
E. Panama, w. Colombia and n.w. Ecuador.	

Family COTINGIDAE COTINGAS

Cotinga amabilis⁹	LOVELY COTINGA
Caribbean slope trop. Mid. Am. s. Mexico to Costa Rica.	
Cotinga ridgwayi	TURQUOISE COTINGA
Pacific slope s.w. Costa Rica and s.w. Panama.	
Cotinga nattererii	BLUE COTINGA
Cent. Panama to Colombia and Ecuador.	
Carpodectes nitidus	SNOWY COTINGA
Caribbean slope Honduras, Nicaragua, Costa Rica and w. Panama.	
Carpodectes antoniae¹⁰	YELLOW-BILLED COTINGA
Pacific slope s.w. Costa Rica and s.w. Panama (Chiriqui).	
Carpodectes hopkei¹⁰	BLACK-TIPPED COTINGA
E. Panama (?) (Darien, presumably this seen, T. Barbour), w. Colombia and Ecuador.	
Attila spadiceus	BRIGHT-RUMPED ATTILA
Trop. Mid. Am. and S. Am.	
Laniocera rufescens¹¹	SPECKLED MOURNER
Guatemala, Honduras, Nicaragua, Costa Rica, Panama and n.w. S. Am.	

⁶ Mid. Am. birds belong to the *altera* group, ranging to n.w. Colombia, perhaps specifically distinct from *leucorrhoea*. "White-throated" Manakin, sometimes used for this species, is the name generally applied to another species, *C. gutturalis* of S. Am.

⁷ If regarded as races of *M. vitellinus*, Golden-collared Manakin is appropriate for the entire complex.

⁸ *Scotothorus* of Ridgway.

⁹ This and the next two forms may be conspecific, perhaps races of the Upper Amazonian *C. maynana*; if so, Blue Cotinga can be used.

¹⁰ Regarded by Hellmayr and Zimmer as races of *C. nitidus*, the Snowy Cotinga.

¹¹ Placed by Ridgway in the family Pipridae.

Rhytipterna holerythra ²	RUFOUS MOURNER
Trop. Mid. Am. (except Brit. Honduras and El Salvador), and n.w. S. Am.	
Lipaugs unirufus ³	RUFOUS PIHA
Trop. Mid. Am. (except El Salvador), and n.w. S. Am.	
Pachyramphus versicolor	BARRED BECARD
Highlands Costa Rica, w. Panama and n.w. S. Am.	
Pachyramphus rufus	CINEREOUS BECARD
E. Panama and S. Am.	
Pachyramphus cinnamomeus	CINNAMON BECARD
Trop. Mid. Am. (except El Salvador), and n. S. Am.	
Pachyramphus polychopterus	WHITE-WINGED BECARD
Guatemala, Honduras, Nicaragua, Costa Rica, Panama and S. Am.	
Pachyramphus albogriseus	BLACK-AND-WHITE BECARD
W. Nicaragua, Costa Rica, Panama and n. S. Am.	
Pachyramphus major	GRAY-COLLARED BECARD
Trop. Mid. Am. from Mexico to e. Nicaragua.	
Platyparis homochrous	ONE-COLORED BECARD
Cent. Panama to w. S. Am.	
Platyparis aglaiae	ROSE-THROATED BECARD
S.w. U. S. and Mid. Am. to n. Costa Rica.	
Tityra semifasciata	MASKED TITYRA
Trop. Mid. Am. and S. Am.	
Tityra ⁴ <i>inquisitor</i>	BLACK-CROWNED TITYRA
Trop. Mid. Am. (except El Salvador), and S. Am.	
Querula purpurata	PURPLE-THROATED FRUITCROW
Costa Rica, Panama and S. Am.	
Cephalopterus glabricollis ⁵	BARE-NECKED UMBRELLABIRD
Costa Rica and w. Panama (chiefly highlands).	
Procnias tricarunculata	THREE-WATTLED BELLBIRD
Nicaragua, Costa Rica and w. Panama (chiefly highlands).	

Family TYRANNIDAE TYRANT FLYCATCHERS

Sayornis phoebe	EASTERN PHOEBE
E. N. Am.; winters to Mexico.	
Sayornis nigricans	BLACK PHOEBE
W. N. Am. and highlands Mid. Am. (except Brit. Honduras) to S. Am.; northern birds winter occasionally to Costa Rica.	

² Ridgway placed this species in the genus *Lipaugs*.

³ Ridgway placed this species in the genus *Lathria*.

⁴ Sometimes placed in a separate genus *Erator*.

⁵ By Hellmayr all Umbrellabirds are deemed races of the S. Am. *C. ornatus*.

<i>Sayornis saya</i>	SAY'S PHOEBE
W. N. Am. and n.w. Mexico; winters to Mexico.	
<i>Colonia colonus</i>	LONG-TAILED TYRANT
S. Honduras, Nicaragua, Costa Rica, Panama and S. Am.	
<i>Fluvicola pica</i>	PIED WATER-TYRANT
E. Panama and S. Am.	
<i>Procephalus rubinus</i>	VERMILION FLYCATCHER
S.w. U. S., Mexico, Brit. Honduras, Guatemala, Honduras, also S. Am.; recorded Panama (once, photographed, E. Eisenmann).	
<i>Muscivora forficata</i>	SCISSOR-TAILED FLYCATCHER
W. U. S.; winters through Mid. Am. (not reported Brit. Honduras).	
<i>Muscivora tyrannus</i>	FORK-TAILED FLYCATCHER
Mid. Am. and S. Am.	
<i>Tyrannus tyrannus</i>	EASTERN KINGBIRD
E. N. Am.; migrates through Mid. Am.; winters from Honduras to S. Am.	
<i>Tyrannus vociferans</i>	CASSIN'S KINGBIRD
W. N. Am. and n. Mexico; winters to Guatemala.	
<i>Tyrannus verticalis</i>	WESTERN KINGBIRD
W. N. Am. to n.w. Mexico; winters to Mexico, Guatemala, El Salvador, Honduras, Nicaragua.	
<i>Tyrannus melancholicus</i>	TROPICAL KINGBIRD
S.w. U. S., Mid. Am. and S. Am.	
<i>Tyrannus dominicensis</i>	GRAY KINGBIRD
S.e. U. S., Bahamas, West Indies; winters to Panama and n. S. Am.; a few records Mexico (Cozumel I. and Cancun I. off Yucatan Pen.), Honduras, Nicaragua, Costa Rica.	
<i>Tyrannus cubensis</i>	GIANT KINGBIRD
West Indies; recorded once from Mexico (Mujeres I. off Yucatan Pen.).	
<i>Tyrannus crassirostris</i>	THICK-BILLED KINGBIRD
W. Mexico to w. Guatemala.	
<i>Legatus leucophaius</i>	PIRATIC FLYCATCHER
Trop. Mid. Am. (not reported El Salvador, Honduras), and S. Am.	
<i>Sirystes⁶ sibilator</i>	SIRYSTES
Cent. Panama to S. Am.	
<i>Myiodynastes luteiventris</i>	SULPHUR-BELLIED FLYCATCHER
S.w. U. S., Mid. Am. to Costa Rica; winters s. to Panama and n.w. S. Am.	
<i>Myiodynastes maculatus</i>	STREAKED FLYCATCHER
Trop. Mid. Am. (not recorded El Salvador and Nicaragua), and S. Am.	

⁶ This monotypic genus may belong in Cotingidae, as Ridgway believed.

Myiodynastes hemichrysus ⁷	GOLDEN-BELLIED FLYCATCHER
Highlands of Costa Rica and w. Panama.	
Megarynchus pitangua	BOAT-BILLED FLYCATCHER
Trop. Mid. Am. and S. Am.	
Coryphotriccus albovittatus ⁸	WHITE-RINGED FLYCATCHER
Costa Rica, Panama, w. Colombia and Ecuador.	
Myiozetetes cayanensis	RUSTY-MARGINED FLYCATCHER
Cent. Panama to S. Am.	
Myiozetetes similis	SOCIAL ⁹ FLYCATCHER
Trop. Mid. Am. and S. Am.	
Myiozetetes granadensis	GRAY-CAPPED FLYCATCHER
Nicaragua, Costa Rica, Panama and S. Am.	
Pitangus sulphuratus	GREAT KISKADEE
S. Texas, Mid. Am. to n.w. Panama (Bocas del Toro), also S. Am.	
Pitangus lictor	LESSER KISKADEE
Cent. Panama to S. Am.	
Myiarchus crinitus	GREAT-CRESTED FLYCATCHER
E. N. Am.; winters through Mid. Am. to Colombia.	
Myiarchus cinerascens	ASH-THROATED FLYCATCHER
W. U. S. and n. Mexico; winters to Guatemala and El Salvador, casually to n.w. Costa Rica.	
Myiarchus nuttingi ¹⁰	PALE-THROATED FLYCATCHER
Trop. Mid. Am. (except Brit. Honduras) to n.w. Costa Rica.	
Myiarchus tyrannulus ¹	BROWN-CRESTED FLYCATCHER
S.w. U. S., Mid. Am. to n.w. Costa Rica, also n. and e. S. Am. and Lesser Antilles.	
Myiarchus yucatanensis	YUCATAN FLYCATCHER
Mexico (n. Yucatan Pen. and Cozumel I.).	
Myiarchus ferox	SHORT-CRESTED FLYCATCHER
S.w. Costa Rica, Panama and S. Am.	
Myiarchus tuberculifer	DUSKY-CAPPED FLYCATCHER
Arizona, Mid. Am. and S. Am.	

⁷ By Hellmayr treated as a race of the S. Am. *M. chrysocephalus*, Golden-crowned Flycatcher.

⁸ By Hellmayr treated as a race of *C. parvus*, of n.e. S. Am.

⁹ "Vermilion-crowned" is sometimes used.

¹⁰ Sometimes regarded as conspecific with *M. cinerascens*.

¹ The A.O.U. Check-List Committee and van Rossem are here followed in the rather questionable lumping with the South American *M. tyrannulus* of the northern *M. magister* group from Arizona to Honduras and El Salvador (at least in winter), and the Central American *M. brachyurus*, from the Pacific slope of Chiapas to n.w. Costa Rica. If more than one species be recognized, Brown-crested Flycatcher applies to the *magister* group. *M. brachyurus* has been called Ometepe Flycatcher.

Nesotriccus ridgwayi	Cocos ISLAND FLYCATCHER
Costa Rica (Cocos I.).	
Deltarhynchus flammulatus	FLAMMULATED FLYCATCHER
S.w. and s. Mexico (Jalisco to Chiapas).	
Nuttallornis borealis	OLIVE-SIDED FLYCATCHER
N. Am. and mts. n.w. Mexico (Baja Calif.); migrates through Mid. Am. (not reported Brit. Honduras); winters chiefly S. Am., occasionally n. to Costa Rica.	
Contopus virens	EASTERN WOOD-PEWEE
E. N. Am.; migrates through Mid. Am. (not reported El Salvador); winters from Nicaragua but chiefly to n.w. S. Am.	
Contopus richardsonii²	WESTERN WOOD-PEWEE
W. N. Am. and highlands Mid. Am. (breeding s. of Guatemala uncertain); migrates through Mid. Am.; winters through Mid. Am. but chiefly in w. S. Am.	
Contopus cinereus³	TROPICAL PEWEE
Trop. Mid. Am. and S. Am.	
Contopus pertinax⁴	GREATER PEWEE
Highlands Arizona, and Mid. Am. s. to n. Nicaragua.	
Contopus lugubris⁴	DARK PEWEE
Highlands Costa Rica and w. Panama.	
Contopus ochraceus	OCHRACEOUS PEWEE
Mts. Costa Rica and w. Panama (Chiriqui, one spec.).	
Epidonax flaviventris	YELLOW-BELLIED FLYCATCHER
N.e. N. Am.; winters through Mid. Am. (not reported Brit. Honduras).	
Epidonax virescens	ACADIAN FLYCATCHER
E. N. Am.; migrates through Mid. Am. (not reported Brit. Honduras, El Salvador); winters in Panama and n. S. Am.	
Epidonax traillii	TRAILL'S FLYCATCHER
N. Am.; winters Mid. Am. (unreported Brit. Honduras) to S. Am.	
Epidonax minimus	LEAST FLYCATCHER
N.e. N. Am.; winters through Mid. Am. to cent. Panama.	
Epidonax hammondii	HAMMOND'S FLYCATCHER
W. N. Am.; winters to Mexico, Guatemala, El Salvador, Honduras, Nicaragua.	

² Perhaps conspecific with *C. virens*. If so, call the complex Wood Pewee. The genus *Contopus* was formerly known as *Myiochanes*.

³ The Middle American birds belong to the *brachytarsus* group.

⁴ Zimmer treats both as races of the South American *C. fumigatus*. If so, the entire complex may be called Greater Pewee.

Empidonax oberholseri⁵	WRIGHT'S FLYCATCHER
W. N. Am.; winters to Mexico.	
Empidonax wrightii⁶	GRAY FLYCATCHER
W. U. S. to (?) n.w. Mexico; winters to Mexico.	
Empidonax affinis⁶	PINE FLYCATCHER
Oak-pine forests of Mexico; winters to Guatemala.	
Empidonax difficilis	WESTERN FLYCATCHER
W. N. Am., mts. Mexico, Guatemala and Honduras.	
Empidonax flavescens	YELLOWISH FLYCATCHER
Highlands of Mid. Am. (except Brit. Honduras) from s. Mexico (Chiapas) to w. Panama.	
Empidonax albicularis	WHITE-THROATED FLYCATCHER
Highlands (chiefly) of Mid. Am. (except Brit. Honduras and El Salvador) to w. Panama.	
Empidonax atriceps	BLACK-CAPPED FLYCATCHER
Mts. Costa Rica and w. Panama.	
Empidonax fulvifrons	BUFF-BREASTED FLYCATCHER
S.w. U. S. and highlands of n. and w. Mexico, Guatemala, El Salvador and Honduras.	
Aechmophorus mexicanus	PILEATED FLYCATCHER
Mexico (highlands of Michoacan to Oaxaca).	
Xenotriccus callizonus	BELTED FLYCATCHER
S. Mexico (Chiapas) and Guatemala.	
Mitrephanes phaeocercus	TUFTED FLYCATCHER
Highlands trop. Mid. Am. (except Brit. Honduras) and w. S. Am.	
Terenotriccus erythrurus	RUDDY-TAILED FLYCATCHER
Guatemala, Honduras, Nicaragua, Costa Rica, Panama and S. Am.	
Aphanotriccus capitalis	TAWNY-CHESTED FLYCATCHER
E. Nicaragua and e. Costa Rica.	
Aphanotriccus⁷ audax	BLACK-BILLED FLYCATCHER
E. Panama and n.w. Colombia.	
Myiobius sulphureipygius⁸	SULPHUR-RUMPED FLYCATCHER
Trop. Mid. Am. (except El Salvador), and n.w. S. Am.	

⁵ The A.O.U. Committee is reluctantly followed in transferring to the bird formerly called *E. griseus* the specific name *wrightii*, and in renaming Wright's Flycatcher *E. oberholseri* (Auk, 70: 360, 1953; Phillips, Auk, 56: 311-312, 1939). Moreover A. Phillips has found that no published records from Guatemala of Wright's Flycatcher are actually that species (Auk, 61: 294, 1944; Condor, 49: 121, 1947).

⁶ This includes *E. "fulvipectus"*, *E. trepidus* and *E. pulverius* (see Moore, Auk, 57: 349-350, 364-365, 1940).

⁷ *Praedo* of some authors.

⁸ Treated as conspecific with *barbatus* of e. S. Am. by some authors.

Myiobius atricaudus	BLACK-TAILED FLYCATCHER
Costa Rica, Panama and S. Am.	
Myiophobus fasciatus	BRAN-COLORED FLYCATCHER
Costa Rica, Panama and S. Am.	
Onychorhynchus mexicanus⁹	NORTHERN ROYAL-FLYCATCHER
Trop. Mid. Am., n. Colombia and n.w. Venezuela.	
Platyrinchus mystaceus¹⁰	WHITE-THROATED SPADEBILL
Trop. Mid. Am. (not reported Brit. Honduras) and S. Am.	
Platyrinchus coronatus	GOLDEN-CROWNED SPADEBILL
Honduras, Nicaragua, Costa Rica, Panama and S. Am.	
Cnemotriccus subbrunneus	BROWNISH FLYCATCHER
E. Panama and S. Am.	
Tolmomyias sulphurescens¹	YELLOW-OLIVE FLYCATCHER
Trop. Mid. Am. (not reported Brit. Honduras) and S. Am.	
Tolmomyias assimilis²	YELLOW-MARGINED FLYCATCHER
Costa Rica, Panama and S. Am.	
Rhynchocyclus olivaceus	OLIVACEOUS FLATBILL
Panama and S. Am.	
Rhynchocyclus brevirostris	EYE-RINGED FLATBILL
Trop. Mid. Am. (except Brit. Honduras) and n.w. S. Am.	
Todirostrum nigriceps³	BLACK-HEADED TODY-FLYCATCHER
Costa Rica, Panama and n.w. S. Am.	
Todirostrum cinereum	COMMON TODY-FLYCATCHER
Trop. Mid. Am. and S. Am.	
Todirostrum sylvia	SLATE-HEADED TODY-FLYCATCHER
Trop. Mid. Am. (except El Salvador), and n. S. Am.	
Oncostoma cinereigulare	NORTHERN BENTBILL
Trop. Mid. Am. to cent. Panama.	
Oncostoma olivaceum⁴	SOUTHERN BENTBILL
Cent. Panama to Colombia.	

⁹ All Royal Flycatchers may be races of *coronatus* of e. S. Am.

¹⁰ Birds from Mexico to w. Costa Rica, *P. cancrinomimus*, are by some regarded as a species, Stub-tailed Spadebill.

¹ Includes *flavolivaceus* and *cinereiceps*, (placed in *Rhynchocyclus* by Ridgway, who called the true members of that genus, *Craspedopriion*). Unlike true *Rhynchocyclus*, whose breeding behavior is like the Pipridae, *Tolmomyias* behaves normally (Skutch, *Ibis* 1953: 4, 33-34). As these genera may not be closely allied, it seems best to reserve the special group name "Flatbill" to the aberrant *Rhynchocyclus*. Ridgway's name "Yellow-olive Flycatcher" is favored because "Sulphury Flycatcher" has been used for another species.

² Called *T. flavotectus* by Hellmayr and *Rhynchocyclus marginatus* by Ridgway.

³ Considered conspecific with *chrysocrotaphum* of e. S. Am. by Zimmer.

⁴ Both Bentbills are considered races of *cinereigulare* by Hellmayr.

Lophotriccus pileatus	SCALE-CRESTED PYGMY-TYRANT
Highlands Costa Rica, Panama and n.w. S. Am.	
Atalotriccus pilaris	WHITE-EYED PYGMY-TYRANT
Panama and n. S. Am.	
Perissotriccus atricapillus ⁵	BLACK-CAPPED PYGMY-TYRANT
Costa Rica, Panama, n.w. Colombia and Ecuador.	
Pseudotriccus pelzelni	STREAK-CROWNED PYGMY-TYRANT
Mts. e. Panama (Darien) and n.w. S. Am.	
Phylloscartes flavovirens ⁶	YELLOW-GREEN TYRANNULET
Panama (Canal Zone eastward).	
Phylloscartes ⁷ <i>superciliaris</i>	RUFOUS-BROWED TYRANNULET
Mts. Costa Rica, Panama and n. S. Am.	
Capsiempis flaveola	YELLOW TYRANNULET
Nicaragua, Costa Rica, Panama and S. Am.	
Serpophaga cinerea	TORRENT TYRANNULET
Rapid streams (chiefly highlands) Costa Rica, w. Panama and w. S. Am.	
Elaenia flavogaster	YELLOW-BELLIED ELAENIA
Trop. Mid. Am., S. Am., s. Lesser Antilles.	
Elaenia martinica ⁸	CARIBBEAN ELAENIA
Lesser Antilles, other Caribbean islands, Mexico (is. off Yucatan Pen.), (?) Brit. Honduras (Half Moon Cay).	
Elaenia chiriquensis	LESSER ELAENIA
S.w. Costa Rica, Panama and S. Am.	
Elaenia frantzii ⁹	MOUNTAIN ELAENIA
Mts. Guatemala, El Salvador, Honduras, Nicaragua, Costa Rica, w. Panama, n. S. Am.	
Myiopagis ¹⁰ <i>gaimardii</i>	FOREST ELAENIA
Cent. Panama through S. Am.	
Myiopagis viridicata	GREENISH ELAENIA
Trop. Mid. Am. and S. Am.	

⁵ Zimmer and others merge *Perissotriccus* in *Myiornis* and treat this form as a race of the gray-headed *ecaudatus* of n. and e. S. Am., Short-tailed Pygmy-Tyrant.

⁶ Ridgway placed this species in the genus *Leptopogon*; Hellmayr treated it as a race of *P. ventralis* of e. S. Am. (But cf. Zimmer, Am. Mus. Nov. 1095: 2-3, 1940).

⁷ Placed in the genus *Mecocerculus* by Hellmayr.

⁸ Includes *chinchorrensis* of Chinchorro Bank off the Yucatan Peninsula.

⁹ Considered by Hellmayr (but not Zimmer) conspecific with *obscura* of cent. S. Am., the Dusky Elaenia.

¹⁰ *Myiopagis* is merged in *Elaenia* by Hellmayr, but considered distinct by Zimmer. Ridgway considered *gaimardii* a member of Cotingidae and erected for it the genus *Elainopsis*.

Sublegatus arenarum ¹	SCRUB FLYCATCHER
Costa Rica, Panama and S. Am.	
Phaeomyias murina	MOUSE-COLORED TYRANNULET
Semi-arid areas, Panama and S. Am.	
Campstostoma imberbe	NORTHERN BEARDLESS TYRANNULET ²
S.w. U. S. and Mid. Am. to n.w. Costa Rica.	
Campstostoma obsoletum	SOUTHERN BEARDLESS TYRANNULET
Costa Rica, Panama and S. Am.	
Phyllomyias griseiceps	SOOTY-HEADED TYRANNULET
N. S. Am. to e. Panama (Darien).	
Tyranniscus vilissimus	PALTRY TYRANNULET
Trop. Mid. Am. (from Chiapas south, except Brit. Honduras) and S. Am.	
Tyrannulus ³ elatus	YELLOW-CROWNED TYRANNULET
Panama and S. Am.	
Acrochordopus ³ zeledoni	WHITE-FRONTED TYRANNULET
Highlands Costa Rica, w. Panama and n. S. Am.	
Ornithion ³ semiflavum	YELLOW-BELLIED TYRANNULET
Trop. Mexico, Guatemala, Honduras, Nicaragua and Pac. slope of Costa Rica.	
Ornithion brunneicapillum ⁴	BROWN-CAPPED TYRANNULET
Caribbean slope of Costa Rica and Panama (Pac. slope also in e. half), and n. S. Am.	
Leptopogon superciliaris	SLATY-CAPPED FLYCATCHER
Highlands of Costa Rica, Panama and n. S. Am.	
Leptopogon amaurocephalus	SEPIA-CAPPED FLYCATCHER
Trop. Mid. Am. (except Brit. Honduras and El Salvador), and S. Am.	
Mionectes olivaceus	OLIVE-STRIPED FLYCATCHER
Costa Rica, Panama and S. Am.	
Pipromorpha oleaginea	OCHRE-BELLIED FLYCATCHER
Trop. Mid. Am. and S. Am.	

Family OXYRUNCIDAE SHARPBILLS

Oxyruncus cristatus	SHARPBILL
Mts. Costa Rica, w. and e. Panama, and locally in S. Am.	

¹ Considered by Hellmayr (but not Zimmer) conspecific with *modestus* of s. S. Am.

² Hellmayr's group name "Tyrannulet" is preferable to "Flycatcher" because of the behavior and small size of the genus, though the A.O.U. Check-List Committee uses "Flycatcher".

³ Ridgway placed the species of these genera in the family Cotingidae.

⁴ By some deemed conspecific with the gray-capped *semiflavum*; if so, Yellow-bellied Tyrannulet is the best name for the species. *Microtriccus* is used as the generic name by Ridgway, Hellmayr and others; Zimmer is here followed.

Family ALAUDIDAE LARKS

Eremophila alpestris HORNERD LARK
Eurasia, N. Am. to Mexico (s. to Oaxaca); Colombia.

Family HIRUNDINIDAE SWALLOWS

Progne subis⁵	PURPLE MARTIN
N. Am., Mexico, West Indies; the e. N. Am. race winters chiefly in Brazil, with a few records from e. Mexico, Guatemala, Brit. Honduras, Honduras, Nicaragua.	
Progne chalybea	GRAY-BREASTED MARTIN
Texas, Mid. Am. and S. Am.	
Phaeoprogne⁶ tapera	BROWN-CHESTED MARTIN
S. Am.; the southern race <i>fusca</i> regularly migrates n. to Panama during the Southern Hemisphere winter.	
Petrochelidon pyrrhonota	CLIFF SWALLOW
N. Am. and Mexico; migrates through Mid. Am. (not reported Brit. Honduras and Nicaragua); winters in S. Am.	
Petrochelidon fulva	CAVE SWALLOW
West Indies, Texas, e. Mexico (s. to Chiapas and Yucatan).	
Hirundo rustica	BARN SWALLOW
Old World, N. Am. to cent. Mexico; migrates through Mid. Am.; winters in S. Am., occasionally in Mid. Am.	
Stelgidopteryx ruficollis	ROUGH-WINGED SWALLOW
N. Am., Mid. Am. and S. Am.; the northern races winter through Mid. Am.	
Neochelidon tibialis	WHITE-THIGHED SWALLOW
Cent. Panama through S. Am.	
Pygochelidon cyanoleuca	BLUE-AND-WHITE SWALLOW
Highlands Costa Rica, w. Panama, S. Am. The s. S. Am. form <i>patagonica⁷</i> sometimes winters n. to cent. Panama, recorded from Nicaragua (<i>fide</i> T. R. Howell).	
Notiochelidon pileata	BLACK-CAPPED SWALLOW
Highlands Mexico (Chiapas), Guatemala and El Salvador.	
Riparia riparia	BANK SWALLOW
Old World, N. Am.; migrates through Mid. Am. (not reported Brit. Honduras), winters in S. Am.	

⁵ Includes *sinaloae* of w. Mexico (Sinaloa, Jalisco), which occurs also in Guatemala. Some authors regard this as a race of the West Indian *P. dominicensis*, which they hold to be specifically distinct and which might be called Snowy-bellied Martin.

⁶ Some authors merge this genus in *Progne*.

⁷ This form, the Patagonian Swallow, may be a distinct species. De Schauensee merges the genus *Pygochelidon* in *Atticora*.

Iridoprocne bicolor	TREE SWALLOW
N. Am.; winters to Mexico, Brit. Honduras, Guatemala, Honduras, Nicaragua.	
Iridoprocne albilinea	MANGROVE SWALLOW
Trop. Mid. Am. and w. S. Am.	
Tachycineta thalassina	VIOLET-GREEN SWALLOW
W. N. Am. and Mexico (south to Oaxaca and Veracruz); winters to Guatemala, El Salvador, and Honduras; recorded Costa Rica.	

Family CORVIDAE CROWS AND JAYS

Corvus corax	COMMON RAVEN
Eurasia, N. Am., Mexico (western and plateau regions), Guatemala, El Salvador, Honduras, Nicaragua.	
Corvus cryptoleucus	WHITE-NECKED RAVEN
Deserts s.w. U. S. and Mexico (to Guanajuato).	
Corvus brachyrhynchos⁸	COMMON CROW
N. Am. to n.w. Mexico.	
Corvus imparatus⁹	MEXICAN CROW
N. Mexico (south to San Luis Potosi and Colima).	
Nucifraga columbiana	CLARK'S NUTCRACKER
W. N. Am. and mts. n.w. Mexico.	
Gymnorhinus cyanocephalus	PINYON JAY
Mts. w. U. S. to n.w. Mexico (Baja Calif., once Chihuahua).	
Calocitta formosa¹⁰	MAGPIE JAY
S.w. Mexico (Colima southward), Guatemala, El Salvador, Honduras, Nicaragua, and n.w. Costa Rica.	
Psilorhinus morio¹	PLAIN-TAILED BROWN JAY
E. Mexico (Nuevo Leon to Tabasco).	
Psilorhinus mexicanus¹	WHITE-TIPPED BROWN JAY
E. and s.e. Mexico, Guatemala, Honduras, Nicaragua, Costa Rica and n.w. Panama.	
Cyanocorax affinis	BLACK-CHESTED JAY
Costa Rica, Panama, Colombia and Venezuela.	
Cyanocorax dickeyi	TUFTED JAY
Mexico (mts. s.e. Sinaloa).	

⁸ Regarded by Hellmayr as conspecific with the Old World *C. corone*.

⁹ Hellmayr treats this as a race of the N. Am. Fish Crow, *C. ossifragus*.

¹⁰ Some authors divide the Magpie Jays into two species: *C. colliei*, Black-throated Magpie-Jay of n.w. Mexico, and *C. formosa*, White-throated Magpie-Jay of s.w. Mexico to Costa Rica.

¹ These are probably partly localized color phases. If so, the name is *P. morio*, Brown Jay.

Cyanocorax ² <i>yncas</i>	GREEN JAY
Texas, Mexico, Brit. Honduras, Guatemala, Honduras, also high-lands n. and w. S. Am.	
Cissilopha san-blasiana	BLACK-AND-BLUE JAY
S.w. Mexico (Nayarit to Guerrero).	
Cissilopha yucatanica ³	YUCATAN JAY
Mexico (Yucatan Pen. and Tabasco), Guatemala (Peten), and Brit. Honduras.	
Cissilopha beecheii	PURPLISH-BACKED JAY
N.w. Mexico (Sonora, Sinaloa and Nayarit).	
Cissilopha melanocyanea	BUSHY-CRESTED JAY
Highlands Guatemala, El Salvador, Honduras and Nicaragua.	
Cyanolyca cucullata ⁴	AZURE-HOODED JAY
Highlands S. Mexico, Guatemala, Honduras, Costa Rica and w. Panama.	
Cyanolyca mirabilis	WHITE-THROATED JAY
Mts. s.w. Mexico (Omilteme, Guerrero).	
Cyanolyca nana	DWARF JAY
Mts. s.e. Mexico (Veracruz, Mexico and Oaxaca).	
Cyanolyca pumilo	BLACK-THROATED JAY
Mts. s. Mexico (Chiapas), Brit. Honduras (<i>fide</i> Salvin and Godman), Guatemala, Honduras and El Salvador.	
Cyanolyca argenticula	SILVERY-THROATED JAY
Mts. Costa Rica and w. Panama.	
Aphelocoma coerulescens	SCRUB JAY
Florida, w. U. S. and plateau of Mexico.	
Aphelocoma ultramarina ⁵	GRAY-BREASTED JAY
Mts. s.w. U. S. and Mexico.	
Aphelocoma unicolor	UNICOLORED JAY
Mts. s. Mexico, Guatemala, El Salvador and Honduras.	
Cyanocitta stelleri	SELLER'S JAY
W. N. Am., highlands Mexico, Brit. Honduras, Guatemala, El Salvador, Honduras, and Nicaragua.	

Family PARIDAE TITMICE

Parus sclateri	MEXICAN OR GRAY-SIDED CHICKADEE
Mts. s.w. U. S. and Mexico (s. to Oaxaca).	

² Often placed in a separate genus *Xanthoura*.

³ Perhaps conspecific with *san-blasiana*; if so, use Black-and-blue Jay.

⁴ Hellmayr treats this as conspecific with the S. Am. *pulchra*; Pitelka disagrees.

⁵ Hellmayr uses the name *sordida* for this species.

Parus gambeli	MOUNTAIN CHICKADEE
Mts. w. N. Am. to n.w. Mexico (Baja Calif.).	
Parus atricristatus⁶	BLACK-CRESTED TITMOUSE
Texas and e. Mexico (to Veracruz).	
Parus inornatus	PLAIN TITMOUSE
W. U. S. and n.w. Mexico (Baja Calif.).	
Parus wollweberi	BRIDLED TITMOUSE
Mts. s.w. U. S. and Mexico (s. to Oaxaca).	
Auriparus flaviceps	VERDIN
S.w. U. S. and n. and w. Mexico (s. to Jalisco).	
Psaltiparus minimus	PLAIN BUSHTIT
W. U. S. and highlands n.w. Mexico (Baja Calif. and Sonora).	
Psaltiparus melanotis⁷	BLACK-EARED BUSHTIT
Highlands s.w. U. S., Mexico and Guatemala.	

Family SITTIDAE NUTHATCHES

Sitta carolinensis	WHITE-BREASTED NUTHATCH
N. Am. to mts. Mexico (s. to Oaxaca).	
Sitta canadensis	RED-BREASTED NUTHATCH
N. Am. and Mexico (Guadalupe I.).	
Sitta pygmaea⁸	PYGMY NUTHATCH
W. N. Am. to highlands cent. Mexico.	

Family CERTHIIDAE CREEPERS

Certhia familiaris⁹	BROWN CREEPER
Eurasia, N. Am., mts. Mexico, Guatemala, Honduras, Nicaragua.	

Family CHAMAEIDAE¹⁰ WRENTITS

Chamaea fasciata	WRENTIT
Pac. U. S. to n.w. Mexico (n.w. Baja Calif.).	

Family CINCLIDAE DIPPERS

Cinclus mexicanus	AMERICAN DIPPER
Rapid mt. streams w. N. Am., and Mid. Am. to w. Panama (except Brit. Honduras and El Salvador).	

⁶ Possibly a race of *P. bicolor*, the Tufted Titmouse of e. and s. U. S.

⁷ Considered by Hellmayr conspecific with *minimus*; if so, Bushtit suffices as the name of the complex.

⁸ Possibly conspecific with *S. pusillus* of s.e. U. S.

⁹ Hellmayr treats the American birds as a distinct species, *C. americana*.

¹⁰ Several recent authors merge this family with the Old World Timaliidae.

Family TROGLODYTIDAE WRENS

Cistothorus platensis	SEDGE WREN
Local: E. N. Am., highlands Mexico, Brit. Honduras, Guatemala, Nicaragua, Costa Rica, w. Panama and S. Am.	
Telmatodytes¹ palustris	MARSH WREN
N. Am.; winters to Mexico.	
Campylorhynchus² albo-brunneus	WHITE-HEADED WREN
Cent. Panama to Colombia.	
Campylorhynchus megalopterus	GRAY-BARRED WREN
Mexico (s. Plateau region).	
Campylorhynchus zonatus	BAND-BACKED WREN
Trop. Mid. Am. (s. Mexico to w. Panama) and n.w. S. Am.	
Campylorhynchus rufinucha^{2a}	RUFOUS-NAPED WREN
Trop. Mid. Am. to n.w. Costa Rica.	
Campylorhynchus chiapensis	GIANT WREN
Mexico (Chiapas).	
Campylorhynchus brunneicapillus	CACTUS WREN
Arid areas s.w. U. S. and Mexico.	
Campylorhynchus jocosus	SPOTTED WREN
N. and w. Mexico.	
Thryothorus ludovicianus	CAROLINA WREN
E. and s. U. S. and n.e. Mexico.	
Thryothorus leucotis	BUFF-BREASTED WREN
Cent. Panama to S. Am.	
Thryothorus modestus	PLAIN WREN
Trop. Mid. Am. to cent. Panama	
Thryothorus zeledoni²	CANE-BRAKE WREN
Caribbean Nicaragua, Costa Rica and n.w. Panama.	
Thryothorus sinaloa	BAR-VENTED WREN
W. Mexico (Sonora and Chihuahua to Guerrero).	
Thryothorus rufalbus	RUFOUS-AND-WHITE WREN
Trop. Mid. Am. (except Brit. Honduras) from s. Mexico (Chiapas) to n. S. Am.	

¹ Hellmayr merges this genus in *Cistothorus*. .

² Hellmayr used *Heleodytes*, and calls most forms of the genus "Cactus Wren". Many are woodland birds, so we reserve that name for *brunneicapillus*.

^{2a} Birds from Chiapas southward, the *C. capistratus* group, may be a separate species, called Rufous-backed Wren by L. I. Davis.

³ Considered by Hellmayr a race of *modestus*.

Thryothorus thoracicus⁴	STRIPE-BREASTED WREN
Nicaragua, Costa Rica and w. Panama.	
Thryothorus leucopogon⁴	STRIPE-THROATED WREN
E. Panama and n.w. S. Am.	
Thryothorus nigricapillus⁵	BLACK-CAPPED WREN
E. Panama and n. w. S. Am.	
Thryothorus castaneus⁵	BAY WREN
Caribbean slope Nicaragua, Costa Rica to cent. Panama (also Pacific slope).	
Thryothorus semibadius⁵	RIVERSIDE WREN
Pacific slope s.w. Costa Rica and s.w. Panama.	
Thryothorus pleurostictus	BANDED WREN
Trop. Mid. Am. (except Brit. Honduras) from s.w. Mexico to n.w. Costa Rica.	
Thryothorus atrogularis	BLACK-THROATED WREN
Caribbean slope Nicaragua, Costa Rica and n.w. Panama.	
Thryothorus spadix⁶	SOOTY-HEADED WREN
Highlands e. Panama (Darien) and n.w. Colombia.	
Thryothorus fasciato-ventris	BLACK-BELLIED WREN
S.w. Costa Rica, Panama and Colombia.	
Thryothorus albinucha	WHITE-BROWED WREN
Mexico (Yucatan Pen.), Guatemala (Peten), Nicaragua.	
Thryothorus felix	HAPPY WREN
W. Mexico (Sonora to Oaxaca, Tres Marias Is.).	
Thryothorus maculipectus⁷	SPOT-BREASTED WREN
Trop. Mexico, Brit. Honduras, Guatemala, El Salvador, Honduras, n. Nicaragua.	
Thryothorus rutilus⁷	RUFOUS-BREASTED WREN
S.w. Costa Rica, Panama, Colombia, Venezuela, Trinidad.	
Thryomanes bewickii	BEWICK'S WREN
U. S. and Mexico (s. to Veracruz and Oaxaca).	

⁴ Hellmayr treats these as one species, calling them *thoracicus*. If so, Stripe-throated Wren may be used for all, as *leucopogon* is plain-breasted.

⁵ Hellmayr treats these as one species, calling them *nigricapillus*. If so, Bay Wren may be used for all, for *semibadius* is not black-capped.

⁶ Hellmayr treats this as a race of *atrogularis*, Black-throated Wren.

⁷ Hellmayr treats these (as well as the *sclateri* and *paucimaculatus* group of s. Colombia to Peru) as one species, using the name *T. rutilus*. The *rutilus* group is bright rufous below, with speckling usually restricted to the throat, the other groups lack rufous and tend to have the speckling everywhere except the throat. If Hellmayr's broad species concept be adopted "Speckled Wren" should be used as the species name.

Thryomanes sissonii	SOCORRO WREN
Mexico (Socorro I.).	
Troglodytes aedon	NORTHERN HOUSE-WREN
N. Am. to n.w. Mexico (mts. Baja Calif.); winters to s. Mexico.	
Troglodytes tanneri	CLARIION ISLAND WREN
Mexico (Clarion I. in Revilla Gigedo group).	
Troglodytes musculus⁸	SOUTHERN HOUSE-WREN
Trop. Mid. Am., Lesser Antilles and S. Am.	
Troglodytes brunneicollis	BROWN-THROATED WREN
Mts. Arizona to s. Mexico (Oaxaca).	
Troglodytes rufociliatus	RUFOUS-BROWED WREN
Mts. s. Mexico (Chiapas), Guatemala, El Salvador and Honduras.	
Troglodytes ochraceus⁹	OCHRACEOUS WREN
Highlands Costa Rica and Panama.	
Thryorchilus browni	TIMBERLINE WREN
Mt. peaks Costa Rica and w. Panama.	
Henicorhina leucosticta	WHITE-BREASTED WOOD-WREN
Trop. Mid. Am. (except El Salvador) and S. Am.	
Henicorhina leucophrys	GRAY-BREASTED WOOD-WREN
Highlands trop. Mid. Am. (unrecorded Brit. Honduras and Nicaragua), and S. Am.	
Uropsila leucogastra	WHITE-BELLIED WREN
E. and s.w. Mexico, Brit. Honduras and Guatemala.	
Salpinctes obsoletus	ROCK WREN
W. N. Am., Mexico, Guatemala, El Salvador, Honduras, Nicaragua and n.w. Costa Rica.	
Catherpes mexicanus	CANYON WREN
Highlands w. N. Am. through Mexico (to Chiapas).	
Hylorchilus sumichrasti	SLENDER-BILLED WREN
Mexico (Veracruz).	
Microcerculus philomela	NIGHTINGALE WREN
S. Mexico (Chiapas), Guatemala, Honduras, Nicaragua, Costa Rica, Panama and n.w. S. Am.	
Cyphorhinus¹⁰ phaeocephalus	SONG WREN
Honduras, Nicaragua, Costa Rica, Panama, Colombia and Ecuador.	

⁸ Includes the *martinicensis* group (to which *beani*, the Cozumel I. form belongs), by some regarded as a separate species, the Antillean House-Wren.

⁹ Hellmayr treats this group as races of the S. Am. *solstitialis*, and suggests that *rufociliatus* and *brunneicollis* may also be conspecific. If so, the technical name would be *brunneicollis*, and "Mountain Wren" may be used for all.

¹⁰ Hellmayr uses *Leucolepis* for this genus.

Family MIMIDAE MOCKINGBIRDS AND THRASHERS

Toxostoma longirostre¹	LONG-BILLED THRASHER
Texas and e. Mexico (s. to Veracruz and Puebla).	
Toxostoma guttatum	COZUMEL THRASHER
Mexico (Cozumel I. off Yucatan Pen.).	
Toxostoma ocellatum	OCELLATED THRASHER
South-central Mexico (Hidalgo and Mexico to Oaxaca).	
Toxostoma cinereum	GRAY THRASHER
Mexico (Baja Calif.).	
Toxostoma bendirei	BENDIRE'S THRASHER
Deserts s.w. U. S. and n.w. Mexico (Sonora); winters to Sinaloa.	
Toxostoma curvirostre	CURVE-BILLED THRASHER
Arid s.w. U. S. and Mexico (to Veracruz and Oaxaca).	
Toxostoma redivivum	CALIFORNIA THRASHER
Calif. and Mexico (Baja Calif.).	
Toxostoma lecontei	LECONTE'S OR DESERT THRASHER
Deserts s.w. U. S. and n.w. Mexico (Sonora and Baja Calif.).	
Toxostoma dorsale	CRISSAL THRASHER
S.w. U. S. and n.w. Mexico (Baja Calif. to Coahuila and Hidalgo).	
Melanotis caerulescens	BLUE MOCKINGBIRD
Mexico (Sonora and Chihuahua, to Veracruz and Oaxaca).	
Melanotis hypoleucus²	BLUE-AND-WHITE MOCKINGBIRD
Highlands s. Mexico (Chiapas), Guatemala, El Salvador and Honduras.	
Melanoptila glabrirostris	BLACK CATBIRD
Mexico (coast of Yucatan Pen. and adjacent islands), Brit. Honduras, Guatemala (Peten), and Honduras.	
Dumetella carolinensis	COMMON CATBIRD
N. Am.; winters through Mid. Am. (except El Salvador) to cent. Panama.	
Mimodes graysoni	SOCORRO THRASHER
Mexico (Socorro I. of Revilla Gigedo Is.).	
Mimus polyglottos	COMMON MOCKINGBIRD
U. S., Bahamas, Greater Antilles, and Mexico (s. to Veracruz and Oaxaca).	
Mimus gilvus	TROPICAL MOCKINGBIRD
Local: trop. Mid. Am. s.e. Mexico, Brit. Honduras, Guatemala, El Salvador, Honduras, cent. Panama (common, introduced [?]), n. S. Am. and s. Lesser Antilles; recorded from Costa Rica.	

¹ *T. rufum*, the Brown Thrasher of e. N. Am. is believed by L. Griscom likely to winter to e. Mexico (Modern Bird Study, p. 109), but there are no positive records.

² Hellmayr and Blake treat this as a race of *caerulescens*; Wetmore disagrees.

Oreoscoptes montanus	SAGE THRASHER
W. N. Am. to n.w. Mexico.	
Donacobius atricapillus	BLACK-CAPPED MOCKINGTHRUSH
E. Panama (Darien) and S. Am.	
Family TURDIDAE THRUSHES	
Turdus migratorius^{2a}	AMERICAN ROBIN
N. Am. and highlands of Mexico (to Veracruz and Oaxaca); winters to Guatemala.	
Turdus rufitorques	RUFOUS-COLLARED ROBIN
Mts. s.e. Mexico (Chiapas), Guatemala, and El Salvador.	
Turdus rufo-palliatus	RUFOUS-BACKED ROBIN
W. Mexico (Sonora to Oaxaca and Tres Marias Is.).	
Turdus assimilis	WHITE-THROATED ROBIN
Trop. Mid. Am. (chiefly in the highlands), w. Colombia and Ecuador.	
Turdus grayi	CLAY-COLORED ROBIN
Trop. Mid. Am. and n. Colombia.	
Turdus fumigatus³	PALE-VENTED ROBIN
Caribbean Costa Rica, Panama (also Pac. slope in eastern part) and S. Am.	
Turdus plebejus⁴	MOUNTAIN ROBIN
Mts. trop. Mid. Am. (except Brit. Honduras) from s. Mexico (Chiapas) to w. Panama.	
Turdus infuscatus	BLACK ROBIN
Mts. e. Mexico (Tamaulipas to Chiapas), Guatemala, El Salvador and Honduras.	
Turdus nigrescens	SOOTY ROBIN
High mts. Costa Rica and w. Panama.	
Ixoreus naevius	VARIED THRUSH
W. N. Am.; winters to Mexico (n. Baja Calif.).	
Myadestes townsendi	TOWNSEND'S SOLITAIRE
Mts. w. N. Am. and n. Mexico (Chihuahua and Durango).	
Myadestes obscurus	BROWN-BACKED SOLITAIRE
Mts. Mexico, Guatemala, El Salvador and Honduras.	
Myadestes melanops⁵	BLACK-FACED SOLITAIRE
Mts. Costa Rica and w. Panama.	
Myadestes coloratus⁶	VARIED SOLITAIRE
Mts. e. Panama (Darien).	

^{2a} Includes *confinis*, San Lucas Robin, of Cape district of Baja California.

³ Hellmayr is followed in including the *obsoletus* group, Costa Rica to Ecuador.

⁴ Hellmayr lumps this group with the lowland S. Am. *ignobilis*, Black-billed Robin; Wetmore disagrees.

⁵ Hellmayr deems these races of *ralloides*, the Andean Solitaire of w. S. Am.

Myadestes unicolor	SLATE-COLORED SOLITAIRE
Mts. s.e. Mexico (San Luis Potosi south), Guatemala, El Salvador, Honduras and Nicaragua.	
Hylocichla⁶ mustelina	WOOD THRUSH
E. N. Am.; winters through Mid. Am. to cent. Panama.	
Hylocichla guttata	HERMIT THRUSH
N. Am. to n.w. Mexico (mts. Baja Calif.); winters to Mexico and Guatemala.	
Hylocichla ustulata	SWAINSON'S OR OLIVE-BACKED THRUSH
N. N. Am.; migrates through Mid. Am.; the eastern populations winter chiefly in S. Am., the western <i>ustulata</i> in Mid. Am. south to Costa Rica.	
Hylocichla minima	GRAY-CHEEKED THRUSH
N. N. Am.; winters in West Indies and n. S. Am.; recorded on migration Mexico (Cozumel I.), Brit. Honduras, Guatemala, Honduras, Costa Rica, Panama.	
Hylocichla fuscescens	VEERY
N. Am.; winters in S. Am.; recorded on migration Mexico (Yucatan, Veracruz), Brit. Honduras, Honduras (Bay Is.), Costa Rica, Panama.	
Catharus dryas	SPOTTED NIGHTINGALE-THRUSH
Highlands Mexico (Chiapas), Guatemala, Honduras and w. S. Am.	
Catharus mexicanus	BLACK-HEADED NIGHTINGALE-THRUSH
Highlands Mid. Am. (except Brit. Honduras and El Salvador) to w. Panama.	
Catharus fuscater	SLATY-BACKED NIGHTINGALE-THRUSH
Highlands Costa Rica, Panama and w. S. Am.	
Catharus occidentalis	RUSSET NIGHTINGALE-THRUSH
Mts. of Mexico (s. to Oaxaca).	
Catharus frantzii⁷	RUDDY-CAPPED NIGHTINGALE-THRUSH
Mts. Mid. Am. (except Brit. Honduras) from s. Mexico (Guerrero eastward) to w. Panama.	
Catharus aurantiirostris⁸	ORANGE-BILLED NIGHTINGALE-THRUSH
Highlands Mid. Am. (except Brit. Honduras) to w. Panama and n. S. Am.	

⁶ Some writers merge this genus in *Catharus*.

⁷ Treated by Hellmayr and Blake as conspecific with *occidentalis*; but cf. Griscom, *Auk*, 54: 198, 1937. If Hellmayr is followed, call the species Russet Nightingale-Thrush.

⁸ Includes the *griseiceps* group of s.w. Costa Rica, Panama and Colombia (see Zimmer, *Auk*, 61: 405, 1944), called Gray-headed Nightingale-Thrush.

Catharus gracilirostris	BLACK-BILLED NIGHTINGALE-THRUSH
High mts. Costa Rica and w. Panama.	
Ridgwayia pinicola	AZTEC THRUSH
High mts. Mexico (Chihuahua to Veracruz and Oaxaca).	
Sialia sialis	COMMON BLUEBIRD
N. Am., Mexico, and highlands Brit. Honduras, Guatemala, El Salvador, Honduras and Nicaragua.	
Sialia mexicana	WESTERN BLUEBIRD
W. N. Am. and highlands Mexico (s. to Michoacan, Puebla and Veracruz).	
Sialia currucoides	MOUNTAIN BLUEBIRD
Mts. w. N. Am. to Mexico (Chihuahua); winters to n.w. Mexico (Sonora and Baja Calif.).	

Family **ZELEDONIIDAE**⁹ WREN-THRUSHES

Zeledonia coronata	WREN-THRUSH
Mts. Costa Rica and w. Panama.	

Family **SYLVIIDAE** OLD WORLD WARBLERS,
KINGLETS AND GNATCATCHERS

Polioptila ¹⁰ <i>caerulea</i>	BLUE-GRAY GNATCATCHER
U. S., Bahamas, Mexico, Guatemala (?); in winter to Brit. Honduras, Guatemala and Honduras (Bay Is.).	
Polioptila albiloris ¹	WHITE-LORED GNATCATCHER
Arid areas s. Mexico (chiefly Pac. slope and Yucatan Pen.), Guatemala, El Salvador, Honduras, Nicaragua and n.w. Costa Rica.	
Polioptila plumbea	TROPICAL GNATCATCHER
Trop. Mid. Am. (except El Salvador) from s.e. Mexico (Quintana Roo) south to S. Am.	
Polioptila nigriceps	BLACK-CAPPED GNATCATCHER
W. Mexico (Sonora to Jalisco).	
Polioptila melanura	BLACK-TAILED GNATCATCHER
S.w. U. S. (chiefly desert) and n.w. Mexico (Baja Calif. to Durango).	
Polioptila schistaceigula	SLATE-THROATED GNATCATCHER
E. Panama (Darien) and n.w. S. Am.	

⁹ Perhaps better included in Turdidae.

¹⁰ This genus and the next two are occasionally placed in a separate family Polioptilidae.

¹ Treated by Hellmayr as a race of *P. plumbea*, the Tropical Gnatcatcher; but Zimmer and Brodkorb consider it a distinct species.

Ramphocaenus rufiventris ²	LONG-BILLED GNATWREN
Trop. Mid. Am. and n.w. S. Am.	
Microbates cinereiventris	HALF-COLLARED GNATWREN
Nicaragua, Costa Rica, Panama and n.w. S. Am.	
Regulus satrapa ³	GOLDEN-CROWNED KINGLET
N. Am. through mts. of Mexico to Guatemala.	
Regulus calendula	RUBY-CROWNED KINGLET
N. Am. and Mexico (Guadalupe I.); winters through Mexico to Guatemala.	

Family MOTACILLIDAE WAGTAILS AND PIPITS

Motacilla alba	PIED WAGTAIL
N. Eurasia; casual in Aleutians and Alaska; accidental once in Mexico (Baja Calif.).	
Anthus spinoletta	WATER PIPIT
Old World and n. N. Am.; winters through Mexico, Guatemala, El Salvador.	
Anthus cervinus	RED-THROATED PIPIT
N. Eurasia; accidental once in Mexico (Baja Calif.).	
Anthus spragueii	SPRAGUE'S PIPIT
Great Plains N. Am.; winters to Mexico (s. to Puebla and Veracruz).	
Anthus lutescens ⁴	YELLOWISH PIPIT
Panama and S. Am.	

Family BOMBYCILLIDAE WAXWINGS

Bombycilla cedrorum	CEDAR WAXWING
N. Am.; winters through Mid. Am. to cent. Panama.	

Family PTILOGONATIDAE⁵ SILKY-FLYCATCHERS

Ptilogonyx cinereus	GRAY SILKY-FLYCATCHER
Mts. Mexico (Chihuahua southwards) and Guatemala.	
Ptilogonyx caudatus	LONG-TAILED SILKY-FLYCATCHER
Mts. Costa Rica and w. Panama.	

Phainopepla nitens	PHAINOPEPLA
S.w. U. S. to s. Mexico (to Puebla and Veracruz).	

² Zimmer considers this conspecific with the S. Am. *melanura*. This genus and the next were formerly placed with the Antwrens of the family Formicariidae.

³ Treated by Hellmayr as conspecific with the Old World *R. regulus*. The genus *Regulus* is placed by Wetmore in a separate family, Regulidae.

⁴ Zimmer considers that *chii* has priority as the specific name.

⁵ By some this family is merged in Bombycillidae.

Phainoptila melanoxantha BLACK-AND-YELLOW SILKY-FLYCATCHER
Mts. Costa Rica and w. Panama.

Family **LANIIDAE** SHRIKES

Lanius ludovicianus LOGGERHEAD SHRIKE
N. Am. and Mexico (s. to Oaxaca).

Family **STURNIDAE** STARLINGS

Sturnus vulgaris COMMON STARLING
N. Old World. Introduced U. S., now N. Am. to n.e. Mexico
(Tamaulipas).

Family **CYCLARHIDAE⁶** PEPPERSHRIKES

Cyclarhis gujanensis RUFOUS-BROWED PEPPERSHRIKE
Trop. Mid. Am. (from s.e. Mexico southward), and S. Am.

Family **VIREOLANIIDAE⁸** SHRIKE-VIREOS

Vireolanius melitophrys CHESTNUT-SIDED SHRIKE-VIREO
Mts. s. cent. Mexico (Michoacan to Veracruz) and Guatemala.

Smaragdolanius pulchellus GREEN SHRIKE-VIREO
S.e. Mexico (Veracruz and Chiapas), Guatemala, Nicaragua, Costa
Rica, Panama.

Smaragdolanius eximus⁷ YELLOW-BROWED SHRIKE-VIREO
E. Panama (Darien) and Colombia.

Family **VIREONIDAE** VIREOS

Vireo atricapillus BLACK-CAPPED VIREO
S.w. U. S.; winters to cent. Mexico.

Vireo bairdi COZUMEL VIREO
Mexico (Cozumel I. off Yucatan Pen.).

Vireo griseus WHITE-EYED VIREO
E. U. S. and n.e. Mexico (to San Luis Potosi and Hidalgo);
winters U. S. to Mexico, Brit. Honduras, Guatemala and Hon-
duras.

Vireo perquisitor⁸ VERACRUZ VIREO
Mexico (Veracruz).

⁶ By some these families are merged in Vireonidae.

⁷ Hellmayr treats this as conspecific with *pulchellus*.

⁸ Treated by Hellmayr as a race of *griseus*, the White-eyed Vireo.

Vireo pallens ⁹	MANGROVE VIREO
Trop. Mid. Am. from Yucatan Pen. and Sonora s. to n.w. Costa Rica.	
Vireo huttoni	HUTTON'S VIREO
W. N. Am., mts. Mexico and Guatemala.	
Vireo carmioli	YELLOW-WINGED VIREO
Mts. Costa Rica and w. Panama.	
Vireo hypochryseus	GOLDEN VIREO
W. Mexico (Sonora and Chihuahua to Oaxaca, Tres Marias Is.).	
Vireo vicinior	GRAY VIREO
S.w. U. S. and n.w. Mexico (s. to Durango).	
Vireo nelsoni ¹⁰	DWARF VIREO
S.w. Mexico (Michoacan).	
Vireo belli	BELL'S VIREO
W. U. S. and n. Mexico (s. to Guanajuato); winters Mexico to Guatemala, El Salvador; recorded in Nicaragua.	
Vireo flavifrons	YELLOW-THROATED VIREO
E. N. Am.; winters through Mid. Am. to Colombia.	
Vireo solitarius	SOLITARY VIREO
Mts. N. Am., Mexico, Brit. Honduras, Guatemala, Honduras and El Salvador; northern birds winter s. U. S. to n. Nicaragua.	
Vireo olivaceus ¹	RED-EYED VIREO
N. Am. and n. Mexico (n.e. Coahuila); migrates through Mid. Am., winters in n. S. Am.	
Vireo flavoviridis ²	YELLOW-GREEN VIREO
Texas and Mid. Am.; winters to n. S. Am.	
Vireo magister	YUCATAN VIREO
Mexico (Yucatan Pen. and adjacent islands), Brit. Honduras, Honduras (Bay Is.).	
Vireo altiloquus	BLACK-WHISKERED VIREO
Florida Keys, West Indies; winters to n. S. Am.; recorded from Caribbean Honduras and Panama.	
Vireo philadelphicus	PHILADELPHIA VIREO
E. N. Am.; winters in Mid. Am. (not recorded Brit. Honduras) to w. Panama.	
Vireo gilvus	WARBLING VIREO
N. Am. to mts. n. and w. Mexico (Chihuahua and Tamaulipas to Guerrero); winters Mexico to Guatemala and El Salvador.	

⁹ Includes the *ochraceus* group (Mexico to e. Nicaragua). All are deemed races of *griseus* by Hellmayr.

¹⁰ Formerly *nanus*. See Bond (Proc. Ac. Nat. Sci. Phila., 86: 399–402, 1934).

¹ Hellmayr uses the name *virescens*.

² Hellmayr and many others consider this, as well as the wide-ranging S. Am. *chivi* group, to be races of the Red-eyed Vireo, *V. olivaceus*.

Vireo leucophrys ³	BROWN-CAPPED VIREO
Mts. s.e. Mexico (s. Veracruz and Chiapas), Costa Rica, w. Panama and S. Am.	
Neochloe brevipennis	SLATY VIREO
Mts. Mexico (Veracruz and Guerrero).	
Hylophilus aurantiifrons	GOLDEN-FRONTED GREENLET
Cent. Panama to n. S. Am.	
Hylophilus flavipes ⁴	SCRUB GREENLET
Costa Rica, Panama and n. S. Am.	
Hylophilus ochraceiceps	TAWNY-CROWNED GREENLET
Trop. Mid. Am. (except El Salvador) and S. Am.	
Hylophilus decurtatus	GRAY-HEADED GREENLET
Trop. Mid. Am. to cent. Panama.	
Hylophilus minor ⁵	LESSER GREENLET
Cent. Panama eastward to Colombia and Ecuador	

Family COEREVIDAE⁶ HONEYCREEPERS

Diglossa baritula	CINNAMON-BELLIED FLOWER-PIERCER
Highlands trop. Mexico, Guatemala, El Salvador and Honduras.	
Diglossa plumbea ⁷	SLATY FLOWER-PIERCER
Highlands Costa Rica and w. Panama.	
Chlorophanes spiza	GREEN HONEYCREEPER
Trop. Mid. Am. (except El Salvador) from Chiapas to S. Am.	
Cyanerpes cyaneus	RED-LEGGED HONEYCREEPER
Trop. Mid. Am., S. Am., Cuba.	
Cyanerpes lucidus ⁸	SHINING HONEYCREEPER
Mexico (Chiapas), Guatemala, Honduras, Nicaragua, Costa Rica, Panama and n.w. Colombia.	
Dacnis cayana	BLUE DACNIS
Nicaragua, Costa Rica, Panama and S. Am.	

³ Zimmer considers this group conspecific with the Warbling Vireo, *V. gilvus*.

⁴ The Costa Rica-Panama form is *viridiflavus*, perhaps a species distinct from the S. Am. *flavipes*. Ridgway used *Pachysylvia* for the genus *Hylophilus*.

⁵ Deemed by Hellmayr conspecific with *decurtatus*. If so, call both Lesser Greenlet, as *minor* is not gray-capped.

⁶ Beecher argues that this family be dropped and that, of the Middle American genera here listed, *Coereba* be transferred to Parulidae and the others to Thraupidae.

⁷ Treated by Hellmayr and Zimmer (together with the S. Am. *sittoides* group) as races of *baritula*. If so, the entire complex may be called Slaty Flower-piercer, for they all have slaty backs but *plumbea* lacks rufous underparts.

⁸ If Hellmayr is correct in treating this as conspecific with the S. Am. Purple Honeycreeper, *caeruleus*, "Yellow-legged Honeycreeper" would be appropriate for the complex; but de Schauensee says races of both *caeruleus* and *lucidus* occur in the same area in n.w. Colombia, near the Panama boundary.

Dacnis viguieri	VIRIDIAN DACNIS
E. Panama (Darien) and n.w. Colombia.	
Dacnis venusta	SCARLET-THIGHED DACNIS
Costa Rica, Panama, Colombia and Ecuador.	
Coereba flaveola	BANANAQUIT
Bahamas, West Indies, trop. Mid. Am. (not reported El Salvador and Honduras) and S. Am., casual in Fla.	

Family PARULIDAE WOOD-WARBLERS

Ateleodacnis⁹ leucogenys	WHITE-EARED CONEBILL
E. Panama (Darien) and Colombia and n.w. Venezuela.	
Mniotilla varia	BLACK-AND-WHITE WARBLER
E. N. Am.; winters through Mid. Am. to n. S. Am.	
Protonotaria citrea	PROTHONOTARY WARBLER
S.e. U. S.; recorded from Mexico (Yucatan Pen. and Cozumel I.), Honduras, and winters in Nicaragua, Costa Rica, Panama and n. S. Am.	
Limnothlypis swainsonii	SWAINSON'S WARBLER
S.e. U. S.; winters in West Indies, Mexico (Yucatan Pen.), Brit. Honduras; recorded from e. Mexico, and off Honduras (Swan I.).	
Helmitheros vermivorus	WORM-EATING WARBLER
E. U. S.; winters in Mid. Am. to e. Panama.	
Vermivora chrysoptera¹⁰	GOLDEN-WINGED WARBLER
E. N. Am.; winters Mid. Am. (not recorded El Salvador or Brit. Honduras) to n. S. Am.	
Vermivora pinus	BLUE-WINGED WARBLER
E. U. S.; winters s.e. Mexico, Guatemala, e. Nicaragua, Costa Rica, Panama, and casually to Colombia.	
Vermivora peregrina	TENNESSEE WARBLER
E. N. Am.; winters Mid. Am. to n. S. Am.	
Vermivora celata	ORANGE-CROWNED WARBLER
W. N. Am. and n.w. Mexico (Todos Santos Is. off Baja Calif.); winters U. S. to Mexico and Guatemala.	
Vermivora ruficapilla	NASHVILLE OR GRAY-CAPPED WARBLER
N. N. Am.; winters Texas to Mexico and Guatemala.	
Vermivora virginiae	VIRGINIA'S WARBLER
Mts. w. U. S.; winters in w. Mexico (s. to Guerrero).	
Vermivora crissalis	COLIMA WARBLER
Mts. s.w. Texas and n.e. Mexico; winters Mexico (s. to Michoacan).	

⁹ Zimmer merges this genus in *Conirostrum*. Hellmayr puts it in Coerebidae.

¹⁰ Brewster's Warbler (*V. leucobronchialis*), a hybrid between *chrysoptera* and *pinus*, has been taken in Costa Rica (Bonilla).

Vermivora luciae	LUCY'S WARBLER
S.w. U. S. and n.w. Mexico (n. Baja Calif.); winters in w. Mexico (to Jalisco).	
Vermivora gutturalis	FLAME-THROATED WARBLER
Mts. Costa Rica and w. Panama.	
Vermivora superciliosa	CRESCENT-CHESTED WARBLER
Highlands Mexico, Guatemala, El Salvador, Honduras and Nicaragua.	
Parula¹⁰ americana	AMERICAN PARULA
E. N. Am.; winters chiefly in West Indies, also e. Mexico, Brit. Honduras, Guatemala, Honduras, El Salvador (once), Nicaragua, Costa Rica (once).	
Parula pitiayumi	TROPICAL PARULA
S. Texas, Mexico, Guatemala, Honduras, Nicaragua, Costa Rica, Panama and S. Am.	
Parula graysoni¹	SOCORRO PARULA
Mexico (Socorro I., Revilla Gigedo group); casually Baja Calif.	
Peucedramus taeniatus²	OLIVE OR TAWNY-HEADED WARBLER
Mts. s.w. U. S., Mexico, Guatemala, El Salvador, Honduras and Nicaragua.	
Dendroica aestiva³	YELLOW WARBLER
N. Am. and Mexico (s. to Michoacan and Morelos), winters through Mid. Am. to S. Am.	
Dendroica petechia⁴	GOLDEN WARBLER
Florida Keys, West Indies, Mexico (Cozumel I. off Yucatan Pen.), Costa Rica (Cocos I.), Galapagos Is., and Pacific coast of s. Colombia to n. Peru.	
Dendroica erithachorides⁵	MANGROVE WARBLER
Coastal mangrove areas on both coasts of trop. Mid. Am. (from Tamaulipas and Baja Calif. southwards) and n. Colombia.	
Dendroica magnolia	MAGNOLIA WARBLER
N. Am.; winters through Mid. Am. to cent. Panama.	
Dendroica tigrina	CAPE MAY WARBLER
N. N. Am.; winters in West Indies; casual in Mexico (Yucatan Pen.), Brit. Honduras, Honduras (Roatan I.).	

¹⁰ Sometimes known as *Compsothlypis*.

¹ Hellmayr treats this as a race of *pitiayumi*.

² Hellmayr uses the name *olivaceus*.

³ Hellmayr treated the *erithachorides* group as races of *petechia*, calling them Golden Warbler. The A.O.U. Check-List Committee now goes further, treating the *aestiva* group as races of *petechia*. If all three groups are deemed conspecific the name Yellow Warbler seems best.

Dendroica caerulescens	BLACK-THROATED BLUE WARBLER
N.e. N. Am.; winters in West Indies, occurs in Mexico (Cozumel I.), Brit. Honduras, in Guatemala, Swan I. off Honduras, Colombia.	
Dendroica coronata	MYRTLE WARBLER
N. N. Am.; winters from U. S. through Mid. Am., casually to Colombia.	
Dendroica auduboni^{2*}	AUDUBON'S WARBLER
Mts. w. N. Am. to n.w. Mexico; winters to Mexico and Guatemala; casual in Costa Rica (sight, C. H. Rogers).	
Dendroica nigrescens	BLACK-THROATED GRAY WARBLER
Mts. w. N. Am. and n.w. Mexico (n. Baja Calif.); winters to Mexico and Guatemala.	
Dendroica townsendi	TOWNSEND'S WARBLER
Mts. n.w. N. Am.; winters to Mexico, Guatemala, El Salvador, Honduras, Nicaragua.	
Dendroica virens	BLACK-THROATED GREEN WARBLER
E. N. Am.; winters through Mid. Am. to Colombia.	
Dendroica chrysoparia	GOLDEN-CHEEKED WARBLER
Highlands Texas; winters Mexico, Guatemala, Honduras, Nicaragua.	
Dendroica occidentalis	HERMIT WARBLER
Mts. w. N. Am.; winters Mexico, Guatemala, El Salvador, Honduras, n. Nicaragua.	
Dendroica cerulea	CERULEAN WARBLER
E. N. Am.; migrates through Mid. Am. (unrecorded El Salvador), winters n. S. Am.	
Dendroica fusca	BLACKBURNIAN WARBLER
E. N. Am.; migrates through Mid. Am., winters from Costa Rica to n.w. S. Am.	
Dendroica dominica	YELLOW-THROATED WARBLER
E. U. S.; the eastern race <i>dominica</i> winters in Fla. and West Indies; <i>albilora</i> winters in Mid. Am. to Costa Rica.	
Dendroica graciae	GRACE'S WARBLER
Mts. s.w. U. S. and highlands Mexico, Brit. Honduras, Guatemala, Honduras, El Salvador and n. Nicaragua.	
Dendroica pensylvanica	CHESTNUT-SIDED WARBLER
E. N. Am.; migrates through Mid. Am. (unrecorded in El Salvador), winters Nicaragua through Panama.	
Dendroica castanea	BAY-BREASTED WARBLER
N.e. N. Am.; recorded through Mid. Am. (except El Salvador),	

* Probably conspecific with *coronata*.

but records few north of Panama; winters in Panama, Colombia and Venezuela.

Dendroica breviunguis⁴ **BLACK-POLLED WARBLER**

N. N. Am.; migrates through West Indies, winters n. S. Am.; once Mexico (Tehuantepec, Oaxaca).

Dendroica pinus **PINE WARBLER**

E. N. Am. and West Indies; the northern race winters to n.e. Mexico (Tamaulipas).

Dendroica discolor **PRairie WARBLER**

E. U. S.; winters in Fla. and West Indies and, at least casually, islands off Caribbean coast of Mexico, Honduras and Nicaragua; once El Salvador.

Dendroica vitellina **VITELLINE WARBLER**

Small Caribbean islets off Central American coast including Honduras (Swan I.).

Dendroica palmarum **PALM WARBLER**

N. N. Am.; winters to West Indies and e. Mexico (Yucatan Pen.), Brit. Honduras, Honduras (Bay Is.).

Seiurus aurocapillus **OVENBIRD**

E. N. Am.; winters through Mid. Am. to Colombia and Venezuela.

Seiurus noveboracensis **NORTHERN WATERTHRUSH**

N. N. Am.; winters in West Indies, through Mid. Am. and n. S. Am.

Seiurus motacilla **LOUISIANA WATERTHRUSH**

E. U. S.; winters in West Indies, and through Mid. Am. to Colombia.

Oporornis formosus **KENTUCKY WARBLER**

E. U. S.; winters through Mid. Am. to n. Colombia.

Oporornis philadelphia⁵ **MOURNING WARBLER**

E. N. Am.; migrates through Mid. Am.; winters Nicaragua s. to n.w. S. Am.

Oporornis tolmiei **MACGILLIVRAY'S WARBLER**

W. N. Am.; winters through Mid. Am. to Panama and Colombia.

Geothlypis trichas **COMMON YELLOWTHROAT**

N. Am. and Mexico (to Veracruz and Oaxaca); winters s. U. S., West Indies, through Mid. Am. to w. Panama, casually to Colombia.

⁴ This is the name under the International Rules, but under A.O.U. Rules the name is *D. striata*. The Tehuantepec specimen, often attributed to *D. castanea* in the literature, is actually a Black-polled Warbler (*fide* H. Friedmann).

⁵ A supposed migrant record of *O. agilis*, the Connecticut Warbler, from Costa Rica turns out to be an error for *O. philadelphia* (*fide* L. Griscom).

Geothlypis beldingi	PENINSULAR YELLOWTHROAT
Mexico (Baja Calif.).	
Geothlypis flavovelata	YELLOW-CROWNED YELLOWTHROAT
E. Mexico (Tamaulipas and n. Veracruz).	
Geothlypis chapalensis	CHAPALA YELLOWTHROAT
Mexico (Lake Chapala and lower Lerma R. in Jalisco).	
Geothlypis chiriquensis*	CHIRIQUI YELLOWTHROAT
W. Panama (base of Chiriqui volcano).	
Geothlypis semiflava	OLIVE-CROWNED YELLOWTHROAT
Caribbean slope Honduras, Nicaragua, Costa Rica and n.w. Panama (Bocas del Toro), w. Colombia and Ecuador.	
Geothlypis speciosa	BLACK-POLLED YELLOWTHROAT
Highland marshes s. cent. Mexico (Michoacan, Puebla, Veracruz).	
Geothlypis nelsoni	HOODED YELLOWTHROAT
Mts. s.e. Mexico (San Luis Potosi to Oaxaca).	
Chamaethlypis poliocephala	GRAY-CROWNED YELLOWTHROAT
Texas, Mid. Am. to s.w. Panama (Chiriqui).	
Icteria virens	YELLOW-BREASTED CHAT
N. Am. and n. Mexico; winters through Mid. Am. to w. Panama.	
Granatellus venustus	RED-BREASTED CHAT
W. Mexico (Sinaloa to Chiapas).	
Granatellus francescae?	TRES MARIAS CHAT
W. Mexico (Tres Marias Is.).	
Granatellus sallaei	GRAY-THROATED CHAT
E. Mexico (Veracruz to Oaxaca and Yucatan Pen.) and Guate- mala.	
Wilsonia citrina	HOODED WARBLER
E. U. S.; winters in West Indies and through Mid. Am. to cent. Panama.	
Wilsonia pusilla	PILEOLATED OR BLACK-CAPPED WARBLER
N. Am.; winters through Mid. Am. (not reported Brit. Honduras).	
Wilsonia canadensis	CANADA WARBLER
E. N. Am.; migrates through Mid. Am.; winters from Honduras (rarely) to n.w. S. Am.	
Cardellina rubrifrons	RED-FACED WARBLER
Mts. s.w. U. S., Mexico and Guatemala.	
Setophaga ruticilla	AMERICAN REDSTART
N. Am.; winters through West Indies, Mid. Am. and n. S. Am.	

* Hellmayr considers this a race of the S. Am. *G. aequinoctialis*, Masked Yellow-throat, which name may be used if he be followed.

? Probably a race of *G. venustus*, and so treated by Blake.

<i>Setophaga picta</i>	PAINTED REDSTART
Mts. Ariz., N. Mex., Mexico, Guatemala, El Salvador, Honduras and Nicaragua.	
<i>Myioborus miniatus</i>	SLATE-THROATED REDSTART
Highlands trop. Mid. Am. and S. Am.	
<i>Myioborus torquatus</i>	COLLARED REDSTART
Mts. Costa Rica and w. Panama.	
<i>Euthlypis lachrymosa</i>	FAN-TAILED WARBLER
Mexico, Guatemala, El Salvador, Honduras, and n.w. Nicaragua.	
<i>Ergaticus ruber</i>	RED WARBLER
Mts. Mexico (Sinaloa and Chihuahua to Veracruz and Oaxaca).	
<i>Ergaticus versicolor</i>⁸	PINK-HEADED WARBLER
Mts. s. Mexico (Chiapas) and Guatemala.	
<i>Basileuterus tristriatus</i>⁹	THREE-STRIPED WARBLER
Highlands Costa Rica, Panama and S. Am.	
<i>Basileuterus culicivorus</i>	GOLDEN-CROWNED WARBLER
Mid. Am. (chiefly highlands) and S. Am.	
<i>Basileuterus melanogenys</i>¹⁰	BLACK-CHEEKED WARBLER
Mts. Costa Rica and Panama.	
<i>Basileuterus belli</i>	GOLDEN-BROWED WARBLER
Mexico, Guatemala, El Salvador and Honduras.	
<i>Basileuterus rufifrons</i>¹	RUFOUS-CAPPED WARBLER
Chiefly highlands Mexico, Brit. Honduras, Guatemala and Honduras.	
<i>Basileuterus delatrii</i>²	CHESTNUT-CAPPED WARBLER
W. Guatemala, El Salvador, Honduras, Nicaragua, Costa Rica, Panama and Colombia.	
<i>Basileuterus fulvicauda</i>³	BUFF-RUMPED WARBLER
Honduras, Nicaragua, Costa Rica and Panama and w. S. Am.	

⁸ Hellmayr treats this as conspecific with *E. ruber*, the Red Warbler.

⁹ Includes *B. tacarcunae*, the Tacarcuna Warbler of Mt. Tacarcuna, Darien, Panama.

¹⁰ Includes *B. ignotus*, the Pirri Warbler, of Mt. Pirri, Darien, Panama.

¹ Includes *B. salvini*, Salvin's Warbler, of s.e. Mexico (from s. Veracruz e.) to Brit. Honduras, n. cent. Guatemala and n. Honduras, which some authors regard as entitled to specific rank because of apparent overlap in range with *rufifrons*. True *rufifrons* (including its undoubted subspecies) ranges more widely in the highlands of Mexico and into w. Guatemala.

² Van Rossem suggests that *delatrii* may be conspecific with *rufifrons*; on that view, use Rufous-capped Warbler for the complex.

³ Zimmer treats this group as races of *rivularis*, of e. S. Am.; on that view, use River Warbler for the complex.

Family PLOCEIDAE⁴ HOUSE SPARROWS AND WEAVERS

Passer domesticus HOUSE SPARROW
Old World, introduced N.Am., West Indies, Mexico, s. S. Am.

Family ICTERIDAE AMERICAN ORIOLES AND BLACKBIRDS

Zarhynchus wagleri	CHESTNUT-HEADED OROPENDOLA
Trop. Mid. Am. (except El Salvador), Colombia and Ecuador.	
Gymnostinops montezuma	MONTEZUMA OROPENDOLA
Trop. Mid. Am. (except El Salvador) to central Panama.	
Gymnostinops guatimozinus	BLACK OROPENDOLA
E. Panama and Colombia.	
Psarocolius⁵ decumanus	CRESTED OROPENDOLA
Panama and S. Am.	
Cacicus cela⁶	YELLOW-RUMPED CACIQUE
Panama and S. Am.	
Cacicus uropygialis⁷	SCARLET-RUMPED CACIQUE
Nicaragua, Costa Rica, Panama and n.w. S. Am.	
Cassiculus melanicterus	YELLOW-WINGED CACIQUE
W. Mexico (s. Sonora to Chiapas).	
Amblycercus holosericeus	YELLOW-BILLED CACIQUE
Mid. Am. and n. and w. S. Am.	
Psomocolax⁸ oryzivorus	GIANT COWBIRD
Trop. Mid. Am. (except El Salvador), and S. Am.	
Tangavarius aeneus	BRONZED OR RED-EYED COWBIRD
S.w. U. S. and Mid. Am. to w. Panama.	
Molothrus ater	BROWN-HEADED COWBIRD
N. Am. and Mexico (s. to Veracruz and Oaxaca).	
Molothrus bonariensis	SHINY COWBIRD
E. Panama (one record, Darien), S. Am., Lesser Antilles.	
Quiscalus quiscula	PURPLE GRACKLE
E. N. Am.; winters to s. Texas; the race <i>aeneus</i> reported from Mexico (Tamaulipas, sight H. C. Oberholser).	

⁴ Some authorities place *Passer* and its close allies in a separate family, Passeridae.

⁵ Hellmayr uses the name *Xanthornus*; *Ostินops* has generally been used, but *Psarocolius* appears to have priority (Miller, *Auk*, 64: 373, 1947).

⁶ Includes *vittellinus* of Panama and n.w. Colombia, often treated as a distinct species. If considered distinct, Saffron-rumped Cacique may be used.

⁷ Includes *microrhynchus* of Nicaragua to e. Panama, perhaps specifically distinct, and called Small-billed Cacique. An undoubted race of *uropygialis* occurs on the Pacific coast of Darien.

⁸ Parkes contends that *Scaphidura* has priority (*Condor*, 56: 229, 1954).

Cassidix mexicanus	BOAT-TAILED GRACKLE
S. U. S., Mid. Am. and Colombia.	
Cassidix palustris	SLENDER-BILLED GRACKLE
Mexico (marshes near Mexico City).	
Cassidix nicaraguensis	NICARAGUAN GRACKLE
Nicaragua (Lakes Nicaragua and Managua).	
Euphagus carolinus	RUSTY BLACKBIRD
N. N. Am.; in winter accidental to n.w. Mexico (Baja Calif.).	
Euphagus cyanocephalus	BREWER'S BLACKBIRD
W. N. Am. and n.w. Mexico (n. Baja Calif.); winters to Mexico and Guatemala.	
Dives dives	MELODIOUS BLACKBIRD
E. Mexico, Brit. Honduras, Guatemala, Honduras and Nicaragua. ⁹	
Icterus spurius¹⁰	ORCHARD ORIOLE
E. U. S.; winters through Mid. Am. to n. S. Am.	
Icterus fuertesi¹¹	OCHRE ORIOLE
E. Mexico (Tamaulipas, Veracruz).	
Icterus prosthemelas	BLACK-COWLED ORIOLE
Trop. Mid. Am. (except El Salvador) to n.w. Panama (Bocas del Toro). ²	
Icterus wagleri	BLACK-VENTED ORIOLE
Highlands Mexico, Guatemala, El Salvador, Honduras and n. Nicaragua.	
Icterus parisorum	SCOTT'S ORIOLE
S.w. U. S. and Mexico (highlands).	
Icterus maculi-alatus	BAR-WINGED ORIOLE
Mexico (Chiapas), Guatemala and El Salvador.	
Icterus cucullatus	HOODED ORIOLE
S.w. U. S., Mexico, Brit. Honduras.	
Icterus auricapillus	ORANGE-CROWNED ORIOLE
E. Panama (Darien) and n. S. Am.	
Icterus mesomelas	YELLOW-TAILED ORIOLE
Trop. Mid. Am. (except El Salvador) and n. and w. S. Am.	
Icterus chrysater	YELLOW-BACKED ORIOLE
Trop. Mid. Am. and n. S. Am.	

⁹ Hellmayr considers *warszewiczi* and *kalinowskii* of s.w. Ecuador and w. Peru races of *D. dives*.

¹⁰ This and the following seven species are placed in the genus *Bananivorus* by Beecher (Wilson Bull., 62: 60-68, 1950).

¹¹ Perhaps a race of the Orchard Oriole, *I. spurius*.

² The Bahaman *northropi* is treated as a subspecies by Hellmayr, but Bond and Beecher regard it as a race of the Greater Antillean *dominicensis*, which may be conspecific with *prosthemelas*, and is the earlier name.

<i>Icterus nigrogularis</i> ³	YELLOW ORIOLE
N. S. Am. (chiefly arid Caribbean) and Panama (?) (one doubtful record).	
<i>Icterus graduacauda</i> ⁴	BLACK-HEADED ORIOLE
Texas, Mexico (Tamaulipas and highlands) and n.w. Guatemala.	
<i>Icterus pectoralis</i>	SPOT-BREASTED ORIOLE
Arid. s.e. Mexico (Oaxaca and Chiapas), Guatemala, Honduras, El Salvador, Nicaragua and n.w. Costa Rica.	
<i>Icterus gularis</i>	BLACK-THROATED ORIOLE
Texas and Mid. Am. s. to Nicaragua.	
<i>Icterus galbula</i>	BALTIMORE ORIOLE
E. N. Am.; winters through Mid. Am. to Colombia.	
<i>Icterus bullockii</i> ⁵	BULLOCK'S ORIOLE
W. N. Am. to cent. highlands of Mexico; winters to Guatemala.	
<i>Icterus auratus</i>	ORANGE ORIOLE
Mexico (Yucatan and Meco I.).	
<i>Icterus pustulatus</i> ⁶	FLAME-HEADED ORIOLE
Mexico (Sonora to Veracruz and Guerrero); recorded in s. Arizona.	
<i>Icterus graysonii</i> ⁶	TRES MARIAS ORIOLE
W. Mexico (Tres Marias Is.).	
<i>Icterus sclateri</i> ⁶	STREAK-BACKED ORIOLE
Trop. Mid. Am. (except Brit. Honduras and Panama) from s. Mexico (Oaxaca and Chiapas) to n.w. Costa Rica.	
<i>Agelaius tricolor</i>	TRICOLORED BLACKBIRD
Pacific U. S. and n.w. Mexico (n. Baja Calif.).	
<i>Agelaius phoeniceus</i>	RED-WINGED BLACKBIRD
N. Am., Bahamas, Cuba, Mexico, Guatemala, El Salvador, Honduras, w. Nicaragua and n.w. Costa Rica.	
<i>Xanthocephalus xanthocephalus</i>	YELLOW-HEADED BLACKBIRD
W. N. Am. and n.w. Mexico; winters in Mexico.	
<i>Leistes militaris</i>	RED-BREASTED BLACKBIRD
Panama and S. Am.	
<i>Sturnella magna</i>	COMMON MEADOWLARK
E. and s. N. Am., Cuba, Mid. Am. and n. S. Am.	

³ Formerly called *xanthornus*.

⁴ Formerly called *melanocephalus*.

⁵ Includes *abeillei*, the Black-backed Oriole, of s. Mexican plateau (San Luis Potosi to Jalisco and Veracruz).

⁶ Hellmayr and Blake treat *graysonii* and members of the *sclateri* group as races of *pustulatus*. On that view Streak-backed Oriole may be used for the complex, for the heads of the southern races are neither scarlet nor even orange.

<i>Sturnella neglecta</i>	WESTERN MEADOWLARK
W. N. Am. and n. Mexico; winters in Mexico (s. to Jalisco, Guanajuato and Veracruz).	
<i>Dolichonyx oryzivorus</i>	BOBOLINK
N. Am.; migrates chiefly through West Indies; recorded on migration Mexico (Yucatan Pen. and Cozumel I.), Brit. Honduras, Honduras, Nicaragua, Costa Rica, Panama; winters in S. Am.	

Family TERSINIDAE SWALLOW-TANAGERS

<i>Tersina viridis</i>	SWALLOW-TANAGER
E. Panama (Darien) and S. Am.	

Family THRAUPIDAE TANAGERS

<i>Chlorophonia occipitalis</i>	BLUE-CROWNED CHLOROPHONIA
Mts. trop. Mexico, Guatemala, El Salvador, Honduras, and n. Nicaragua.	
<i>Chlorophonia callophrys</i>⁷	GOLDEN-BROWED CHLOROPHONIA
Mts. Costa Rica and w. Panama.	
<i>Tanagra elegantissima</i>⁸	BLUE-HOODED EUPHONIA
Highlands Mid. Am. to w. Panama.	
<i>Tanagra xanthogaster</i>	ORANGE-BELLIED EUPHONIA
E. Panama (Darien) and S. Am.	
<i>Tanagra anneae</i>	TAWNY-CAPPED EUPHONIA
Highlands Costa Rica and Panama.	
<i>Tanagra fulvicrissa</i>	FULVOUS-VENTED EUPHONIA
Costa Rica, Panama and n.w. S. Am.	
<i>Tanagra minuta</i>	WHITE-VENTED EUPHONIA
Guatemala, Brit. Honduras, Honduras, Nicaragua, Costa Rica, Panama and S. Am.	
<i>Tanagra godmani</i>⁹	PALE-VENTED EUPHONIA
W. Mexico (Sonora to Colima).	
<i>Tanagra affinis</i>	SCRUB EUPHONIA
Trop. Mid. Am. from e. Mexico to n.w. Costa Rica (chiefly in semi-arid areas).	

⁷ Treated by Hellmayr as a race of *occipitalis*. If so, Blue-crowned Chlorophonia may stand as the name of the entire complex.

⁸ Treated by Hellmayr (and Blake) as conspecific with the West Indian *musica* group, in which he also includes several S. Am. forms.

⁹ Blake considers this a race of *T. affinis*, the Scrub Euphonie.

Tanagra luteicapilla	YELLOW-CROWNED EUPHONIA
E. Nicaragua, Costa Rica and Panama.	
Tanagra laniirostris	THICK-BILLED EUPHONIA
Costa Rica, Panama and S. Am.	
Tanagra lauta	YELLOW-THROATED EUPHONIA
Trop. Mid. Am. (chiefly in the highlands) to w. Panama (Chiriqui).	
Tanagra gouldi	OLIVE-BACKED EUPHONIA
Trop. Mid. Am. (except El Salvador) to n.w. Panama (Bocas del Toro).	
Tanagra imitans¹⁰	SPOT-CROWNED EUPHONIA
S. Costa Rica and w. Panama (Chiriqui).	
Tangara¹ florida	EMERALD TANAGER
Costa Rica, Panama and Colombia.	
Tangara chrysophrus	SPECKLED² TANAGER
Costa Rica, Panama and n. S. Am.	
Tangara icterocephala	SILVER-THROATED TANAGER
Highlands Costa Rica, Panama and n.w. S. Am.	
Tangara larvata³	GOLDEN-MASKED TANAGER
Trop. Mid. Am. (except El Salvador) and n.w. S. Am.	
Tangara cabanisi	AZURE-RUMPED TANAGER
S. Mexico (Chiapas) and Guatemala.	
Tangara palmeri	GRAY-AND-GOLD TANAGER
E. Panama (Darien), w. Colombia and Ecuador.	
Tangara inornata	PLAIN-COLORED TANAGER
Cent. Panama to n.w. Colombia.	
Tangara gyrota	BAY-HEADED TANAGER⁴
Costa Rica, Panama and trop. S. Am.	
Tangara lavinia	RUFOUS-WINGED TANAGER
Guatemala, Honduras, Nicaragua, Costa Rica, Panama and n.w. S. Am.	
Tangara dowii	SPANGLE-CHEEKED TANAGER
Mts. Costa Rica and w. Panama.	

¹⁰ Formerly known as *gracilis*.

¹ This genus was formerly called *Calliste*, and by Ridgway and Hellmayr *Calospiza*.

³ This much more suitable name, suggested by Skutch, replaces Ridgway's "Yellow-browed" Tanager, which name Ridgway and Hellmayr also used for another Middle American species *Chlorothraupis olivacea*. "Spotted" Tanager has been used for several other species.

[•] Treated by Hellmayr and Blake as conspecific with the blue-headed *T. nigro-cincta* of S. Am. east of the Andes. On that view the species may be called Masked Tanager.

⁴ "Blue-rumped Green" Tanager, used for one of the Middle American races, is not appropriate for the species, as several races, including *gyrota*, have green rumps.

Tangara fucosa	GREEN-NAPED TANAGER
Mts. e. Panama (Mt. Pirri, Darien).	
Bangsia arcaei	BLUE-AND-GOLD TANAGER
Foothills Carib. Costa Rica and w. Panama (Veraguas).	
Thraupis episcopus⁵	BLUE-GRAY TANAGER
Trop. Mid. Am. and S. Am.	
Thraupis abbas	YELLOW-WINGED TANAGER
Trop. Mid. Am. from Mexico to n. Nicaragua.	
Thraupis palmarum	PALM TANAGER
Nicaragua, Costa Rica, Panama and trop. S. Am.	
Spindalis zena	STRIPE-HEADED TANAGER
Bahamas, Greater Antilles, and Mexico (Cozumel I.).	
Ramphocelus dimidiatus	CRIMSON-BACKED TANAGER
Panama and n.w. S. Am.	
Ramphocelus passerinii	SCARLET-RUMPED TANAGER
Trop. Mid. Am. (except El Salvador) to w. Panama.	
Ramphocelus icteronotus	YELLOW-RUMPED TANAGER
Panama, w. Colombia and Ecuador.	
Phlogothraupis sanguinolenta	CRIMSON-COLLARED TANAGER
Trop. Mid. Am. (except El Salvador) to n.w. Panama (Bocas del Toro).	
Piranga rubra	SUMMER TANAGER
S. U. S. and n. Mexico; winters through Mid. Am. to S. Am.	
Piranga flava	HEPATIC TANAGER
Highlands s.w. U. S. and Mid. Am. through most of S. Am.	
Piranga roseo-gularis	ROSE-THROATED TANAGER
Mexico (Yucatan Pen. and adjacent islands) and Guatemala (Peten).	
Piranga olivacea	SCARLET TANAGER
E. N. Am.; migrates through Mid. Am.; winters S. Am.	
Piranga leucoptera	WHITE-WINGED TANAGER
Highlands trop. Mid. Am. to w. Panama; also S. Am.	
Piranga ludoviciana	WESTERN TANAGER
W. N. Am. and n.w. Mexico; winters to Mexico, Guatemala, El Salvador, Honduras, Costa Rica (Miravalles).	
Piranga bidentata	FLAME-COLORED TANAGER
Mts. trop. Mid. Am. (not reported Brit. Honduras) to w. Panama.	
Piranga erythrocephala	RED-HEADED TANAGER
W. Mexico (Sonora and Chihuahua to Oaxaca).	

* By some called *virens*.

<i>Chlorothraupis olivacea</i>	LEMON-BROWED TANAGER⁶
E. Panama (Darien), w. Colombia and Ecuador.	
<i>Chlorothraupis carmioli</i>	OLIVE TANAGER
Nicaragua, Costa Rica, Panama, Peru.	
<i>Habia rubica</i>	RED-CROWNED ANT-TANAGER
Trop. Mid. Am. and S. Am.	
<i>Habia fuscicauda</i>⁷	DUSKY-TAILED ANT-TANAGER
Trop. Mid. Am. and Colombia.	
<i>Habia atrimaxillaris</i>⁷	BLACK-CHEEKED ANT-TANAGER
Pac. slope s. Costa Rica (Osa Pen. and vicinity).	
<i>Lanio aurantius</i>⁸	BLACK-THROATED SHRIKE-TANAGER
Trop. s. Mexico, Brit. Honduras, Guatemala, Honduras.	
<i>Lanio leucothorax</i>⁸	WHITE-THROATED SHRIKE-TANAGER
Nicaragua, Costa Rica and w. Panama.	
<i>Tachyphonus rufus</i>	WHITE-LINED TANAGER
Costa Rica, Panama and trop. S. Am.	
<i>Tachyphonus luctuosus</i>	WHITE-SHOULDERED TANAGER
Honduras, Nicaragua, Costa Rica, Panama and n.w. S. Am.	
<i>Tachyphonus delatrii</i>	TAWNY-CRESTED TANAGER
Nicaragua, Costa Rica, Panama and n.w. S. Am.	
<i>Heterospingus rubrifrons</i>⁹	SULPHUR-RUMPED TANAGER
Costa Rica and Panama.	
<i>Eucometis penicillata</i>	GRAY-HEADED TANAGER
Trop. Mid. Am. (except El Salvador) to trop. S. Am.	
<i>Mitrospingus cassinii</i>	DUSKY-FACED TANAGER
Costa Rica, Panama and n.w. S. Am.	
<i>Rhodinocichla rosea</i>	ROSE-BREASTED THRUSH-TANAGER
Trop. w. Mexico (Sinaloa to Colima), Costa Rica, Panama and n. S. Am.	
<i>Hemithraupis flavicollis</i>	YELLOW-BACKED TANAGER
E. Panama (Darien) and trop. S. Am.	
<i>Chrysotlypis chrysomelas</i>	BLACK-AND-YELLOW TANAGER
Highlands Costa Rica and Panama.	

⁶ "Yellow-browed Tanager", used by Ridgway and Hellmayr for this bird, was also used by Ridgway and Skutch for *Tangara chrysophrys*. To avoid confusion a modification of the name seems desirable.

⁷ Hellmayr and others consider these forms races of *H. gutturalis* of the Magdalena Valley of Colombia. If so, the species name should be Red-throated Ant-Tanager.

⁸ Hellmayr treats these as conspecific under the name *aurantius*; if so the name should be Great Shrike-Tanager.

⁹ Regarded by Hellmayr as a "hen-feathered" race of *H. xanthopygius* of n.w. S. Am.

Chlorospingus ophthalmicus ¹⁰	COMMON BUSH-TANAGER
Highlands trop. Mid. Am. (except Brit. Honduras), and S. Am.	
Chlorospingus pileatus	SOOTY-CAPPED BUSH-TANAGER
Mts. Costa Rica and w. Panama.	
Chlorospingus zeledoni ¹	VOLCANO BUSH-TANAGER
Mts. Costa Rica (Irazu and Turrialba volcanoes).	
Chlorospingus inornatus	MOUNT PIRRI BUSH-TANAGER
Highlands e. Panama (Darien).	
Chlorospingus flavigularis	YELLOW-THROATED BUSH-TANAGER
Highlands w. Panama (chiefly Carib. slope), n.w. S. Am.	
Chlorospingus canigularis	ASHY-THROATED BUSH-TANAGER
Highlands Carib. slope Costa Rica, and n.w. S. Am.	

Family **FRINGILLIDAE** FINCHES, GROSBEAKS, BUNTINGS

Saltator atriceps	BLACK-HEADED SALTATOR
Trop. Mid. Am. to cent. Panama.	
Saltator maximus	BUFF-THROATED SALTATOR
Trop. Mid. Am. (except El Salvador) and S. Am.	
Saltator coerulescens	GRAYISH SALTATOR
Trop. Mid. Am. to cent. Costa Rica; also n. and e. S. Am.	
Saltator albicollis	STREAKED SALTATOR
S.w. Costa Rica, Panama, n. and w. S. Am. and Lesser Antilles.	
Rhodothraupis celaeno	CRIMSON-COLLARED GROSBEAK
E. Mexico (s. Nuevo Leon to Veracruz and Puebla).	
Caryothraustes canadensis	GREEN GROSBEAK
E. Panama (Darien) and trop. S. Am. e. of Andes.	
Caryothraustes poliogaster	BLACK-FACED GROSBEAK
Trop. Mid. Am. (except El Salvador) to cent. Panama.	
Pitylus grossus	SLATE-COLORED GROSBEAK
Nicaragua, Costa Rica, Panama, S. Am.	
Richmondena cardinalis ²	COMMON CARDINAL
E. and s. U. S., Mexico and Brit. Honduras.	
Pyrrhuloxia sinuata	PYRRHULOXIA
S.w. U. S. and w. Mexico (s. to Puebla).	
Pheucticus chrysopaeplus	YELLOW GROSBEAK
Highlands w. Mexico and Guatemala.	

¹⁰ Zimmer is followed in treating *punctulatus* of w. Panama and *tacarcunae* of e. Panama as races of *ophthalmicus*; Hellmayr treated *tacarcunae* as a race of *flavigularis*.

¹ Carriker has suggested that this may be a color phase of *pileatus*.

² Includes *carnea* of s.w. Mexico (Colima to Oaxaca), by some regarded as a distinct species, which may be called Long-crested Cardinal.

Pheucticus tibialis ³	BLACK-THIGHED GROSBEAK
Highlands Costa Rica and w. Panama.	
Pheucticus ludovicianus	ROSE-BREASTED GROSBEAK
E. N. Am.; winters through Mid. Am. from s. Mexico to Colombia.	
Pheucticus melanocephalus	BLACK-HEADED GROSBEAK
W. N. Am., n. and w. Mexico (chiefly in mts., Baja Calif. to Tamaulipas s. to Oaxaca and Veracruz); winters in Mexico.	
Guiraca caerulea	BLUE GROSBEAK
S. U. S. and Mid. Am. (except Brit. Honduras) to n.w. Costa Rica; northern birds winter through Mid. Am. to w. Panama.	
Cyanocompsa cyanooides	BLUE-BLACK GROSBEAK
Trop. Mid. Am. (except El Salvador) and S. Am.	
Cyanocompsa parellina	BLUE BUNTING
Trop. Mexico, Guatemala, El Salvador, Honduras and Nicaragua.	
Passerina cyanea	INDIGO BUNTING
E. N. Am.; winters through Mid. Am. to cent. Panama.	
Passerina amoena	LAZULI BUNTING
W. N. Am. and n.w. Mexico; winters in Mexico.	
Passerina versicolor	VARIED BUNTING
S.w. U. S., Mexico and Guatemala.	
Passerina rositae	ROSE-BELLIED BUNTING
S. Mexico (Oaxaca and Chiapas).	
Passerina ciris	PAINTED BUNTING
S. U. S. and n. Mexico; winters through Mid. Am. to w. Panama.	
Passerina leclancherii	ORANGE-BREASTED BUNTING
S.w. Mexico (Colima to Chiapas).	
Tiaris olivacea	YELLOW-FACED GRASSQUIT
Greater Antilles, trop. Mid. Am. (not reported Brit. Honduras) and n. S. Am.	
Spiza americana	DICKCISSEL
Cent. N. Am.; migrates through Mid. Am.; winters from s. Mexico to n. S. Am.	
Pinaroloxias inornata	COCOS ISLAND FINCH
Costa Rica (Cocos Island).	
Hesperiphona vespertina	EVENING GROSBEAK
N. N. Am. and in mts. to Mexico (s. to Veracruz and Oaxaca).	
Hesperiphona abeillei	HOODED GROSBEAK
Mts. Mexico (Chihuahua and Tamaulipas s.) to Guatemala.	
Carpodacus purpureus	PURPLE FINCH
N. N. Am. and in mts. to w. Mexico (n. Baja Calif.).	

³ Hellmayr considers this and several S. Am. forms to be races of the Yellow Grosbeak, *P. chrysopeplus*.

Carpodacus cassini	CASSIN'S FINCH
Mts. w. N. Am. to w. Mexico (n. Baja Calif.); winters to s. cent. Mexico.	
Carpodacus mexicanus⁴	HOUSE FINCH
W. N. Am. to Mexico (chiefly in west and in highlands elsewhere).	
Sporophila schistacea	SLATE-COLORED SEEDEATER
Costa Rica, Panama and n. S. Am.	
Sporophila torqueola	WHITE-COLLARED SEEDEATER
S. Texas and Mid. Am. s. to Costa Rica.	
Sporophila aurita⁵	VARIABLE SEEDEATER
Trop. Mid. Am. (except El Salvador) and n.w. S. Am.	
Sporophila nigricollis⁶	YELLOW-BELLIED SEEDEATER
S.w. Costa Rica, Panama, S. Am. and s. Lesser Antilles.	
Sporophila minuta	RUDDY-BREASTED SEEDEATER
Pac. slope Mexico, Guatemala, El Salvador, Nicaragua, Costa Rica, Panama, and S. Am.	
Amaurospiza relicta⁷	SLATE-BLUE SEEDEATER
Mexico (Mts. Guerrero).	
Amaurospiza concolor	BLUE SEEDEATER
Nicaragua, Costa Rica, Panama and n.w. S. Am.	
Oryzoborus nuttingi⁸	NICARAGUAN SEED-FINCH
Caribbean slope of Nicaragua.	
Oryzoborus funereus⁹	THICK-BILLED SEED-FINCH
Trop. Mid. Am. (except El Salvador) and n.w. S. Am.	
Volatinia jacarina	BLUE-BLACK GRASSQUIT
Trop. Mid. Am., S. Am., s. Lesser Antilles (Grenada).	
Spinus pinus	PINE SISKIN
N. N. Am. and mts. of Mexico (s. to Chiapas).	
Spinus atriceps	BLACK-CAPPED SISKIN
Mts. s. Mexico (Chiapas) and w. Guatemala.	
Spinus notatus	BLACK-HEADED SISKIN
Mts. s.e. Mexico, Guatemala, El Salvador, Honduras and Nicaragua.	

⁴ The San Benito House Finch (*mccgregori*) of San Benito I., and the Guadalupe House Finch (*amplus*) of Guadalupe I., Mexico are by some ornithologists regarded as distinct species. On that view *mexicanus* would be called Common House Finch.

⁵ Includes *corvina*, Carib. slope of Mid. Am. to n.w. Panama, sometimes deemed a distinct species, the Black Seedeater. All are regarded by de Schauensee as races of the S. Am. *S. americana*.

⁶ Formerly called *S. gutturalis*.

⁷ Described as *Amaurospizopsis relicta*; closely related to *A. concolor*.

⁸ Treated by Hellmayr as a race of the S. Am. *O. crassirostris*, Large-billed Seed-Finch.

⁹ Regarded by de Schauensee as a race of *O. angolensis* of n. and e. S. Am.

Spinus xanthogaster	YELLOW-BELLIED SISKIN
Mts. Costa Rica, w. Panama and n.w. S. Am.	
Spinus tristis	AMERICAN GOLDFINCH
N. Am. and n.w. Mexico; winters to Mexico. ¹⁰	
Spinus psaltria	LESSER OR DARK-BACKED GOLDFINCH
W. N. Am., Mid. Am. (not reported Brit. Honduras) and n.w. S. Am.	
Spinus lawrencei	LAWRENCE'S GOLDFINCH
California to n.w. Mexico (n. Baja Calif.); winters to n.w. Mexico.	
Loxia curvirostra	RED CROSSBILL
Holarctic region, in America follows coniferous forests (chiefly in mts.) through Mexico, Brit. Honduras, Guatemala, Honduras and n. Nicaragua.	
Sicalis flaveola	SAFFRON FINCH
S. Am.; introduced in Jamaica, cent. Panama (Gatun, photographed, R. T. Scholes, seen E. Eisenmann).	
Sicalis luteola	YELLOW GRASS-FINCH
S.e. Mexico, Guatemala, Panama (Coclé), S. Am., s. Lesser Antilles.	
Spodiornis rusticus	SLATY FINCH
Highlands s. Mexico (Veracruz, Chiapas), Costa Rica, w. Panama, n. and w. S. Am.	
Acanthidops bairdi	PEG-BILLED FINCH
Mts. Costa Rica.	
Pezopetes capitalis	LARGE-FOOTED FINCH
Mts. Costa Rica and w. Panama (Chiriquí).	
Pselliophorus tibialis	YELLOW-THIGHED FINCH
Highlands Costa Rica and w. Panama (w. Chiriquí).	
Pselliophorus luteoviridis	YELLOW-GREEN FINCH
Highlands w. Panama (Cerro Flores, e. Chiriquí).	
Atlapetes pileatus	RUFous-CAPPED BRUSH-FINCH
Mts. of Mexico (Chihuahua and Tamaulipas to Guerrero).	
Atlapetes albinucha	WHITE-NAPED BRUSH-FINCH
Highlands s.e. Mexico (Veracruz to Chiapas), also Colombia.	
Atlapetes gutturalis	YELLOW-THROATED BRUSH-FINCH
Highlands trop. Mid. Am. (except Brit. Honduras) Chiapas to N. S. Am.	
Atlapetes apertus¹	PLAIN-BREASTED BRUSH-FINCH
Highlands s.e. Mexico (s. Veracruz).	

¹⁰ The one Guatemalan record appears to be an immature *S. psaltria* (*fide* A. R. Phillips).

¹ Perhaps a race of *brunnei-nucha* (see Parkes, Condor, 56: 132, 1954).

Atlapetes brunnei-nucha	CHESTNUT-CAPPED BRUSH-FINCH
Highlands trop. Mid. Am. (except Brit. Honduras), and n. S. Am.	
Atlapetes virenticeps²	GREEN-STRIPED BRUSH-FINCH
S. part of Mexican plateau.	
Atlapetes assimilis²	GRAY-STRIPED BRUSH-FINCH
Highlands s.w. Costa Rica and w. Panama, also n.w. S. Am.	
Atlapetes atricapillus	BLACK-HEADED BRUSH-FINCH
Highlands e. Panama (Darien) and Colombia.	
Lysurus crassirostris	SOOTY-FACED FINCH
Highlands Costa Rica and w. Panama.	
Arremon aurantiirostris	ORANGE-BILLED SPARROW
Trop. Mid. Am. (except El Salvador), and n.w. S. Am.	
Arremonops rufivirgatus	OLIVE SPARROW
Texas, Mexico, s.w. Nicaragua and n.w. Costa Rica.	
Arremonops conirostris	GREEN-BACKED SPARROW
Trop. Mid. Am. (except El Salvador), and n. S. Am.	
Chlorura chlorura	GREEN-TAILED TOWHEE
W. U. S.; winters to cent. Mexico.	
Pipilo ocai	COLLARED TOWHEE
Mts. s. Mexico (Jalisco and Veracruz to Guerrero and Oaxaca).	
Pipilo macronyx³	OLIVE-BACKED TOWHEE
Mts. cent. Mexico (Michoacan and Hidalgo to Oaxaca).	
Pipilo maculatus³	SPOTTED TOWHEE
W. N. Am. and highlands (mts. farther south) Mexico to Guatema-la.	
Pipilo fuscus	BROWN TOWHEE
W. U. S. and Mexico (Baja Calif., Sonora and interior highlands).	
Pipilo albicollis^{3a}	WHITE-THROATED TOWHEE
Mts. s. Mexico (Guerrero, Puebla, Oaxaca).	
Pipilo aberti	ABERT'S TOWHEE
S.w. U. S. and n.w. Mexico (n. Baja Calif. and n.w. Sonora).	
Melozone kieneri	RUSTY-CROWNED GROUND-SPARROW
Highlands w. Mexico (Sonora and Chihuahua to Chiapas), Guate-mala, El Salvador, Honduras, and Costa Rica.	

² Considered by Hellmayr and others races of the S. Am. *A. torquatus*; if so, the entire complex may be called Striped Brush-Finch. These birds, as well as *brunnei-nucha*, were placed in a genus *Buarremom* by Ridgway.

³ The A.O.U. Check-List Committee has recently accepted Sibley's arguments for treating these forms as races of *P. erythrophthalmus* of eastern North America (Auk, 71: 312; Sibley, Univ. Calif. Publ. Zool., 56: 116-120). The A.O.U. Committee has adopted Rufous-sided Towhee as the English name for the entire complex.

^{3a} Often called *rutilus* (but cf. Stresemann, Condor, 1954: 91).

Melozone leucotis	WHITE-EARED GROUND-SPARROW
Highlands s. Mexico (Chiapas), Guatemala, El Salvador, Nicaragua and Costa Rica.	
Oriturus⁴ superciliosus	STRIPED SPARROW
Mexican tableland (Sonora and Chihuahua s. to Michoacan and Veracruz).	
Calamospiza melanocorys	LARK BUNTING
Cent. N. Am.; winters to Mexico (Baja Calif. and cent. Plateau).	
Passerculus sandwichensis	SAVANNAH SPARROW
N. Am. and Mexico (n.w. coast and interior highlands), Guatemala (high mts.); winters to Guatemala and El Salvador.	
Ammodramus savannarum	GRASSHOPPER SPARROW
N. Am. and locally through Mid. Am. to n.w. S. Am.; northern birds winter to El Salvador.	
Ammodramus bairdii	BAIRD'S SPARROW
Cent. N. Am.; winters to Mexico.	
Xenospiza baileyi	SIERRA MADRE SPARROW
Mts. s.w. Mexico (Durango, Jalisco, Distrito Federal).	
Pooecetes gramineus	VESPER SPARROW
N. Am.; winters to Mexico and Guatemala.	
Chondestes grammacus	LARK SPARROW
Cent. and w. N. Am. and highlands n. Mexico (Chihuahua, Coahuila and Durango); winters to Mexico, Guatemala and El Salvador.	
Aimophila quinquestriata	FIVE-STRIPED SPARROW
W. Mexico (Sonora and Chihuahua to Jalisco).	
Aimophila mystacalis	BRIDLED SPARROW
Mts. s. Mexico (Veracruz, Puebla, Oaxaca).	
Aimophila humeralis	BLACK-CHESTED SPARROW
Highlands s. Mexico (Colima to Morelos and Puebla).	
Aimophila ruficauda	STRIPE-HEADED SPARROW
Arid areas s.w. Mexico (Durango s.), Guatemala, El Salvador, Honduras, Nicaragua, and n.w. Costa Rica.	
Aimophila carpalis	RUFOUS-WINGED SPARROW
Arid areas Arizona and n.w. Mexico (Sonora and Sinaloa).	
Aimophila sumichrasti	CINNAMON-TAILED SPARROW
Arid. s. Mexico (Oaxaca).	
Aimophila notosticta	OAXACA SPARROW
S.e. Mexico (Oaxaca and possibly Puebla).	
Aimophila rufescens	RUSTY SPARROW
Arid areas trop. Mid. Am. s. to n.w. Costa Rica.	

⁴ *Plagiospiza superciliosa* is preferred by Hellmayr and Ridgway.

Aimophila ruficeps	RUFOUS-CROWNED SPARROW
S.w. U. S. and highlands Mexico (s. to Veracruz and Oaxaca).	
Aimophila botterii	BOTTERI'S SPARROW
S.w. U. S. and Mexico (highlands chiefly, lowlands in Tamaulipas), and w. Guatemala.	
Aimophila petenica⁵	YELLOW-CARPALLED SPARROW
S.e. Mexico, Guatemala, w. Nicaragua and n.w. Costa Rica.	
Aimophila cassinii	CASSIN'S SPARROW
S.w. U. S. to n. Mexico; winters in Mexico.	
Amphispiza bilineata	BLACK-THROATED SPARROW
Arid areas of s.w. U. S. and n. and w. Mexico (s. to Durango and Hidalgo).	
Amphispiza belli	SAGE SPARROW
Arid areas of w. U. S. and n.w. Mexico (Baja Calif.); winters to n.w. Mexico.	
Junco hyemalis⁶	SLATE-COLORED JUNCO
N. N. Am.; winters casually to n.w. Mexico.	
Junco oreganus⁷	OREGON JUNCO
W. N. Am. and in mts. to n.w. Mexico (n. Baja Calif.); winters to n.w. Mexico.	
Junco caniceps⁸	GRAY-HEADED JUNCO
Rocky Mts. of U. S.; winters to n.w. Mexico.	
Junco phaeonotus⁹	YELLOW-EYED JUNCO
Mts. s. Arizona through Mexico to Guatemala.	
Junco vulcani	VOLCANO JUNCO
Volcanic summits near or above timberline in Costa Rica and w. Panama.	
Spizella passerina	CHIPPING SPARROW
N. Am. s. through highlands Mid. Am. to n.e. Nicaragua.	
Spizella pallida	CLAY-COLORED SPARROW
Cent. N. Am.; winters to Mexico and casually Guatemala.	
Spizella breweri	BREWER'S SPARROW
Arid areas Great Basin and mts. w. N. Am.; winters to Mexico.	

* Hellmayr suggests this group may be races of Botteri's Sparrow, *A. botterii*.

* Many ornithologists would treat as races of *hyemalis*, the members of the *oreganus* and *caniceps* groups.

⁷ Includes the Guadalupe Junco, *insularis* (Guadalupe I.), regarded by A. Miller as a species.

⁸ Includes *dorsalis* (formerly regarded as a race of *phaeonotus*), which winters to Mexico.

⁹ Includes Baird's Junco, *bairdi* (Cape Dist., Baja Calif.), Chiapas Junco, *fulvescens* (Chiapas highlands), and Guatemala Junco, *alticola* (mts. of Guatemala and adjacent e. Chiapas), each regarded as a species by A. Miller.

Spizella pusilla	FIELD SPARROW
N. Am. e. of Rockies; winters to n.e. Mexico.	
Spizella wortheni ¹⁰	WORTHEN'S SPARROW
S.w. U. S. to n. Mexico (Tamaulipas); winters to s. Mexico.	
Spizella atrogularis	BLACK-CHINNED SPARROW
S.w. U. S. and highlands w. Mexico (to Guerrero and Puebla).	
Zonotrichia leucophrys	WHITE-CROWNED SPARROW
N. Am.; winters to highlands cent. Mexico.	
Zonotrichia atricapilla ¹	GOLDEN-CROWNED SPARROW
N.w. N. Am.; winters to n.w. Mexico (Baja Calif.).	
Zonotrichia albicollis	WHITE-THROATED SPARROW
N.e. N. Am.; winters to n.e. Mexico, casual Guadalupe I. off Baja Calif.	
Zonotrichia capensis ^{1*}	RUFOUS-COLLARED SPARROW
Highlands Mid. Am. (except Brit. Honduras), from Chiapas to cent. Panama, most of S. Am. (montane in trop. latitudes), Hispaniola, Curaçao, Aruba.	
Passerella iliaca	FOX SPARROW
N. N. Am. s. in mts. to California; winters to n.w. Mexico.	
Melospiza ² <i>lincolnii</i>	LINCOLN'S SPARROW
N. N. Am.; winters Mexico, Brit. Honduras, Guatemala, El Salvador, Honduras; one Panama spec.	
Melospiza georgiana	SWAMP SPARROW
E. N. Am.; winters to Mexico (Jalisco and Tamaulipas).	
Melospiza melodia	SONG SPARROW
N. Am. and Mexico (Baja Calif. and plateau region to Michoacan and Puebla).	
Emberizoides herbicola	WEDGE-TAILED GROUND-FINCH
S.w. Costa Rica, w. Panama, and S. Am. (chiefly e. of the Andes).	
Rhynchophanes mccownii	McCOWN'S LONGSPUR
Cent. N. Am.; winters to n. Mexico (Chihuahua and Durango).	
Calcarius ornatus	CHESTNUT-COLLARED LONGSPUR
Cent. N. Am.; winters to Mexico (Sonora, Chihuahua and Vera-cruz).	

¹⁰ May be conspecific with *S. pusilla*, the Field Sparrow.

¹ Formerly known as *Z. coronata*.

^{1*} Sometimes placed in a separate genus *Brachyspiza*.

² Some ornithologists merge this genus in *Passerella*.

BIBLIOGRAPHY

The following list is by no means complete. Only works of special usefulness for distribution or identification are included. A very few ecological papers are noted. Regional check-lists mentioning earlier literature and summarizing prior distributional information, are marked by an asterisk; those covering a whole country by a double asterisk. Few papers published prior to 1900 are included, because the locality data are usually repeated in the Ridgway or Hellmayr works listed below, or in the regional check-lists mentioned.

GENERAL

- AMERICAN ORNITHOLOGISTS' UNION COMMITTEE.** 1931. Check-list of North American birds. 4th. ed. This work and its Supplements, Nineteenth to Twenty-ninth inclusive (published between 1944 and 1954 in the *Auk*), involve the nomenclature and distribution of many Middle American species.
- BENT, A. C.** 1924-1953. Life histories of North American birds. U. S. Natl. Mus. Bull. The recent volumes contain data (often not elsewhere published) on occurrences in Middle America of North American species.
- CHAPMAN, F. M.** 1917. The Distribution of Bird-Life in Colombia. Bull. Amer. Mus. Nat. Hist., 36: 1-729. The first part of this work (and the next), describing distribution of forests, the life-zones, and the extension of the Subtropical Zone into Central America (pp. 70-158), is of the greatest interest to a student of Middle American bird distribution.
- CHAPMAN, F. M.** 1926. The Distribution of Bird-Life in Ecuador. Bull. Amer. Mus. Nat. Hist., 55: 1-784. See particularly pp. 23-133.
- GRISCOM, L.** 1945. Modern Bird Study. Harvard Univ. Press. A very able, and more up-to-date, discussion of the distribution and origin of birds in Middle America is contained in chapters VII and VIII.
- HARROWER, D. E.** 1936. The habits of the passerine birds of Central America, with particular reference to their breeding. Unpublished doctoral dissertation, Cornell Univ. Copy in Cornell Univ. Library. A very useful compilation.
- HELLMAYR, C. E.** 1924-1949. Catalogue of Birds of the Americas. Field Mus. Nat. Hist., Zool. Ser. 13, pts. 1-11. (Begun by C. B. Cory and completed by B. Conover.) Lists each form, giving distribution, synonymy, often discussing taxonomy and distinguishing characters of critical subspecies. The volumes written by Hellmayr are the chief modern authority on neotropical taxonomy.
- MURPHY, R. C.** 1936. Oceanic Birds of South America. Vols. 1-2. Amer. Mus. Nat. Hist. The information regarding the factors controlling sea-bird distribution is not limited to South America. There is also much detail as to Cocos I., Costa Rica, and the Pearl Is., Panama.
- PETERS, J. L.** 1931-1951. Check-List of Birds of the World. Vols. 1-7. Harvard Univ. Press. The volumes so far published cover Struthionidae to Rhinocryptidae. One of the chief modern authorities on technical nomenclature.
- RIDGWAY, R.** 1901-1950. Birds of North and Middle America. U. S. Natl. Mus. Bull. 50, pts. 1-11. (Continued by H. Friedmann, who prepared pts. 9-11.)

Though the nomenclature of the Ridgway parts is somewhat outmoded and the work is still incomplete (Tinamiformes to Anseriformes remain to be treated), this work is of basic importance and usefulness. It provides full descriptions of almost all Middle American forms, with drawings of generic characters, and also gives detailed distributional data, and synonyms.

- SALVIN, O. AND F. D. GODMAN. 1879-1904. *Biologia Centrali Americana: Aves*. Vols. 1-3. Despite the outmoded nomenclature, this expensive work, with many colored plates, simple species descriptions (in Latin), and often discussion in English of behavior, habitat and Middle American distribution, is still useful. The introduction in volume 1 (published last) gives in tabular form the distribution, as then understood, of each species by countries.
- SHARPE, R. B. AND OTHERS. 1888-1898. *Catalogue of Birds of the British Museum*. Vols. 1-28. For species not treated in Ridgway's work, brief descriptions usually can be found here.
- SHELFORD, V. E. AND OTHERS. 1926. *Naturalist's Guide to the Americas*. Williams & Wilkins Co. Section 3A, Mexico and Central America, gives concise descriptions of topography, climate and major ecological features.
- SKUTCH, A. F. 1954. *Life histories of Central American Birds*. Cooper Orn. Soc. Pac. Coast Avifauna, no. 31. Detailed accounts of 40 species of Coerebidae, Parulidae, Thraupidae, Icteridae and Fringillidae, with drawings of each, many photographs, and much comparative material.
- ZIMMER, J. T. 1931-1953. *Studies of Peruvian Birds*. Nos. 1-65. Amer. Mus. Novit. nos. 500 *et seq.* This series of taxonomic papers, though primarily relating to Peruvian birds, contains important discussions regarding taxonomy and nomenclature of many species found in Middle America, and often lists localities in that area not elsewhere published.

MEXICO

Entire country

- **BLAKE, E. R. 1953. *Birds of Mexico: A Guide for Field Identification*. Univ. of Chicago Press. A pocket guide, providing concise descriptions of 967 species, with many line drawings. It serves also as a Mexican check-list, for technical names and Mexican ranges of each subspecies (some 2000) are stated. Indispensable in Mexico, useful elsewhere in Middle America.
- DAVIS, L. I. *et al.* 1947-1954. Mexican breeding-bird censuses. *Aud. Field Notes*, 1-8 (no. 6). Valuable studies made each year in different localities and habitats.
- **EDWARDS, E. P. 1955. *Finding Birds in Mexico*. 101 pp. E. P. Edwards & Co., Amherst, Va. Where to find birds along the highways, with ecological and other useful data, including a list of almost all Mexican species.
- **FRIEDMANN, H., L. GRISCOM AND R. T. MOORE. 1950. *Distributional Check-List of the Birds of Mexico*. Pt. 1 (Tinamidae to Trochilidae). Pac. Coast Av. no. 29. Cooper Orn. Club. Gives localities where each form has been collected. When complete it will be the basic distributional work on Mexico.
- GOLDMAN, E. A. 1951. *Biological Investigations in Mexico*. Smith. Misc. Coll. vol. 115. Describes topography and vegetation of each Mexican state as found on expeditions between 1892-1906, with photographs. Discusses life zones and biotic provinces, listing characteristic plants, birds and mammals.
- GOLDMAN, E. A. AND R. T. MOORE. 1945. The biotic provinces of Mexico. *Journ. Mam.*, 26: 347-360. Describes the biotic provinces.

- GRISCOM, L.** 1950. Distribution and origin of the birds of Mexico. Bull. Mus. Comp. Zool. 103, no. 6: 341-382. Discusses the origin and analyzes the various elements of the Mexican avifauna.
- LEOPOLD, A. S.** 1950. The vegetation zones of Mexico. Ecology, 31(4): 507-518. A valuable ecological study, with a vegetation map.
- SURTON, G. M.** 1951. Mexican Birds. Univ. of Okla. Press. Beautiful plates, many drawings and an appendix summarizing the Mexican avifauna make this book much more than a charming ornithological travelogue of "First Impressions".

The following groupings of Mexican states are simply for bibliographic convenience and do not imply bio-geographic relationship (*cf.* Goldman and Moore, *supra*). In a few cases a paper listed under one grouping may contain important data relating to other states; the broader scope is indicated in the title of the paper or in the comment.

Northwest Pacific:¹ Baja California, Sonora, Sinaloa, Nayarit

- BANCROFT, G.** 1930. The breeding birds of central Lower California. Condor, 32: 20-49.
- ***BOND, J. AND R. M. DE SCHAUENSEE.** 1944. The birds of the Tres Marias Islands. In Results of the Fifth George Vanderbilt Expedition (1941). Ac. Nat. Sci. Phila. Mon. No. 6: 45-56.
- GRAYSON, A. J. AND G. N. LAWRENCE.** 1871. On the physical geography and natural history of the islands of the Tres Marias and of Socorro, off the western coast of Mexico. Proc. Boston Soc. Nat. Hist., 14: 261-302.
- ***GRINNELL, J. A.** 1928. A distributional summation of the ornithology of Lower California. Univ. Calif. Publ. Zool., 32: 1-300. The basic Baja California check-list.
- HILL, H. M. AND I. L. WIGGINS.** 1948. Ornithological notes from Lower California. Condor, 50: 155-161.
- ***HOWELL, T. R. AND T. J. CADE.** 1954. The birds of Guadalupe Island in 1953. Condor, 56: 283-294. A complete check-list.
- KENYON, K. W.** 1947. Notes on occurrences of birds in Lower California. Condor, 49: 210.
- MCELLAN, M. E.** 1926. Expedition to the Revillagigedo Islands, Mexico, in 1925. The birds and mammals. Proc. Cal. Ac. Sci., 15: 279-322.
- MCELLAN, M. E.** 1927. Notes on birds of Sinaloa and Nayarit, Mexico in the Fall of 1926. Proc. Cal. Ac. Sci., 16: 1-15.
- MILLER, W. DEW.** 1905. List of birds collected in southern Sinaloa, Mexico by J. H. Batty during 1903-1905. Bull. Amer. Mus. Nat. Hist., 21: 339-369.
- MOORE, R. T.** 1938. Unusual birds and extensions of range in Sonora, Sinaloa and Chihuahua. Condor, 46: 9-14.
- NEFF, J. A.** 1947. Notes on some birds of Sonora, Mexico. Condor, 49: 32-34.
- PHILLIPS, A. R. AND D. AMADON.** 1952. Some birds of northwestern Sonora, Mexico. Condor, 54: 163-168.
- VAN ROSSEM, A. J.** 1932. The avifauna of Tiburon Island, Sonora, Mexico. Trans. San Diego Soc. Nat. Hist., 7(12): 119-150.

¹ For birds of Clipperton Island (a French possession) see Gifford (1913, 1919) and Snodgrass and Heller (1902) under "Costa Rica" below.

*VAN ROSSEM, A. J. 1945. A distributional survey of the birds of Sonora, Mexico. Occ. Pap. Mus. Zool. La. State Univ., 21: 1-379. The basic Sonora check-list, with full bibliography.

VAURIE, C. 1953. Summer records and observations on the Island of Tiburon, Sonora, Mexico. Condor, 55: 217-218.

West central: Chihuahua, Durango, Zacatecas, Aguascalientes

ALLEN, J. A. 1893. List of mammals and birds collected in northeastern Sonora and northwestern Chihuahua, Mexico on the Lumholtz archaeological expedition, 1890-1892. Bull. Amer. Mus. Nat. Hist., 5: 27-42.

BAILEY, A. M. AND H. B. CONOVER. 1935. Notes from the state of Durango, Mexico. Auk, 52: 421-424.

JOUY, P. L. 1894. Notes on birds of central Mexico, with descriptions of forms believed to be new. Proc. U. S. Nat. Mus., 16: 771-791.

KNOBLICH, I. W. 1950. Una lista de pajaros reconocidos en el Estado de Chihuahua. Ann. Inst. Biol. Mexico, 21: 155-157.

MILLER, W. DEW. 1906. List of birds collected in northwestern Durango, Mexico, by J. H. Batty, during 1903. Bull. Amer. Mus. Nat. Hist., 22: 161-183.

STAGER, K. E. 1954. Birds of the Barranca de Cobre Region of southwestern Chihuahua, Mexico. Condor, 56: 21-34.

VAN ROSSEM, A. J. 1934. Critical notes on Middle American birds. Bull. Mus. Comp. Zool., 77: 424-490. Part C of this paper, "A systematic report on the Brewster collection of Mexican birds" deals largely with Chihuahua and adjacent Sonora.

WEBSTER, J. D. AND R. T. ORR. 1952. Notes on Mexican birds from the states of Durango and Zacatecas. Condor, 54: 309-313.

WEBSTER, J. D. AND R. T. ORR. 1954. Summering birds of Zacatecas, Mexico, with a description of a new race of Worthen Sparrow. Condor, 56: 155-160.

WEBSTER, J. D. AND R. T. ORR. 1954. Miscellaneous notes on Mexican birds. Wilson Bull., 66: 267-269. Chiefly Durango and Guerrero, but also Revilla Gigedo Is., Sinaloa, San Luis Potosi and Jalisco.

South Pacific: Jalisco, Colima, Michoacan, Guerrero, Oaxaca, Chiapas

ALVAREZ DEL TORO, M. 1952. Contribucion al conocimiento de la oología y nidología de las aves chiapanecas. Ateneo, 4: 11-21.

ALVAREZ DEL TORO, M. 1952. New records of birds from Chiapas, Mexico. Condor, 54: 112-114.

ALVAREZ DEL TORO, M. 1952. Los Animales Silvestres de Chiapas. Tuxtla Gutierrez, Chiapas, Mexico. Though a popular work, the chapter "Aves" (pp. 91-180) contains original information as to habitat and behavior of many birds.

ALVAREZ DEL TORO, M. 1954. Notes on the occurrence of birds in Chiapas, Mexico. Condor, 56: 365.

BANGS, O. AND J. L. PETERS. 1928. A collection of birds from Oaxaca. Bull. Mus. Comp. Zool., 68: 305-404.

BERLIOZ, J. 1939. Etude d'une collection d'oiseaux du Chiapas (Mexique). Bull. Mus. Hist. Nat. Paris, ser. 2, 11: 360-377. Relates to Comitan district.

BLAKE, E. R. 1950. Report on a collection of birds from Guerrero, Mexico. Fieldiana Zool. 31, no. 39: 375-393.

BLAKE, E. R. 1950. Report on a collection of birds from Oaxaca, Mexico. Fieldiana Zool. 31, no. 40: 395-419.

BLAKE, E. R. AND H. C. HANSON. 1942. Notes on a collection of birds from Michoacan, Mexico. Field Mus. Nat. Hist. Zool. Ser., 22, no. 9: 513-551.

- BRODKORB, P. 1938. New birds from the district of Soconusco, Chiapas. Occ. Pap. Mus. Zool. Univ. Mich. no. 369: 1-7.
- BRODKORB, P. 1939. Rediscovery of *Heleodytes chiapensis* and *Tangara cabanisi*. Auk, 56: 447-450.
- BRODKORB, P. 1940. New birds from southern Mexico. Auk, 57: 542-549.
- BRODKORB, P. 1943. Notes on two rare birds in Chiapas, Mexico. Auk, 60: 281-281.
- DAVIS, J. 1953. Birds of the Tzitzio Region, Michoacan, Mexico. Condor, 55: 90-98.
- DAVIS, W. B. 1944. Notes on summer birds of Guerrero. Condor, 46: 9-14.
- EDWARDS, E. P. AND R. B. LEA. 1955. Birds of the Montserrat Area, Chiapas, Mexico. Condor, 57: 31-54.
- HOFFMEISTER, D. F. 1951. A western record of the Quetzal, *Pharomachrus mocinno*, and Chachalaca, *Penelopina nigra*, in Mexico. Auk, 68: 507-508.
- *GRISCOM, L. 1934. The ornithology of Guerrero, Mexico. Bull. Mus. Comp. Zool., 75: 367-422. The basic Guerrero check-list.
- GRISCOM, L. 1937. A collection of birds from Omitlame, Guerrero. Auk, 54: 192-199.
- LEA, F. B. AND E. P. EDWARDS. 1952. Notes on birds of the Lake Patzcuaro Region, Michoacan, Mexico. Condor, 52: 260-271.
- MARTIN DEL CAMPO, R. 1948. Contribucion para el conocimiento de la fauna ornitologica del estado de Guerrero. Ann. Inst. Biol. Mex., 19(1): 241-266. Lists Guerrero specimens, with localities, in the Instituto Biologico, adding a few to the Guerrero avifauna.
- SUTTON, G. M. 1953. New birds for the state of Michoacan, Mexico. Wilson Bull., 64: 221-223.
- TASHIAN, R. E. 1952. Birds from the Palenque region of northeastern Chiapas, Mexico, Auk, 69: 60-66.
- ZIMMERMAN, D. A. AND G. B. HARRY. 1951. Summer birds of Autlan, Jalisco. Wilson Bull., 63: 302-314.

Northeast: Coahuila, Nuevo Leon, Tamaulipas, San Luis Potosi.

- AMADON, D. AND A. R. PHILLIPS. 1947. Notes on Mexican birds, Auk, 64: 576-581. Chiefly Coahuila; some Durango and Hidalgo data.
- BURLEIGH, T. D. AND G. H. LOWERY, JR. 1942. Notes on the birds of southeastern Coahuila. Occ. Pap. Mus. Zool. La. State Univ., 12: 185-212.
- DAVIS, L. I. 1952. Winter bird census at Xilitla, San Luis Potosi, Mexico. Condor, 54: 345-355.
- EATON, S. W. AND E. P. EDWARDS. 1948. Notes on the birds of the Gomez Farias region of Tamaulipas. Wilson Bull., 60: 109-114.
- EVENDEN, F. G., JR. 1952. Notes on Mexican bird distribution. Wilson Bull., 64: 112-113. Chiefly observations from the northeast, but also some in Zacatecas, Durango and Hidalgo.
- GRABER, R. R. AND J. W. GRABER. 1954. Yellow-headed Vulture in Tamaulipas, Mexico. Condor, 56: 165-166.
- PHILLIPS, J. C. 1911. A year's collecting in the state of Tamaulipas, Mexico. Auk, 28: 67-89.
- MARTIN, P. S., C. R. ROBINS AND W. B. HEED. 1954. Birds and biogeography of the Sierra de Tamaulipas, an isolated pine-oak habitat. Wilson Bull., 66: 38-57.
- ROBINS, C. R., P. S. MARTIN AND W. B. HEED. 1951. Frigatebird, Oystercatcher, Upland Plover, and various terns on the coast of Tamaulipas, Mexico. Wilson Bull., 63: 336.

- ROBINS, C. R. AND W. B. HEED. 1951. Bird notes from La Joya de Salas, Tamaulipas. *Wilson Bull.*, 63: 263-270.
- SUTTON, G. M. AND T. D. BURLEIGH. 1939. A list of birds observed on the 1938 Semple Expedition to northeastern Mexico. *Occ. Pap. Mus. Zool. La. State Univ.*, 3: 15-46.
- SUTTON, G. M. AND T. D. BURLEIGH. 1940. Birds of Valles, San Luis Potosi, Mexico. *Condor*, 42: 259-262.
- SUTTON, G. M. AND T. D. BURLEIGH. 1940. Birds of Tamazunchale, San Luis Potosi. *Wilson Bull.*, 52: 221-233.
- SUTTON, G. M., O. S. PETTINGILL, JR., AND R. B. LEA. 1942. Notes on the birds of the Monterrey District of Nuevo Leon, Mexico. *Wilson Bull.*, 54: 199-203.
- SUTTON, G. M. AND O. S. PETTINGILL, JR. 1943. Birds of Linares and Galeana, Nuevo Leon, Mexico. *Occ. Pap. Mus. Zool. La. State Univ.*, 16: 273-290.
- SUTTON, G. M. AND O. S. PETTINGILL, JR. 1944. Birds of the Gomez Farias region, Tamaulipas. *Auk*, 59: 1-34.
- SUTTON, G. M., R. B. LEA AND E. P. EDWARDS. 1950. Notes on the ranges and breeding habits of certain Mexican birds. *Bird-Banding*, 21: 45-59.

South central: Guanajuato, Queretaro, Hidalgo, Mexico, Distrito Federal, Morelos, Puebla.

- MOORE, R. T. 1945. The Transverse Volcanic Biotic Province of Central Mexico and its relationship to adjacent provinces. *Trans. San Diego Soc. Nat. Hist.*, 10: 217-236. Covers more than the states listed above.
- NEWMAN, R. J. 1954. *Toxostoma ocellatum* and *Diglossa baritula* in Hidalgo. *Condor*, 56: 361.
- PAYNTER, R. A., JR. 1952. Birds from Popocatepetl and Ixtaccihuatl, Mexico. *Auk*, 69: 293-301.
- PITELKA, F. A. 1948. Notes on the distribution and taxonomy of Mexican game birds. *Condor*, 50: 113-128. Contains many records from south central area, also Sonora, Guerrero, Chiapas and elsewhere.
- SUTTON, G. M. AND T. D. BURLEIGH. 1941. Birds recorded in the state of Hidalgo, Mexico, by the Semple Expedition of 1939. *Ann. Carn. Mus.*, 28: 169-186.
- SUTTON, G. M. AND T. D. BURLEIGH. 1942. Birds recorded in the Federal District and states of Puebla and Mexico by the 1939 Semple Expedition. *Auk*, 59: 418-423.

So. Gulf and Caribbean: Veracruz, Tabasco, Campeche, Yucatan, Quintana Roo.

- AMADON, D. AND D. R. ECKELBERRY. 1955. Observations on Mexican birds. *Condor*, 57: 65-80. Ill. Chiefly notes on Veracruz, Chiapas, Mexico, Oaxaca, Tamaulipas, Puebla.
- BANGS, O. AND J. L. PETERS. 1927. Birds of the rain forest region of Vera Cruz. *Bull. Mus. Comp. Zool.*, 67: 471-487.
- *BRODKORB, P. 1943. Birds of the Gulf Lowlands of southern Mexico. *Misc. Publ. Mus. Zool. Univ. Mich.*, 55: 1-88. A basic check-list of Tabasco, with notes on adjacent parts of Veracruz, Campeche and Chiapas.
- BRODKORB, P. 1948. Some birds of the lowlands of central Veracruz, Mexico. *Quart. Jour. Florida Acad. Sci.*, 10: 31-38.
- CHAPMAN, F. M. 1896. Notes on birds observed in Yucatan. *Bull. Amer. Mus. Nat. Hist.*, 8: 271-290.
- CHAPMAN, F. M. 1898. Notes on birds observed at Jalapa and Las Vigas, Vera Cruz, Mexico. *Bull. Amer. Mus. Nat. Hist.*, 10: 15-43.

- COLE, L. J. 1906. Aves from Yucatan. Bull. Mus. Comp. Zool. 50: 109-146.
- DAVIS, W. B. 1945. Notes on Veracuzan birds. Auk, 62: 272-286.
- GRISCOM, L. 1926. The ornithological results of the Mason-Spinden expedition to Yucatan. Pt. 1. Introduction. Birds of the mainland of eastern Yucatan. Am. Mus. Nat. Hist. Nov. No. 235: 1-19.
- GRISCOM, L. 1926. The ornithological results of the Mason-Spinden expedition to Yucatan. Pt. 2. Chinchorro Bank and Cozumel Island. Am. Mus. Nat. Hist., Nov. No. 236: 1-13.
- *LOETSCHER, F. W., JR. 1941. Ornithology of the Mexican state of Veracruz, with an annotated list of the birds. Unpublished doctoral dissertation, Cornell Univ. Copy in Cornell Univ. Library. An annotated check-list for Veracruz.
- LOETSCHER, F. W., JR. 1955. North American migrants in the state of Veracruz, Mexico: a summary. Auk, 72: 14-54.
- LOWERY, G. H., JR. AND W. W. DALQUEST. 1951. Birds from the state of Veracruz, Mexico. Univ. Kansas Publ., Mus. Nat. Hist., 3: 531-649. An important paper on the area.
- LOWERY, G. H., JR. AND R. J. NEWMAN. 1949. New birds from the state of San Luis Potosi and the Tuxtla Mountains of Veracruz, Mexico. Occ. Pap. Mus. Zool. La. State Univ., 22: 1-10.
- PAYNTER, R. A. JR. 1950. Rare migration and wintering records from the Yucatan Peninsula, Postilla, 2: 1-7.
- PAYNTER, R. A., JR. 1953. Migrants on the Campeche Bank. Auk, 70: 338-349.
- PAYNTER, R. A., JR. 1954. Two species new to the Mexican avifauna. Auk, 71: 204.
- *PAYNTER, R. A., JR. 1955. The Ornithogeography of the Yucatan Peninsula. Peabody Mus. Bull., 9: 1-347. Yale Univ.
- PETERS, J. L. 1913. List of birds collected in the territory of Quintana Roo, Mexico, in the winter and spring of 1912. Auk, 30: 367-380. Lists 132 species.
- SUTTON, G. M. AND T. D. BURLEIGH. 1940. Birds of Las Vigas, Veracruz. Auk, 57: 234-243.
- TRAYLOR, M. A., JR. 1941. Birds from the Yucatan Peninsula. Field Mus. Nat. Hist. Zool. Ser. 24, no. 19: 195-225. Lists 217 species from Yucatan and Campeche.
- TRAYLOR, M. A., JR. 1949. Notes on some Veracruz birds. Fieldiana Zool., 31: 269-275.
- VAN TYNE, J. AND TRAUTMAN, M. B. 1941. New birds from Yucatan. Univ. Mich. Occ. Pap. Mus. Zool. no. 439: 1-11.
- WARNER, D. W. AND R. M. MENGEI. 1951. Notes on birds of the Veracruz coastal plain. Wilson Bull., 63: 288-295.
- WETMORE, A. 1943. The birds of southern Veracruz, Mexico, U. S. Nat. Mus. Bull., 93: 215-340. An important paper relating to the Tuxtla area.

BRITISH HONDURAS

- There is no check-list, and the literature specifically devoted to this colony is scanty. The basis for a list to 1904 may be compiled from the tables in volume 1 of Salvin and Godman, *Biologia Centrali Americana, Aves*.
- AUSTIN, O. L., JR. 1929. Birds of the Cayo District, British Honduras. Bull. Mus. Comp. Zool., 69: 363-394. A collection of 157 species, many not theretofore reported; gives bibliography.
- BANGS, O. AND M. E. PECK. 1908. On some rare and new birds from British Honduras. Proc. Biol. Soc. Wash., 21: 43-46.

- *BOND, J. 1954. Birds of Turneffe and Northern Two Cays, British Honduras. *Not. Nat.* no. 260: 1-10.
- LANTZ, D. E. 1899. A list of birds collected by Col. N. S. Goss in Mexico and Central America. *Trans. Kansas Acad. Sci.*, 16: 218-224. A number of the birds listed have British Honduras localities.
- PECK, M. E. 1910. The effect of natural enemies on the nesting habits of some British Honduras birds. *Condor*, 12: 53-60.
- SALVIN, O. 1864. A fortnight amongst the sea-birds of British Honduras. *Ibis*, 1864: 372-387.

GUATEMALA

- CARRIKER, M. A., JR. AND R. M. DE SCHAUENSEE. 1935. An annotated list of two collections of Guatemalan birds in the Academy of Natural Sciences of Philadelphia. *Proc. Ac. Nat. Sci. Phila.*, 87: 411-455.
- **GRISCOM, L. 1932. The distribution of bird-life in Guatemala. *Bull. Amer. Mus. Nat. Hist.*, 64: 1-439. The Guatemalan check-list, with localities, bibliography, discussion of faunistics and taxonomic notes. Lists 736 forms.
- GRISCOM, L. 1935. The birds of Sierra de las Minas, Eastern Guatemala. *Ibis*, 1935: 801-837.
- SAUNDERS, G. B. 1950. The game birds and shorebirds of Guatemala. In *A Fish and Wildlife Survey of Guatemala*. U. S. Dept. Int. Fish and Wildlife Serv. Spec. Sci. Rept. no. 5: 3-98. Gives descriptions and information as to status and behavior of birds treated, banding recoveries in Guatemala, and good bibliography.
- STEYERMARK, J. A. 1950. Flora of Guatemala. *Ecology* 31(3): 368-372. An ecological study, with good map showing the various vegetation regions.
- TASHIAN, R. E. 1953. The birds of southeastern Guatemala. *Condor*, 55: 198-210.
- VAN ROSSEM, A. J. 1934. Critical notes on Middle American birds. *Bull. Mus. Comp. Zool.*, 77: 387-405. Part A of this paper "Notes on some species and subspecies of Guatemala birds" contains a few additions to the avifauna.
- VAN TYNE, J. 1935. The birds of northern Peten, Guatemala. *Univ. Mich. Mus. Zool. Misc. Pub. no. 27*: 1-47. Adds 21 forms to Guatemala.
- WETMORE, A. 1941. Notes on the birds of the Guatemalan highlands. *Proc. U. S. Nat. Mus.* 89: 523-581.

EL SALVADOR

- COOKE, M. T. 1941. Banded birds recovered in El Salvador. *Auk*, 58: 589-590.
- **DICKEY, D. R. AND A. J. VAN ROSSEM. 1938. The birds of El Salvador. *Field Mus. Nat. Hist., Zool. Ser.* 23, *Publ. 406*: 1-609. A masterly annotated check-list, with data on distribution, ecology, moults, taxonomy, behavior, and bibliography. Lists 446 forms.
- MARSHALL, J. T., JR. 1943. Additional information concerning the birds of El Salvador. *Condor*, 45: 21-33. Adds several species.

HONDURAS

- BOND, JAMES. 1936. Resident birds of the Bay Islands of Spanish Honduras. *Proc. Ac. Nat. Sci. Phila.*, 88: 353-364.
- CARR, A. F., JR. 1950. Outline for a classification of animal habitats in Honduras. *Bull. Am. Mus. Nat. Hist.* 94(10): 563-594. An ecological study.

- CARR, M. J. AND J. C. DICKINSON, JR. 1951. The San Geronimo Swift in Honduras, *Wilson Bull.*, 63: 271-273. Refers also to several other species.
- DEIGNAN, H. G. 1936. Notes on a small collection of birds from the Republic of Honduras. *Auk*, 53: 186-193.
- GRISCOM, L. 1932. New Birds from Honduras and Nicaragua. *Proc. New Eng. Zool. Club*, 13: 55-62.
- **STONE, W. 1932. The birds of Honduras. *Proc. Ac. Nat. Sci. Phila.*, 84: 291-342. A distributional check-list, with bibliography. Lists 420 forms.

NICARAGUA

There is no published check-list for Nicaragua, though there is much data scattered through the literature. A basic check-list to 1904 can be compiled from the tables in *Biologia Centrali Americana*.

- GRISCOM, L. 1932. (See under Honduras.)
- HUBER, W. 1932. Birds collected in northeastern Nicaragua in 1922. *Proc. Ac. Nat. Sci. Phila.*, 84: 205-249.
- MILLER, W. DEW. AND L. GRISCOM. 1921. Descriptions of proposed new birds from Central America, with notes on other little known forms. *Am. Mus. Nat. Hist. Nov. No. 25*: 1-13.
- MILLER, W. DEW. AND L. GRISCOM. 1925. Description of new birds from Nicaragua. *Am. Mus. Nat. Hist. Nov. No. 159*: 1-9.
- MILLER, W. DEW. AND L. GRISCOM. 1925. Notes on Central American birds with descriptions of new forms. *Am. Mus. Nat. Hist. Nov. No. 183*: 1-14.
- MILLER, W. DEW. AND L. GRISCOM. 1925. Further notes on Central American birds, with descriptions of new forms. *Am. Mus. Nat. Hist. Nov. No. 184*: 1-16.
- RENDAHL, H. 1919. Notes on a collection of birds from Panama, Costa Rica and Nicaragua. *Arkiv. for Zool.*, 12(8): 1-36. Lists 139 forms from Nicaragua, 73 from Costa Rica, 28 from Panama, collected by C. Bovallius in 1882-1883.
- RICHMOND, C. W. 1893. Notes on a collection of birds from eastern Nicaragua and the Rio Frio, Costa Rica. *Proc. U. S. Nat. Mus.* 16: 479-530.

COSTA RICA

- **CARRIKER, M. A., JR. 1910. An annotated list of the birds of Costa Rica, including Cocos Island. *Ann. Carn. Mus.*, 6, nos. 2-4: 314-915. An excellent check-list, with localities, information about habitat and behavior, and bibliography of Costa Rican ornithology. Lists 753 forms.
- *CHERRIE, G. K. 1891-1892. A preliminary list of the birds of San José, Costa Rica. *Auk*, 8: 271-279. *Auk*, 9: 21-27, 247-251, 322-329. Reports 162 species collected within a radius of two miles of the city, with notes on status and habits.
- FERRY, J. F. 1910. Catalogue of a collection of birds from Costa Rica. *Field Mus. Nat. Hist. Publ., Orn. Ser.*, 1: 257-282.
- GIFFORD, E. W. 1913. The birds of the Galapagos Islands, with observations of the birds of Cocos and Clipperton Islands (Columbiformes to Pelecaniformes). *Proc. Calif. Acad. Sci. Fourth ser.* 2, pt. 1: 1-132.
- GIFFORD, E. W. 1919. Field notes on the land birds of the Galapagos Islands and of Cocos Island, Costa Rica. *Proc. Calif. Acad. Sci. Fourth ser.* 2, pt. 2: 189-258.
- GRISCOM, L. 1933. Notes on the Havemeyer Collection of Central American birds. *Auk*, 50: 297-308. Costa Rican birds on pp. 297-300.

- RENDAHL, H. 1919. (See under Nicaragua.)
- SASSI, M. A. 1938-1939. Die Vogel der oesterreichischen Costa-Rica-Expedition. Erster Teil. *Temminckia*, 3: 279-232. Zweiter Teil. *Temminckia*, 4: 135-222.
- SMITH, A. 1931. New records of several species of birds rare or local in Costa Rica. *Condor*, 33: 249-300.
- SMITH, A. 1932. Some records for Costa Rica. *Auk*, 49: 496-497.
- *SNODGRASS, R. E. AND E. HELLER. 1902. The birds of Clipperton and Cocos Islands. Papers from the Hopkins-Stanford Galapagos Expedition, 1898-1899. Proc. Wash. Acad. Sci., 4: 501-520. Lists species known from these islands up to that time. This paper was apparently overlooked by Carriker.
- SWARTH, H. S. 1931. The avifauna of the Galapagos Islands. Occ. Pap. Calif. Acad. Sci., 18: 1-299. Includes some records from Cocos I., Costa Rica.
- WETMORE, A. 1944. A collection of birds from northern Guanacaste, Costa Rica. Proc. U. S. Nat. Mus. 95: 25-80.

PANAMA

- ALDRICH, J. W. AND B. P. BOLE. 1937. Birds and mammals of the western slope of the Azuero Peninsula, Republic of Panama. Cleveland Mus. Nat. Hist. Sci. Pub. 7: 1-196. Bird part written by Aldrich.
- ALDRICH, J. W. 1945. Notes on Panamanian birds. *Auk*, 62: 111-114. Mentions two species not listed by Griscom.
- ARBIB, R. S. AND F. LOETSCHER, JR. 1935. Some notes from the Panama Canal Zone—Summer 1934. *Auk*, 52: 326. Adds sight reports of a few shore birds not previously recorded from Panama.
- BANGS, O. 1901. On a collection of birds made by W. W. Brown, Jr. at David and Divala, Chiriqui. *Auk*, 23: 356-370.
- BANGS, O. 1902. On a second collection of birds made in Chiriqui by W. W. Brown, Jr. Proc. New England Zool. Club, 3: 15-70.
- BANGS, O. AND BARBOUR, T. 1922. Birds from Darien. Bull. Mus. Comp. Zool. 65, no. 6: 191-229.
- BOND, J. AND R. M. DE SCHAUENSEE. 1944. Results of the Fifth George Vanderbilt Expedition (1951). Ac. Nat. Sci. Phila. Mon., 6: 22-45. A collection of birds from Cape Garachine and Mount Sapo, Darien.
- DAVIDSON, M. E. M. 1938. On some birds rare in, or hitherto unrecorded from, Chiriqui Province, Panama. Proc. Cal. Ac. Sci. 17: 255-261.
- EISENMANN, E. 1948. Surf-bird and Vermilion Flycatcher in Panama. *Auk*, 65: 605-606.
- EISENMANN, E. 1950. Some notes on Panama birds collected by J. H. Batty. *Auk*, 67: 363-366. Highland species reported collected on low islands off Pacific coast.
- EISENMANN, E. 1951. Northern birds summering in Panama. *Wilson Bull.*, 63: 181-185.
- *EISENMANN, E. 1952. Annotated list of birds of Barro Colorado Island, Panama Canal Zone. Smith. Misc. Coll. 117, no. 5: 1-62. Breeding and migration dates, full bibliography; adds to the Canal Zone and Panama avifauna. 306 species listed in 5.6 sq. miles.
- GRISCOM, L. 1924. Descriptions of new birds from Panama and Costa Rica. Amer. Mus. Novit., no. 141. Chiefly from mts. of eastern Chiriqui and coast of Veraguas.

- GRISCOM, L. 1927. Undescribed or little known birds from Panama. Amer. Mus. Novit. no. 280. Coll. in Veraguas, Los Santos and Coclé.
- GRISCOM, L. 1927. An ornithological reconnaissance in eastern Panama in 1927. Amer. Mus. Novit., no. 282.
- GRISCOM, L. 1929. A collection of birds from Darien. Bull. Mus. Comp. Zool., 69, no. 8: 149-190.
- GRISCOM, L. 1932. The ornithology of the Caribbean coast of extreme eastern Panama. Bull. Mus. Comp. Zool., 72, no. 9: 303-372.
- **GRISCOM, L. 1935. The ornithology of the Republic of Panama. Bull. Mus. Comp. Zool. 78, no. 3: 261-382. The basic Panama check-list, with bibliography, and discussion of geography and faunal relations. Over 800 species, 1038 forms listed.
- IMHOFF, T. A. 1950. Additional bird records for Panama. Auk, 67: 265-267.
- MURPHY, R. C. 1945. Notes on Pearl Islands ornithology. Auk, 62: 114-117.
- *PETERS, J. L. 1931. Additional notes on the birds of the Almirante Bay region of Panama. Bull. Mus. Comp. Zool., 71(5): 310. Lists all forms then known from Bocas del Toro province, with references to earlier literature.
- *RENDAHL, H. 1920. A list of birds of the Pearl Islands, Bay of Panama. Arkiv for Zool., 13, no. 4: 1-56. Lists all forms then known from Pearl Islands, a total of 100.
- SCHOLES, R. T. AND K. T. SCHOLES. 1953. Notes from Panama and the Canal Zone. Condor, 56: 166-167.
- *STONE, W. 1918. Birds of the Panama Canal Zone, with special reference to a collection made by Mr. Lindsay L. Jewel. Proc. Ac. Nat. Sci. Phila., 70: 239-280. Lists all species then recorded, contains Canal Zone bibliography, some breeding data.
- *STURGIS, B. B. 1928. Field Book of Birds of the Panama Canal Zone. G. P. Putnam's Sons. A very useful pocket guide to identification, though the nomenclature is somewhat outmoded and not all Canal Zone species are described.
- *WETMORE, A. 1946. The birds of San José and Pedro Gonzales Islands, Republic of Panama. Smith. Misc. Coll. 106, no. 1: 1-60. A list of the birds known from these Pearl Islands; several of the pelagics are new to the Panama list.
- WETMORE, A. 1951. Four additional species for Panama. Auk, 68: 525-526.
- WETMORE, A. 1951. Additional forms of birds from Colombia and Panama. Smith. Misc. Coll., 117, no. 2: 1-11.
- *WETMORE, A. 1952. The birds of the Islands of Taboga, Taboguilla, and Urava, Panama. Smith. Misc. Coll., 121, no. 2: 1-32. A list of all forms reported from these small islands.
- WETMORE, A. 1953. Further additions to the birds of Panama and Colombia. Smith. Misc. Coll., 122, no. 8: 1-12.

NEIGHBORING COUNTRIES

Colombia

DE SCHAUENSEE, R. M. 1948-1952. The birds of the Republic of Colombia. Caldasia, 5, nos. 22-26: 251-1212. A check-list of all forms, giving range, Colombian localities, and keys to the identification of specimens. Useful for eastern Panama.

United States of America

- AMERICAN ORNITHOLOGISTS' UNION COMMITTEE. 1931. Check-list of North American birds. 4th ed. (See *supra* under "General").
- PETERSON, R. T. 1941. A Field Guide to Western Birds. Houghton Mifflin Co., Boston. Very useful in Mexico and, as regards North American migrants, elsewhere in Middle America.
- PETERSON, R. T. 1947. A Field Guide to the Birds. Houghton Mifflin Co., Boston. Second Rev. ed. The indispensable guide to field identification of North American species east of the Rockies. Useful in Middle America for eastern migrants.
- POUGH, R. H. 1946-1951. Audubon Bird Guide. Eastern Land Birds. Water Birds. Doubleday & Co., Inc., Garden City. A set of two books describing the appearance, habits and range of all species found in eastern and central North America, including southern Texas. Each species illustrated in color by Don Eckelberry.

Venezuela

- PHELPS, W. H. AND W. H. PHELPS, JR. 1950. Lista de las Aves de Venezuela. Bol. Soc. Venez. Cien. Nat., 12, no. 75: 1-427. Pt. 2 Passeriformes. A careful check-list of all forms of the order, giving range, Venezuelan localities, and an indication of habitat.

West Indies

- BOND, J. 1947. A Field Guide to Birds of the West Indies. Macmillan Co., New York. An excellent guide, with most resident species depicted in line drawings.
- BOND, J. 1950. Check-list of birds of the West Indies. Third ed. Acad. Nat. Sci. Phila. Gives West Indian locality data for each form, migration dates, and general range information. A First and Second Supplement published in 1951 and 1952.

INDEX TO GENERA AND GROUP NAMES

Where a group name appears on successive pages only the first page is here listed. A page reference in *italic* type indicates the name appears in a footnote.

- | | | |
|---|---|--|
| <p>Abeillia, 47
Acanthidops, 104
Accipiter, 21
Accipitridae, 20
Acestura, 51
Acrochordopus, 72
Actitis, 30
Aechmolophus, 69
Aechmophorus, 10
Aegolius, 43
Aeronautes, 45
Agamia, 15
Agelaius, 96
Agriocaris, 26
Aimophila, 106
Aix, 19
Ajaia, 16
Alaudidae, 73
Albatross, 10
Alcedinidae, 53
Alcidae, 35
Amaurolimnas, 27
Amaurospiza, 103
Amaurospizopsis, 103
Amazilia, 48
Amazona, 39
Amblycercus, 94
Ammodramus, 106
Amphispiza, 107
Anabacerthia, 60
Anas, 18
Anatidae, 17
Androdon, 45
Anhinga, 14
Anhingidae, 14
Ani, 40
Anous, 35
Anser, 17
Antbird, 61</p> | <p>Anthracothorax, 46
Anthus, 84
Antpitta, 63
Antshrike, 61
Ant-Tanager, 100
Antthrush, 62
Antvireo, 61
Antwren, 61
Aphanotriccus, 69
Aphelocoma, 75
Aphriza, 30
Apodidae, 44
Aquila, 23
Ara, 38
Araçari, 55
Aramidae, 26
Aramides, 27
Aramus, 26
Aratinga, 38
Archilochus, 51
Ardea, 14
Ardeidae, 14
Ardeola, 15
Arenaria, 30
Arremon, 105
Arremonops, 105
Asio, 42
Aspatha, 53
Asyndesmus, 56
Atalotriccus, 71
Ateleodacnis, 88
Atlapetes, 104
Atthis, 51
Atticora, 73
Attila, 64
Auklet, 35
Aulacorhynchus, 55
Auriparus, 76
Automolus, 60</p> | <p>Avocet, 31
Aythya, 19
Azurecrown, 48
Baldpate, 19
Bananaquit, 88
Bananivorus, 95
Bangsia, 99
Barbet, 54
Barbtail, 59
Barbthroat, 45
Bartramia, 29
Baryphthengus, 53
Basileuterus, 93
Becard, 65
Bellbird, 65
Belonopterus, 28
Bentbill, 70
Bittern, 15
Blackbird, 95, 96
Bluebird, 83
Bobolink, 97
Bobwhite, 25
Bolborhynchus, 39
Bombycilla, 84
Bombycillidae, 84
Booby, 13
Botaurus, 16
Brachygalba, 53
Brachyramphus, 35
Brachyspiza, 108
Brant, 17
Branta, 17
Brilliant, 50
Brotogeris, 39
Brush-Finch, 104
Buarremont, 104
Bubo, 41
Bubulcus, 15</p> |
|---|---|--|

- Bucconidae, 53
Bucephala, 19
Bufflehead, 19
Bunting, 102
Burhinidae, 32
Burhinus, 32
Busarellus, 22
Bush-Tanager, 101
Bushtit, 76
Buteo, 21
Buteogallus, 22
Butorides, 15

Cacicus, 94
Cacique, 94
Cairina, 18
Calamospiza, 106
Calcarius, 108
Calidris, 30
Callipepla, 25
Calliste, 98
Calocitta, 74
Calospiza, 98
Calothorax, 51
Calypte, 51
Campethophilus, 57
Camptonotoma, 72
Campylopterus, 46
Campylorhamphus, 59
Campylorhynchus, 77
Canvasback, 19
Capella, 30
Capito, 54
Capitonidae, 54
Caprimulgidae, 43
Caprimulgus, 43
Capsiempis, 71
Caracara, 23, 23
Cardellina, 92
Cardinal, 101
Carpodacus, 102
Carpodectes, 64
Caryothraustes, 101
Casmerodius, 15
Cassiculus, 94
Cassidix, 95
Catbird, 80
Catharacta, 32
Cathartes, 20
Cathartidae, 20

Catharus, 82
Catherpes, 79
Catoptrophorus, 30
Celeus, 56
Centurus, 56
Cephalopterus, 65
Cercomacra, 62
Cerorhinca, 35
Certhia, 76
Certhiidae, 76
Ceryle, 53
Chachalaca, 24
Chaetura, 44
Chaeturella, 45
Chalybura, 49
Chamaea, 76
Chamaeidae, 76
Chamaepetes, 24
Chamaethlypis, 92
Charadriidae, 28
Charadrius, 29
Chat, 92
Chen, 17
Chickadee, 75
Chiroxiphia, 63
Chlidonias, 33
Chloroceryle, 53
Chlorophanes, 87
Chlorophonia, 97
Chloropipo, 63
Chlorospingus, 101
Chlorostilbon, 47
Chlothraupis, 100
Chlorura, 105
Chondestes, 106
Chondrohierax, 20
Chordeiles, 43
Chrysoptilus, 55
Chrysotlypis, 100
Chuck-will's-widow, 43
Cicaba, 42
Ciconiidae, 16
Cinclidae, 76
Cinclus, 76
Circus, 23
Cissiophila, 75
Cistothorus, 77
Claravis, 37
Cnemidectes, 70
Coccyzus, 40

Cochleariidae, 16
Cochlearius, 16
Coereba, 88
Coerebidae, 87
Colaptes, 55
Colibri, 46
Colinus, 25
Colonia, 66
Columba, 36
Columbidae, 36
Columbigallina, 37
Colymbidae, 9
Colymbus, 9
Comb-Duck, 18
Compsothlypis, 89
Condor, 20
Conebill, 88
Conirostrum, 88
Contopus, 68
Coot, 27
Coquette, 47
Coragyps, 20
Corapipo, 64
Cormorant, 14
Corvidae, 74
Corvus, 74
Coryphotriccus, 67
Cotinga, 64
Cotingidae, 64
Coturnicops, 27
Cowbird, 94
Cracidae, 24
Crake, 27
Crane, 26
Crane-Hawk, 23
Cranioleuca, 59
Craspedoprion, 70
Crax, 24
Creagrus, 33
Creeper, 76
Crocethia, 30
Crossbill, 104
Crotaphaga, 40
Crow, 74
Crypturellus, 9
Cuckoo, 40
Cuculidae, 40
Curassow, 24
Curlew, 29
Cyanerpes, 87

- Cyanocitta, 75
Cyanocompsa, 102
Cyanocorax, 74
Cyanolyca, 75
Cyclarhidae, 85
Cyclarhis, 85
Cygnus, 17
Cymbilaimus, 61
Cynanthus, 47
Cyphorhinus, 79
Cypseloides, 45
Cyrtonyx, 26

Dacnis, 87
Dactylortyx, 26
Damophila, 47
Daptrius, 23
Deconychura, 58
Deltarhynchus, 68
Dendrocincla, 58
Dendrocolaptes, 58
Dendrocolaptidae, 58
Dendrocopos, 57
Dendrocygna, 17
Dendroica, 89
Dendroplex, 58
Dendroptyx, 25
Dichromanaassa, 15
Dickcissel, 102
Diglossa, 87
Diomedea, 10
Diomedidae, 10
Dipper, 76
Dives, 95
Dolichonyx, 97
Donacobius, 81
Doricha, 50
Doryfera, 45
Dove, 36
Dowitcher, 30
Dromococcyx, 40
Drycopus, 55
Duck, 17
Dumetella, 80
Dunlin, 31
Dysithamnus, 61

Eagle, 22
Eagle-Hawk, 22
Ectopistes, 36

Egret, 15
Egretta, 15
Elaenia, 71
Elainopsis, 71
Elanoides, 20
Elanus, 20
Electron, 53
Elvira, 49
Emberizoides, 108
Emerald, 47, 48, 49
Empidonax, 68
Endomychura, 35
Erator, 65
Eremophila, 73
Ereunetes, 30
Ergaticus, 93
Erolia, 31
Eubucco, 53
Eucometis, 100
Eudocimus, 16
Eugenes, 50
Eumomota, 53
Eupagus, 95
Euperusa, 49
Euphonia, 97
Eupoda, 29
Euptilotis, 52
Eurypyga, 28
Eurypygidae, 28
Euthlypis, 93
Eutoxeres, 46

Falco, 24
Falcon, 23
Falconidae, 23
Finch, 101, 102-5
Finfoot, 28
Flamingo, 17
Flatbill, 70
Flicker, 55
Florida, 15
Florisuga, 46
Flower-piercer, 87
Fluvicola, 66
Flycatcher, 65
Foliage-gleaner, 60
Forest-Falcon, 23
Formicariidae, 61
Formicarius, 62
Formicivora, 62

Forpus, 39
Fregata, 14
Fregatidae, 14
Frigatebird, 14
Fringillidae, 101
Fruitcrow, 65
Fulica, 27
Fulmar, 10
Fulmarus, 10
Furnariidae, 59

Gadwall, 18
Galbula, 53
Galbulidae, 53
Gallinago, 30
Gallinula, 27
Galinule, 27
Gmpsonyx, 20
Gannet, 13
Gavia, 9
Gaviidae, 9
Gelochelidon, 34
Geococcyx, 40
Geothlypis, 91
Geotrygon, 37
Geranospiza, 23
Glaucidium, 42
Glaucis, 45
Glyphorhynchus, 58
Gnatcatcher, 83
Gnatwren, 84
Godwit, 29
Goethalsia, 48
Goldeneye, 19
Goldentail, 48
Goldfinch, 104
Goldmania, 48
Goosander, 19
Goose, 17
Goshawk, 21
Grackle, 94
Grallaria, 63
Grallaricula, 63
Granatellus, 92
Grass-Finch, 104
Grassquit, 102, 103
Grebe, 9
Greenlet, 87
Grosbeak, 101
Ground-Cuckoo, 40

- Ground-Dove**, 37
Ground-Finch, 108
Ground-Sparrow, 105
Gruidae, 26
Grus, 26
Guan, 24
Guiraca, 102
Gull, 32
Gygis, 35
Gymnocichla, 62
Gymnogyps, 20
Gymnopithys, 62
Gymnorhinus, 74
Gymnostinops, 94

Habia, 100
Haematopodidae, 28
Haematopus, 28
Haliaeetus, 23

Halocyptena, 12
Haplophaedia, 50
Harpagus, 20
Harpia, 22
Harpyhaliaetus, 22
Harrier, 23
Hawk, 20, 21
Hawk-Eagle, 22
Heleodytes, 77
Heliodoxa, 50
Heliomaster, 50
Heliornis, 28
Heliornithidae, 28
Heliothrix, 50
Helmitheros, 88
Hemithraupis, 100
Henicorhina, 79
Hermit, 45
Heron, 14
Herpetotheres, 23
Herpsilochmus, 62
Hesperiphona, 102
Heterocnus, 15
Heteroscelus, 30
Heterospingus, 100
Heterospizias, 21
Himantopus, 31
Hirundinidae, 73
Hirundo, 73
Honeycreeper, 87

Hoploxypterus, 28
House-Wren, 79
Hummingbird, 45
Hydranassa, 15
Hydrobatidae, 11
Hydroprogne, 34
Hylocharis, 47
Hylocichla, 82
Hyloctistes, 60
Hylomanes, 53
Hylophilax, 62
Hylophylus, 87
Hylorchilus, 79
Hypomorphnus, 22

Ibis, 16
Ibycter, 23
Icteria, 92
Icteridae, 94
Icterus, 95
Ictinia, 20
Iridoprocne, 74
Ixobrychus, 15
Ixoreus, 81

Jabiru, 16
Jacamar, 53
Jacamerops, 53
Jacana, 28
Jacanidae, 28
Jacobin, 46
Jaeger, 32
Jay, 74
Juliamyia, 47
Junco, 107

Kestrel, 24
Killdeer, 29
Kingbird, 66
Kingfisher, 53
Kinglet, 83
Kiskadee, 67
Kite, 20
Kittiwake, 33
Klais, 46
Knot, 30

Lampornis, 49
Lamprolaima, 50

Lancebill, 45
Laniidae, 85
Lanio, 100
Laniocera, 64
Lanius, 85
Lapwing, 28
Laridae, 32
Lark, 73
Larus, 32
Laterallus, 27
Lathria, 65
Leafscraper, 60
Legatus, 66
Leistes, 96
Lepidocolaptes, 59
Lepidopyga, 47
Leptodon, 20
Leptopogon, 72
Leptotila, 37
Leucolepis, 79
Leucophoyx, 15
Leucopternis, 22
Limnodromus, 30
Limnothlypis, 88
Limosa, 29
Limpkin, 26
Lipaugus, 65
Lobipes, 32
Lochmias, 61
Longspur, 108
Loon, 9
Lophodytes, 19
Lophornis, 47
Lophortyx, 25
Lophostryx, 41
Lophotriccus, 71
Loxia, 104
Eurocalis, 43
Lysurus, 105

Macaw, 38
Magpie-Jay, 74
Malacoptila, 54
Mallard, 18
Manacus, 64
Manakin, 63
Mango, 46
Mareca, 18
Margarornis, 59
Martin, 73

[124]

- Meadowlark, 96
Mecocerculus, 71
Megacyrle, 53
Megarynchus, 67
Melanerpes, 56
Melanitta, 19
Melanoptila, 80
Melanotis, 80
Meleagridae, 26
Meleagris, 26
Merlin, 24
Melospiza, 108
Melozone, 105
Merganser, 19
Mergus, 19
Mesembrinibis, 16
Micrathene, 42
Micrastur, 23
Microbates, 84
Microcerculus, 79
Microchera, 49
Micromonacha, 54
Micropalama, 31
Microrhopias, 62
Microtriccus, 72
Milvago, 23
Mimidae, 80
Mimodes, 80
Mimus, 80
Mionectes, 72
Mitrephanes, 69
Mitrospingus, 100
Mniotilta, 88
Mockingbird, 80
Mockingthrush, 81
Molothrus, 94
Momotidae, 53
Momotus, 53
Monasa, 54
Monklet, 54
Morococcyx, 40
Morphnus, 22
Morus, 13
Motacilla, 84
Motacillidae, 84
Motmot, 53
Mountain-gem, 49
Mourner, 64
Murrelet, 35
Muscivora, 64
Muscovy, 18
Myadestes, 81
Mycteria, 16
Myiarchus, 67
Myiobius, 69
Myioborus, 93
Myiochanes, 68
Myiodynastes, 66
Myiopagis, 71
Myiophobus, 70
Myiornis, 71
Myiozetetes, 67
Myrmeciza, 62
Myrmornis, 63
Myrmotherula, 61
Neochelidon, 73
Neochloe, 86
Neomorphus, 41
Nesotriccus, 68
Nighthawk, 43
Night-Heron, 15
Nightingale-Thrush, 82
Nightjar, 43
Noddy, 35
Nonnula, 54
Notharchus, 54
Nothocercus, 9
Notiochelidon, 73
Nucifraga, 74
Numenius, 29
Nunbird, 54
Nunlet, 54
Nutcracker, 74
Nuthatch, 76
Nuttallornis, 68
Nyctanassa, 15
Nyctibiidae, 43
Nyctibius, 43
Nycticorax, 15
Nyctidromus, 43
Nyctiphrynus, 43
Nystalus, 54
Oceanites, 11
Oceanodroma, 12
Odontophorus, 25
Odontorchis, 20
Oncostoma, 70
Onychorhynchus, 70
Oporornis, 91
Oreophasis, 24
Oreopyra, 50
Oreortyx, 25
Oreoscopetes, 81
Oriole, 95
Oriturus, 106
Ornithion, 72
Oropendola, 94
Ortalis, 24
Oryzoborus, 103
Osprey, 23
Ostินops, 94
Otophanes, 43
Otus, 41
Ovenbird, 59, 91
Owl, 41
Oxyruncidae, 72
Oxyruncus, 72
Oxyura, 19
Oystercatcher, 28
Pachyramphus, 64
Pachysylvia, 87
Pandion, 23
Pandionidae, 23
Panyptila, 45
Panterpe, 47
Paphosia, 47
Parabuteo, 22
Parakeet, 38
Pardirallus, 27
Paridae, 75
Parrot, 38
Parrotlet, 39
Partridge, 25
Parula, 89
Parulidae, 88
Parus, 75
Passer, 94
Passeridae, 94
Passerculus, 106
Passerella, 108
Passerina, 102
Pauraque, 43
Pelagodroma, 12
Pelecanidae, 13
Pelecanus, 13
Pelican, 13
Penelope, 24

- Penelopina, 24
Peppershrike, 85
Perissotriccus, 71
Petrel, 10, 11
Petrochelidon, 73
Peucedramus, 89
Pewee, 68
Pezopetes, 104
Phaeochroa, 46
Phaeomyias, 72
Phaenostictus, 62
Phaeoprogne, 73
Phaethon, 12
Phaethontidae, 12
Phaethornis, 45
Phainopepla, 84
Phainoptila, 85
Phalacrocoracidae, 14
Phalacrocorax, 14
Phalaenoptilus, 43
Phalarope, 31
Phalaropodidae, 31
Phalaropus, 31
Pharomachrus, 52
Phasianidae, 25
Pheucticus, 101
Philodice, 50
Philortyx, 25
Philydor, 60
Phloeoceastes, 57
Phlogothraupis, 99
Phoebe, 65
Phoenicopteridae, 17
Phoenicopterus, 17
Phyllomyias, 72
Phylloscartes, 71
Piaya, 40
Picidae, 55
Piculet, 55
Piculus, 55
Picumnus, 55
Pigeon, 36
Piha, 65
Filherodius, 15
Pinaroloxias, 102
Pintail, 18
Pionopsitta, 39
Pionus, 39
Pipilo, 105
Pipit, 84
Pipra, 63
Pipridae, 63
Piprites, 63
Pipromorpha, 72
Piranga, 99
Pitangus, 67
Pittasoma, 63
Pitylus, 101
Plagiospiza, 106
Platyparis, 65
Platyrinchus, 70
Plegadis, 16
Ploceidae, 94
Plover, 28
Plumeleteer, 49
Pluvialis, 28
Podiceps, 9
Podicipididae, 9
Podilymbus, 10
Polioptila, 83
Polioptilidae, 83
Polyborus, 23
Pooecetes, 106
Poorwill, 43
Popelairia, 47
Porphyruia, 27
Porzana, 27
Potoo, 43
Praedo, 69
Premnoplex, 59
Priocella, 10
Procellariidae, 10
Procnias, 65
Progne, 73
Protonotaria, 88
Psaltiparus, 76
Psarocolius, 94
Pselliophorus, 104
Pseudocolaptes, 60
Pesudotriccus, 71
Psilorhinus, 74
Psittacidae, 38
Psomocolax, 94
Pterodroma, 11
Pteroglossus, 55
Ptilogonatidae, 84
Ptilogonyx, 84
Ptychoramphus, 35
Puffbird, 54
Puffinus, 10
Puffleg, 50
Pulsatrix, 42
Pygmy-Owl, 42
Pygmy-Tyrant, 71
Pygochelidon, 73
Pyrocephalus, 66
Pyrrhuloxia, 101
Pyrrhura, 39
Quail, 25
Quail-Dove, 37
Querula, 65
Quetzal, 52
Quiscalus, 94
Rail, 26
Rallidae, 26
Rallus, 26
Ramphastidae, 55
Ramphastos, 55
Ramphocaenus, 84
Ramphecelus, 99
Raven, 74
Recurvirostra, 31
Recurvirostridae, 31
Redhead, 19
Redstart, 92
Regulidae, 84
Regulus, 84
Rhinocryptidae, 63
Rhinoptynx, 42
Rhodinocichla, 100
Rhodothraupis, 101
Rhynchocyclus, 70
Rhynchophanes, 108
Rhynchopsitta, 38
Rhopoterpe, 63
Rhynchorhynchus, 26
Rhytipterna, 65
Richmondena, 101
Ridgwayia, 83
Riparia, 73
Rissa, 33
Roadrunner, 40
Robin, 81
Rostrhamus, 20
Royal-Flycatcher, 70
Rynchopidae, 35
Rynchops, 35

- Sabrewing, 46
 Salpinctes, 79
 Saltator, 101
 Sanderling, 30
 Sandpiper, 29
 Sapayo, 64
 Sapsucker, 57
 Sarcoramphus, 20
 Sarkidiornis, 18
 Sayornis, 65
 Scaphidura, 94
 Scardafella, 37
 Scaup, 19
 Schiffornis, 64
 Sclerurus, 60
 Scolopacidae, 29
 Scops-Owl, 41
 Scoter, 19
 Scotothorus, 64
 Screech-Owl, 41
 Scytalopus, 63
 Scythebill, 59
 Seedeater, 103
 Seed-Finch, 103
 Seiurus, 91
 Selasphorus, 51
 Selenidera, 55
 Semnornis, 54
 Serpophaga, 71
 Setophaga, 92
 Sharpbill, 72
 Sheartail, 50
 Shearwater, 10
 Shoveler, 18
 Shrike, 85
 Shrike-Tanager, 100
 Shrike-Vireo, 85
 Sialia, 83
 Sicalis, 104
 Sicklebill, 46, 59
 Silky-flycatcher, 84
 Sirystes, 66
 Siskin, 103
 Sitta, 76
 Sittasomus, 58
 Sittidae, 76
 Skimmer, 35
 Skua, 32
 Smaragdolanius, 85
 Snipe, 30
 Snowcap, 49
 Softtail, 59
 Solitaire, 81
 Sora, 27
 Spadebill, 70
 Sparrow, 94, 105, 106
 Spatula, 18
 Speotyto, 42
 Sphyrapicus, 57
 Spindalis, 99
 Spinetail, 59
 Spinus, 103
 Spiza, 102
 Spizaetus, 23
 Spizastur, 22
 Spizella, 107
 Spodiornis, 104
 Spoonbill, 16
 Sporophila, 103
 Squatarola, 28
 Starling, 85
 Starthroat, 50
 Steganopus, 31
 Stelgidopteryx, 73
 Stellula, 51
 Stercorariidae, 32
 Stercorarius, 32
 Sterna, 34
 Stilt, 31
 Stork, 16
 Storm-Petrels, 11
 Streamcreeper, 61
 Streptoprocne, 44
 Strigidae, 41
 Strix, 42
 Sturnella, 96
 Sturnidae, 85
 Sturnus, 85
 Sublegatus, 72
 Sula, 13
 Sulidae, 13
 Sunbittern, 28
 Sungrebe, 28
 Surfbird, 30
 Swallow, 73
 Swallow-Tanager, 97
 Swan, 17
 Swift, 44
 Sylviidae, 83
 Synallaxis, 59
 Syndactila, 60
 Synthliboramphus, 35
 Tachycineta, 74
 Tachyphonus, 100
 Tanager, 97
 Tanagra, 97
 Tangara, 98
 Tangavius, 94
 Tapaculo, 63
 Tapera, 40
 Taraba, 61
 Tattler, 30
 Teal, 18
 Telmatodytes, 77
 Terenotriccus, 69
 Terenura, 62
 Tern, 33
 Tersina, 97
 Thalasseus, 34
 Thalurania, 47
 Thamnistes, 61
 Thamnophilus, 61
 Theristicus, 16
 Thick-knee, 32
 Thorntail, 47
 Thrasher, 80
 Thraupidae, 97
 Thraupis, 99
 Threnetes, 45
 Threskiornithidae, 16
 Thripadectes, 60
 Thrush, 81
 Thrush-Tanager, 100
 Thryomanes, 78
 Thryorchilus, 79
 Thryothorus, 77
 Tiaris, 102
 Tiger-Heron, 15
 Tigrisoma, 15
 Tilmatura, 51
 Timaliidae, 76
 Tinamidae, 9
 Tinamou, 9
 Tinamus, 9
 Titmouse, 76
 Tityra, 65
 Todirostrum, 70
 Tody-Flycatcher, 70
 Tolmomyias, 70

- Totanus**, 30
Toucan, 55
Toucanet, 55
Touit, 39
Towhee, 105
Toxostoma, 80
Tree-Duck, 17
Treehunter, 60
Treerunner, 59
Tringa, 30
Tripsurus, 57
Trochilidae, 45
Troglodytes, 79
Troglodytidae, 77
Trogon, 52
Trogonidae, 52
Tropicbird, 12
Tryngites, 31
Tuftedcheek, 60
Turdidae, 81
Turdus, 81
Turkey, 26
Turnstone, 30
Tyrannidae, 65
Tyranniscus, 72
Tyrannulet, 71
Tyrannulus, 72
Tyrannus, 66
Tyrant, 66
Tyto, 41
Tytonidae, 41
Umbrellabird, 65
Uropsila, 79
Urubitinga, 22
Urubitornis, 22
Veery, 82
Veniliornis, 57
Verdin, 76
Vermivora, 88
Violet-ear, 46
Vireo, 85
Vireolaniidae, 85
Vireolanius, 85
Vireonidae, 85
Volatinia, 103
Vulture, 20
Wagtail, 84
Warbler, 83, 88
Waterthrush, 91
Water-Tyrant, 66
Waxwing, 84
Wedge-bill, 58
Whimbrel, 29
Whip-poor-will, 43
Widgeon, 18
Willet, 30
Wilsonia, 92
Woodcreeper, 57
Woodhaunter, 60
Woodhewer, 57
Woodnymph, 47
Wood-Partridge, 25
Woodpecker, 55
Wood-Pewee, 68
Wood-Quail, 25
Wood-Rail, 27
Woodstar, 50, 51
Wood-Warbler, 88
Wood-Wren, 79
Wren, 77
Wren-Thrush, 83
Xanthocephalus, 96
Xanthornus, 94
Xanthoura, 75
Xema, 33
Xenerpestes, 59
Xenicopsis, 60
Xenicopoides, 60
Xenocistics, 60
Xenops, 60
Xenornis, 61
Xenospiza, 106
Xenotriccus, 69
Xiphocolaptes, 58
Xiphorhynchus, 58
Yellowlegs, 30
Yellowthroat, 91
Zarhynchus, 94
Zeledonia, 83
Zeledoniidae, 83
Zenaida, 36
Zenaidura, 36
Zonotrichia, 108

The species of Middle American birds. A list of all species recorded from Mexico to Panama, with suggested English names, outlines of range, and a distributional bibliography. By Eugene Eisenmann, with the collaboration in the selection of English names of Emmet R. Blake and Edward L. Chalif.

Eisenmann, Eugene, 1906-1981.

[New York, The Linnaean Society of New York] 1955.

<http://hdl.handle.net/2027/coo.31924000092613>

HathiTrust



www.hathitrust.org

Public Domain, Google-digitized

http://www.hathitrust.org/access_use#pd-google

We have determined this work to be in the public domain, meaning that it is not subject to copyright. Users are free to copy, use, and redistribute the work in part or in whole. It is possible that current copyright holders, heirs or the estate of the authors of individual portions of the work, such as illustrations or photographs, assert copyrights over these portions. Depending on the nature of subsequent use that is made, additional rights may need to be obtained independently of anything we can address. The digital images and OCR of this work were produced by Google, Inc. (indicated by a watermark on each page in the PageTurner). Google requests that the images and OCR not be re-hosted, redistributed or used commercially. The images are provided for educational, scholarly, non-commercial purposes.

**LIBRARY
ANNEX
M2**

A. R. MANN
LIBRARY
CORNELL U.
Digitized by Google

Original from
CORNELL UNIVERSITY

**ALBERT R. MANN
LIBRARY**

**NEW YORK STATE COLLEGES
OF
AGRICULTURE AND HOME ECONOMICS**



**AT
CORNELL UNIVERSITY**

Date Due

JAN 16 1968
JAN 16 1968
JAN 12 1968
JAN 12 1970
DEC 14 1972
MAY 14 1973
MAY 16 1974
MAY 18 1974
DEC 11 1975 0

Library Bureau Cat. No. 1137

Cornell University Library
QL 687.E36

The species of Middle American birds. A I



3 1924 000 092 613